

NEGATIVE TRENDS IN BIRD ABUNDANCE ARE STRONGLY CORRELATED TO RAINFALL DECLINE IN A CENTRAL AMERICAN TROPICAL FOREST

ALEXIS CEREZO BLANDÓN^{1,3}, CHANDLER S. ROBBINS², BARBARA DOWELL², MIGUEL RAMÍREZ³, ANTONIO LÓPEZ³ AND OBDULIO JAVIER³

¹*Fundación para la Conservación y el Ecodesarrollo (FUNDAECO). 25 calle 2-39, Zona 1, 0101 Ciudad de Guatemala, Guatemala. a.cerezo@fundaeco.org.gt*

²*Patuxent Wildlife Research Center, Biological Resources Division, United States Geological Survey, 11410 American Holly Drive, 20708-4015 Laurel, Maryland, USA.*

³*Aves Argentinas/Asociación Ornitológica del Plata. Matheu 1246, Buenos Aires, Argentina.*

ABSTRACT.— Climate change is considered one of the main factors threatening biodiversity. Weather is of major importance for bird population dynamics, but the implications of climate change have only recently begun to be addressed, especially for tropical birds. For Northern Central America, climate change scenarios for 2050 predict a reduction in precipitation across the region, with decreases ranging from 4–19% of current rainfall. In this work, we addressed the relationship between temporal changes in precipitation amount and bird community dynamics in eastern tropical Guatemala, for a time period of 18 years (1993–2010). Data consisted of yearly captures and recaptures in four sites, located at elevations between 100–750 masl, and analyses were carried out for total captures and for six foraging guilds. Statistical analyses consisted of Poisson regressions, where estimated abundance (taking into account recapture probability) was modelled as a function of wet-season, dry-season, and annual rainfall, and temporal trend. We detected strong declines in total abundance and in the abundance of nectarivores, omnivores, frugivores, and, to a lesser degree, foliage insectivores. These declines were strongly associated with declines in rainfall amount, generally during the rainy season. A more comprehensive understanding of the effects of climate change on animal abundance in tropical ecosystems is strongly needed to propose conservation and management actions in these biodiverse ecosystems.

KEY WORDS: *capture probability, climate change, community dynamics, feeding guilds, Guatemala, Poisson regression, tropical birds.*

RESUMEN. LAS TENDENCIAS NEGATIVAS EN LA ABUNDANCIA DE AVES ESTÁN FUERTEMENTE CORRELACIONADAS CON LA DISMINUCIÓN DE LAS PRECIPITACIONES EN UN BOSQUE TROPICAL CENTROAMERICANO.— El cambio climático es considerado uno de los principales factores que amenazan a la biodiversidad. El clima es de gran importancia para la dinámica poblacional de las aves, pero las consecuencias del cambio climático no han sido abordadas sino hasta recientemente, especialmente en aves tropicales. Para el norte de América Central, los escenarios de cambio climático para 2050 predicen una reducción en las precipitaciones, con disminuciones de 4–19% de las precipitaciones actuales. En este trabajo se evalúa la relación entre los cambios temporales en la cantidad de precipitaciones y la dinámica de la comunidad de aves en el este tropical de Guatemala para un período de 18 años (1993–2010). Los datos consistieron en capturas y recapturas anuales en cuatro sitios ubicados a altitudes entre 100–750 msnm, con análisis para las capturas totales y para seis gremios de alimentación. Los análisis estadísticos consistieron en regresiones de Poisson en las cuales la abundancia estimada (tomando en cuenta la probabilidad de recaptura) fue modelada en función de las precipitaciones (de la época lluviosa, de la época seca y la anual) y la tendencia temporal. Se detectaron fuertes declinaciones en la abundancia total y en la abundancia de las aves nectarívoras, omnívoras, frugívoras y, en menor grado, insectívoras de follaje. Estas declinaciones estuvieron fuertemente asociadas a la disminución en la cantidad de precipitaciones, principalmente de la época lluviosa. Se necesita un conocimiento más profundo de los efectos del cambio climático sobre la abundancia de animales en los ecosistemas tropicales, con el fin de proponer acciones de conservación y manejo en estos sistemas altamente diversos.

PALABRAS CLAVE: *aves tropicales, cambio climático, dinámica de comunidades, gremios de alimentación, Guatemala, probabilidad de captura, regresión de Poisson.*

Climate change is considered one of the main factors threatening biodiversity (Root and Schneider 2002, Walther et al. 2002, Chen et al. 2004, Thomas et al. 2004) because species strongly depend on interannual fluctuations in precipitation, temperature, and extreme climatic events over ecological and evolutionary time scales (Chen et al. 2004, Parmesan 2006, Beever et al. 2011). Weather is of major importance for the population dynamics of bird populations, and there has thus been a recent, rising need to address the implications of climate change on these populations (Crick 2004, Wolfe and Ralph 2009, Sekercioglu et al. 2012, Faaborg et al. 2013, Foden et al. 2013, Blake and Loiselle 2015). Under the Millennium Ecosystem Assessment scenarios, Jetz et al. (2007) predicted that even under environmentally benign scenarios, at least 400 bird species are projected to suffer >50% range reductions by the year 2050. Also, multiple models based on the Intergovernmental Panel on Climate Change (IPCC) A2 Emissions Scenario predict tropical drying trends, particularly in the Caribbean and Central America region and equatorial South America (Neelin et al. 2006).

Climate-driven changes in abundance, phenology and changes in species range, among others, have been shown by recent studies, mainly for northern temperate latitudes, in general associated with rising temperatures (Root and Schneider 2002, Walther et al. 2002, Gordo and Sanz 2006, Parmesan 2006). However, due to the scarcity of long-term data sets in the tropics, biotic changes are relatively undocumented in the region (Harris et al. 2011), with some notable exceptions: in particular, a long-term study involving captures and observations in undisturbed lowland tropical forest, found that overall captures and observations declined by approximately 40 and 50%, respectively, from 2008 to 2014 (Blake and Loiselle 2015). Others have studied the effects of climate, mainly El Niño–Southern Oscillation (ENSO) effects, on tropical bird communities (Wolfe and Ralph 2009, Styrsky and Brawn 2011, Wolfe et al. 2015), but these studies have focused on one or a few bird species. Consequently, it is crucial to increase our basic knowledge in order to evaluate to what extent climate-driven changes have also occurred in the tropics, because: (1) climatic effects between biomes may be extremely dif-

ferent, and (2) we need to understand and eventually predict the impact of future climatic changes on all biotic communities and ecosystems, and particularly on the diverse tropical ecosystems.

Studies in northern latitudes have shown that insectivorous bird abundance and reproductive success are affected by insect food abundance, and both temperature and precipitation indirectly affect insect abundance and plant growth rates via leaf quality or budding (Sillett et al. 2000, Nott et al. 2002, Jones et al. 2003, Visser et al. 2004), suggesting that bird population dynamics are in turn indirectly driven by regional climate patterns, through their effects on plant and insect biomass. In the tropics, rainfall is the key factor determining the overall primary productivity and its temporal variability (Schloss et al. 1999), and it is positively associated with increases in plant and insect biomass (Schuur 2003, Cao et al. 2004, Cleland et al. 2007, Saatchi et al. 2007, for plants; Denlinger 1980, Lowman 1982, Frith and Frith 1985, Bonebrake et al. 2010, for insects). Likewise, the availability of food resources in Neotropical forests has been positively correlated with bird abundance for several feeding guilds, such as insectivores, frugivores, and nectarivores (Karr and Brawn 1990, Loiselle and Blake 1991, Poulin et al. 1992). Also, plant phenological cycles, in particular leaf flushing, fruiting and flowering periods, are associated (to different degrees) to the onset or end of the rainy season in the tropics (Frankie et al. 1974, Wolda 1978, van Schaik et al. 1993, Cattanio et al. 2004), and other studies from tropical sites have found that insect abundance was positively related to the peak of plant phenological activities (Wolda 1978, Lowman 1982). Because of these relationships between rainfall and plant and insect productivity, a positive association between the amount of rainfall during the rainy season and bird abundance could be expected (Powell et al. 2015). Alternatively, dry seasons are potential periods of scarce resources, resulting in “ecological bottlenecks” that limit species abundances and biotic interactions and processes, and may be important in structuring bird communities in variable ecosystems (Williams and Middleton 2008). Finally, rainfall in both seasons might have independent effects on tropical bird populations, because different demographic

components that determine abundance might depend on either dry-season or wet-season rainfall. Thus, abundance might also be associated with total annual rainfall.

In this work, we addressed the relationship between temporal changes in precipitation and bird community dynamics in eastern tropical Guatemala, using mist-net data from the Bird Monitoring Program of the Caribbean Region of Guatemala, established in 1992 (Cerezo et al. 2012). We evaluated changes in total bird abundance and in the abundance of six foraging guilds, in relation to rainfall amount in the 1993–2010 period (18 years). Specific objectives were: (1) to evaluate if abundance was associated (positively or negatively) to the amount of rainfall in the previous year, either to wet-season rainfall, dry-season rainfall, a combination of these two, or to annual rainfall, and (2) to evaluate the existence of an increasing or decreasing trend in total abundance and of each feeding guild, that was independent of any potential relationship to the amount of rainfall.

METHODS

Study site, bird sampling, and rainfall data

The study was carried out in Cerro San Gil Watershed Protection Reserve, located in eastern tropical Guatemala (Fig. 1). The reserve is divided into three zones: a nuclear zone (full preservation zone, where only biological research and low-impact tourism activities are allowed), a buffer zone (its main objective is to mitigate possible impacts of surrounding

human populations to the nuclear zone; only environmentally sustainable activities are allowed, e.g., reforestation, small-scale farming in established communities), and a multiple-use zone (extant economic activities are allowed, e.g., farming, cattle ranching, commercial plantations, as well as subsistence hunting and selective wood extraction). Large-scale deforestation is not allowed in either the multiple-use or buffer zones. The dominant vegetation type is tropical wet forest, with mean (\pm SD) annual temperature and precipitation of 26.6 ± 0.4 °C and 3284.9 ± 596.5 mm, respectively, for the study period (INSIVUMEH 2012).

Bird sampling was carried out in four sites, located approximately at 100, 200, 250 and 750 masl. All sites were located within the nuclear zone, in relatively undisturbed forest (i.e., with minimum human disturbance), and there have thus been no major habitat changes either in the immediate area or in the broader landscape. The 100 masl site was approximately 1 km away from sites at 200 and 250 masl, these two were very close together, approximately 200 m from each other, and the 750 masl site was approximately 3 km away from the 200 and 250 masl sites. Sampling consists of yearly constant-effort mist netting in each site throughout the duration of the study, between early February and mid-March. In each site, 16 mist nets (36 mm mesh size) were placed at the same net location every year, at an average distance of approximately 50 m from each other. Mist nets were only occasionally shifted a few meters as a result of tree falls. The understory bird community was sampled during three consecutive days, usually between 06:00–16:00 h, for an average (\pm SD) of 430.0 ± 60.6 net hours per site each year. During sampling periods, nets were occasionally closed because of heavy rain or human disturbances.

All captured birds were identified to the species level, and were banded with a serially-numbered aluminium ring, or note was taken of the ring number if it was a recaptured individual (i.e., a bird banded in a previous year to the current sampling period). For all individuals, we determined sex and age (if possible) with criteria described in Wolfe et al. (2009), measured wing length and weight, and took notes on moult and plumage condition. The Bander's Code of Ethics (North Ameri-




Figure 1. Study site (barred rectangle) located within Cerro San Gil Watershed Protection Reserve, eastern Guatemala. Grey shaded area is tropical wet forest.

can Bird Banding Council 2001) was strictly followed during sampling. For the present analysis, we used data derived from the number of captured individuals per species. Taxonomy of birds is according to the North American Classification Committee checklist (American Ornithological Society 2016).

Precipitation data was taken from the nearest meteorological station of the Guatemalan Institute of Seismology, Vulcanology, Meteorology and Hydrology (INSIVUMEH 2012), located in the city of Puerto Barrios, approximately 3 km from the nearest of the four study sites. Other studies have evaluated the ENSO effects on different parameters of tropical bird communities and populations (Wolfe and Ralph 2009, Blake and Loiselle 2015, Wolfe et al. 2015). In the tropics, ENSO typically alters precipitation patterns (Holmgren et al. 2001, Mahli and Wright 2004), so we expected that our rainfall data would reflect abnormal rainfall conditions caused by ENSO, as well as the effects of such unusual conditions on bird abundance.

Statistical analysis

For statistical analyses, we only included species that had more than five captured individuals in at least three years. We considered that there would be a potential for bias in the results if these individuals were included in the analysis. We also excluded Nearctic migrants, because their population dynamics can be altered by other factors outside the wintering grounds. For species with captures in at least 9 of 18 years and more than 30 captures, we calculated a Pearson correlation coefficient as a descriptive measure of association between “year” and observed abundance (following Blake and Loiselle 2015), a negative association indicating a temporal decline in captures.

Because observed captures (i.e., captures that do not take into account capture or recapture probability) may provide biased counts of bird abundance (Pollock et al. 2002, Williams et al. 2002), we estimated yearly bird abundance using the estimator developed by Pollock and Otto (1983), using each one of the three sampling days per site as repeated sampling periods, which are necessary for the estimation process. Pollock and Otto’s (1983) abundance estimator assumes heterogeneity among individuals in capture probability, and

different capture probabilities for first and subsequent captures during the multiple sampling periods. Figure 2 provides graphs of capture and recapture rates, showing that the observed pattern of captures and recaptures conform to the assumptions of this model for estimating abundance (i.e., different capture and recapture probabilities). Estimated yearly abundance data were then expressed as capture rates per 100 net hours (net hours were transformed to vary between 0 and 1).

To analyse the relationship between bird abundance, rainfall variables and temporal trends in populations, we used Poisson regression, a special type of generalized linear model appropriate for the discrete, highly-skewed distribution of dependent variables that are counts (Vincent and Haworth 1983). Abundance was modelled as a function of: (1) total rainfall for the seven-month period prior to captures (July–January) comprising the rainy season and accounting on average for approximately 66% of total yearly rainfall, (2) total rainfall for the five-month period prior to cap-




Figure 2. Decreasing proportion of captures with sampling day (a), and proportion of birds captured in only one day during each sample period (“One capture”), or recaptured in the second or third sampling day (“Recaptured”) (b) in Cerro San Gil Watershed Protection Reserve, eastern Guatemala.

Table 1. Description of predictive variables in fitted models of bird abundance in Cerro San Gil Watershed Protection Reserve, eastern Guatemala.

Model	Predictive variables
Wet + Dry + Trend	Wet- and dry-season rainfall, and temporal trend
Wet + Trend	Wet-season rainfall and temporal trend
Dry + Trend	Dry-season rainfall and temporal trend
Annual + Trend	Annual rainfall and temporal trend
Trend	Temporal trend

tures (February–June) comprising the dry season and accounting on average for approximately 31% of total annual rainfall, (3) annual rainfall (the sum of wet- and dry-season rainfall), and (4) year or temporal trend. Poisson regression analyses were carried out for total abundance and for the abundance of six foraging guilds: frugivores, nectarivores, sallying insectivores, bark insectivores, foliage (gleaning/leaf-tossing) insectivores, and omnivores. Foraging guild classification was made according to Stiles and Skutch (1989) and Terborgh et al. (1990).

Models were compared using an information-theoretic framework. For total abundance and abundance of each foraging guild, we fitted five different models, with different

combinations of wet-season, dry-season and annual rainfall, and temporal trend (Table 1). The second-order Akaike Information Criterion (AIC_c , recommended when $n/K < 40$, where n is sample size and K is the number of estimated parameters) and Akaike weights (w_i) were then used to choose the best-fitting models from the set of candidate models (Burnham and Anderson, 1998, 2001, Anderson et al. 2000). When differences between AIC values are small ($< 2 AIC$ units), Akaike weights can be used as indicators of the strength of evidence for each model. The w_i is interpreted as the approximate probability that model i is the best model in the set of models being considered (Anderson et al. 2000). To compare the relative effects of the




Figure 3. Plots of temporal autocorrelation function for residuals corresponding to fitted models of bird abundance in Cerro San Gil Watershed Protection Reserve, eastern Guatemala. Plots were produced for models in confidence sets, according to the Akaike weight.

three rainfall variables and linear trend, we used multi-model inference, or model averaging (Burnham and Anderson 1998). For each species group we obtained mean model coefficients for each effect, weighted by w_i .

Because rainfall variables were positively correlated, and rainfall variables and temporal trend were negatively correlated (to differing degrees; see *Results*), we used sequential regression (Graham 2003, Dormann et al. 2013) to avoid problems associated with multicollinearity in the estimation of regression parameters. Sequential regression consists in linearly regressing pairs of explanatory variables against each other, and then extracting the residuals for the regressed variable, thus producing a new explanatory variable that expresses variation that is completely independent of the other predictors (Dormann et al. 2013). In our study, we regressed “year” against “rainfall amount” to produce measures of temporal trend that were independent of rainfall amount. All variables were transformed to vary between 0 and 1, so that estimated model coefficients could be compared directly.

Finally, because we modelled a time series, we were possibly violating the statistical assumption of independence between model residuals, which can inflate type-I error rates (Zuur et al. 2009). For each model, we produced a plot of the temporal autocorrelation function for residuals, using the autocorrelation function (“acf”) in R software (R Core Team 2014). The plot consists of a graph of the autocorrelation function of residuals for different time lags (in this case, of one year), and the horizontal dashed lines in graphs are lower and upper 95% confidence limits

(Fig. 3). Values that fall within these limits correspond to the statistical hypothesis of no temporal autocorrelation in residuals (Zuur et al. 2009). Models did not show or showed very little autocorrelation (Fig. 3), and we proceeded with model formulations without autocorrelation structures.

RESULTS

In our study site, rainfall significantly decreased during the study period (Fig. 4). Estimated declines in rainfall were of -38.8 mm/year (-0.11 mm/day per year) for wet-season rainfall, -21.1 mm/year (-0.06 mm/day per year) for dry-season rainfall, and -59.9 mm/year (-0.16 mm/day per year) for annual rainfall. The declines were more evident for wet-season rainfall ($r = 0.55$; Fig. 2). The declines are much higher than those reported in other studies for tropical regions. Malhi and Wright (2004) reported declines in the tropics worldwide during the 1960–1998 period, with an estimated mean decrease of 22 mm/year (0.06 mm/day per year). Declines in rainfall have also been reported for more recent periods for Central America (-0.02 mm/day per year; IPCC 2014). In particular, Neelin et al. (2006) and Rauscher et al. (2008) have reported a significant drying trend in the Central America-Caribbean region. Neelin et al. (2006) report a 5–30% decline in rainfall from the mean values in 1979, and also report expected declines derived from climate models for the region of -0.005 to -0.01 mm/day per year in the next 100 years.

In total, 53 species were included in analyses (Table 2). Six species were classified as frugivores, 9 as nectarivores, 12 as sallying




Figure 4. Temporal trends in wet-season, dry-season, and annual rainfall during the study period in Cerro San Gil Watershed Protection Reserve, eastern Guatemala. Pearson correlation coefficients between year and rainfall are shown.

Table 2. Bird species recorded in Cerro San Gil Watershed Protection Reserve (eastern Guatemala) included in the analysis. Foraging guild, total captures per year, Pearson correlation coefficient (r) as a descriptive measure of temporal change in captures, and number of individuals captured (with percentage between brackets) in each of the four sites (100, 200, 250 and 750 masl) are shown.

	Guild ^a	Year (1993–2010)												r	masl									
		93	94	95	96	97	98	99	00	01	02	03	04		05	06	07	08	09	10	100	200	250	750
<i>Leptotila cassinii</i>	F	5		3	4	2	4	3	4	3	1	4	2	4	5	7	3	3	2	16 (27)	22 (37)	13 (22)	8 (14)	
<i>Geothlypis montana</i>	F	9	2	9	5	5	12	4	16	2	5	13	9	10	12	6	3	10	2	40 (30)	18 (13)	69 (51)	7 (5)	
<i>Florisuga mellivora</i>	N	3	2			1	1	6				1	1			1	1	1	-	8 (44)	6 (33)	2 (11)	2 (11)	
<i>Threnetes ruckeri</i>	N	6	5	5	3	1	1	1	10	6	4	1	3	2	2	1			-0.54	3 (6)	19 (39)	25 (51)	2 (4)	
<i>Phaethornis longirostris</i>	N	86	55	59	94	31	49	56	66	84	54	41	40	47	62	41	14	22	25	312 (34)	199 (21)	293 (32)	122 (13)	
<i>Phaethornis striigularis</i>	N	4	2	1	3	0	1	1	2	2	4	0	1	1	2	2	1	1	-	5 (17)	9 (31)	12 (41)	3 (10)	
<i>Campylopterus hemileucurus</i>	N	104	39	9	76	10	57	30	17	45	14	11	6	9	14	13	1	11	-0.67	134 (29)	166 (36)	44 (9)	122 (26)	
<i>Eupherusa eximia</i>	N	2	3	2	2	6	1	3	4	4	2	1	2	2	2	3	1	2	1	-0.34	1 (2)	1 (2)	41 (95)	
<i>Thalurania colombica</i>	N	26	20	17	24	17	16	31	12	29	4	14	15	7	13	3	1	11	4	69 (26)	61 (23)	45 (17)	89 (34)	
<i>Anazilia candida</i>	N	8	10	9	11	2	9	11	3	7	3	6	3	3	1	2	1	3	3	-0.77	25 (27)	36 (39)	24 (26)	7 (8)
<i>Anazilia tzacatl</i>	N	1	1			1	2	4		2	2	2	1	3	5	1	1	1	-	16 (57)	9 (32)	3 (11)	-	
<i>Hylomanes monrotula</i>	SI	3						1	1	1			2	1		1		1	-	2 (20)	-	-	8 (80)	
<i>Monotus monota</i>	SI	1	2	1	1	2	2	2	2	2	2	2	2	1	1		2	1	-	8 (33)	5 (21)	8 (33)	3 (13)	
<i>Malacoptila panamensis</i>	SI	3		2	2	3			7	2	2	2	2	1	1	1	2	2	-	14 (52)	7 (26)	6 (22)	-	
<i>Picooides fumigatus</i>	BI				2			5					1						-	-	2 (25)	4 (50)	2 (25)	
<i>Dysithamnus mentalis</i>	FI	3	1	5	3	4	4	1	3	3	5	4	5	2	2	2	3	5	0.08	1 (2)	-	-	57 (98)	
<i>Myrmotherula schisticolor</i>	FI	11	7	11	4	7	3	4	5	4	6	5	8	3	4	4	8	4	3	-0.49	1 (1)	4 (4)	10 (10)	86 (85)
<i>Grallaria guatemalensis</i>	FI			1	2		2	1		2	1								-	1 (14)	5 (71)	-	1 (14)	
<i>Formicarius analis</i>	FI	4	4	6	6	7	6	2	5	3	7	2	4	4	4	2	1	2	-	16 (23)	12 (17)	14 (20)	27 (39)	
<i>Sclerurus guatemalensis</i>	FI	11	7	13	5	3	7	10	7	9	13	12	9	11	14	8	11	7	0.26	18 (11)	17 (10)	28 (17)	105 (63)	
<i>Sittasomus griseicapillus</i>	BI							1	1	2		2		1			1	2	-	3 (30)	4 (40)	3 (30)	-	
<i>Dendrocincla anabatina</i>	BI	30	23	13	11	32	23	21	14	28	20	12	38	16	18	12	11	14	8	-0.41	113 (33)	73 (21)	97 (28)	61 (18)
<i>Dendrocincla homochroa</i>	SI	12	11	3	1	8		1	2	11	1	5	2	2	1	5	3	6	7	-0.23	10 (12)	16 (20)	29 (36)	26 (32)
<i>Glyphorhynchus spirurus</i>	BI	43	32	34	19	27	47	31	35	22	22	34	28	20	21	27	18	19	-0.63	134 (27)	67 (14)	234 (47)	61 (12)	
<i>Dendrocolaptes sancithomae</i>	BI	2	5		1	1	2	2	4	2	2	2	2	3	3	2	2	1	1	-0.05	11 (33)	7 (21)	15 (45)	-
<i>Xiphorhynchus flavigaster</i>	BI	1	3	5	4		9	6	9	8	3	6	1	2	6	2	5	2	4	-0.04	29 (38)	18 (24)	29 (38)	-
<i>Xiphorhynchus erythropygius</i>	BI			2					1		1		1			2	1	2	-	-	-	-	7 (100)	

^a F: frugivore, N: nectarivore, SI: sallying insectivore, BI: bark insectivore, FI: foliage insectivore, O: omnivore.

Table 2. Continuation.

	Guild ^a	Year (1993–2010)												r	masl								
		93	94	95	96	97	98	99	00	01	02	03	04		05	06	07	08	09	10	100	200	250
<i>Xerops minutus</i>	FI	4	2	7	7	2	2	4	6	4	4	2	2	1	3	2	2	2	2	7 (12)	17 (29)	9 (16)	25 (43)
<i>Automolus ochrolaemus</i>	FI	10	2	4	5	1	6	3	10	4	6	4	6	4	8	4	6	2	5	-	5 (6)	2 (2)	83 (92)
<i>Mionectes olegineus</i>	O	100	101	128	104	50	115	63	70	40	41	48	66	37	31	46	64	23	36	416 (36)	397 (34)	222 (19)	128 (11)
<i>Leptopogon amarocephalus</i>	SI	8	6	2	6	8	2	8	8	7	1	4	4	4	8	1	6	7	7	28 (30)	38 (41)	24 (26)	3 (3)
<i>Oncostoma cinereigulare</i>	SI	9	5	8	2	8	9	8	4	5	3	5	3	5	3	3	3	3	1	9 (11)	52 (62)	13 (15)	10 (12)
<i>Platyrinchus canerominus</i>	SI	13	10	17	14	14	14	13	14	11	10	7	7	11	11	5	9	13	10	44 (22)	47 (23)	46 (23)	66 (33)
<i>Oryzorchyriacus coronatus</i>	SI	3	5	3	3	3	4	2	1	2	1	4	1	1	1	2	2	1	1	20 (57)	9 (26)	4 (11)	2 (6)
<i>Terenotriccus erythrorus</i>	SI	9	6	5	6	7	3	4	5	6	7	3	8	4	8	5	6	5	2	47 (47)	22 (22)	17 (17)	13 (13)
<i>Miyobius sulphureipygus</i>	SI	9	5	9	8	9	16	17	5	18	8	9	10	4	7	4	5	4	3	44 (29)	14 (9)	35 (23)	57 (38)
<i>Attila spadiceus</i>	SI	5	3	3	7	13	7	7	7	10	14	4	5	8	3	2	5	5	2	26 (24)	25 (23)	39 (35)	20 (18)
<i>Ceratopipa mentalis</i>	F	100	73	81	49	57	106	60	70	51	35	38	49	12	28	23	23	14	21	258 (29)	335 (38)	203 (23)	94 (11)
<i>Manacus candei</i>	F	1	2	4	6	7	5	10	5	3	2	2	2	2	2	3	1	1	1	5 (9)	34 (62)	14 (25)	2 (4)
<i>Turdichornis ochraceiceps</i>	FI	29	16	17	24	18	35	27	21	23	15	11	18	10	11	14	13	4	16	85 (26)	61 (19)	95 (30)	81 (25)
<i>Microcerculus philomela</i>	FI	1	2	2	1	1	1	2	1	2	1	1	1	1	2	1	1	1	1	-	2 (12)	1 (6)	14 (82)
<i>Pheugopedius maculipectus</i>	FI	2	2	1	1	2	4	3	4	5	2	2	2	2	2	2	2	1	1	-	11 (42)	11 (42)	4 (15)
<i>Henicorhina leucosticta</i>	FI	27	34	28	26	23	19	29	31	28	29	28	23	20	21	16	7	20	12	93 (22)	64 (15)	92 (22)	172 (41)
<i>Ramphocaelus melanurus</i>	FI	2	2	5	2	4	1	4	4	4	2	1	1	1	2	2	1	1	1	5 (15)	14 (41)	5 (15)	10 (29)
<i>Miyadestes unicolor</i>	F	1	2	1	11	1	7	4	4	4	4	4	4	4	4	4	1	1	1	12 (33)	7 (19)	6 (17)	11 (31)
<i>Turdus assimilis</i>	O	6	1	3	18	4	5	1	7	7	2	1	11	3	9	1	1	7	7	-0.33	16 (20)	25 (31)	31 (39)
<i>Basileuterus culicivorus</i>	FI	1	3	3	2	1	2	2	6	2	3	1	1	1	1	4	2	6	6	-0.07	-	-	39 (100)
<i>Lanio aurantius</i>	SI	1	1	4	1	1	1	1	1	1	1	1	3	3	3	3	1	1	1	-	-	-	-
<i>Arremon aurantirostris</i>	O	4	6	6	1	9	12	2	7	2	6	9	6	4	4	2	4	3	6	2 (13)	3 (20)	4 (27)	6 (40)
<i>Habia rubica</i>	FI	4	4	5	5	9	9	12	2	7	2	6	9	6	4	4	2	4	6	-0.03	10 (30)	11 (33)	-
<i>Habia fuscicauda</i>	FI	4	5	6	3	1	2	8	11	13	8	9	10	7	2	8	1	4	1	-0.20	6 (6)	6 (6)	25 (27)
<i>Cyanoloxia cyanoides</i>	O	4	6	3	4	7	3	8	1	1	3	4	2	4	2	1	4	2	2	-0.05	28 (27)	26 (25)	48 (47)
<i>Euphonia gouldi</i>	F	60	18	26	49	31	50	27	40	34	19	16	17	24	27	14	6	25	19	-0.50	20 (36)	26 (47)	9 (16)
																				-0.62	137 (27)	84 (17)	180 (36)

^a F: frugivore, N: nectarivore, SI: sallying insectivore, BI: bark insectivore, FI: foliage insectivore, O: omnivore.

insectivores, 7 as bark insectivores, 15 as foliage insectivores, and 4 as omnivores. We captured a total of 8435 individuals for these 53 species. The foraging guild with the higher number of captures was the nectarivore guild (1915), followed by frugivores (1676), foliage insectivores (1607), omnivores (1331), bark insectivores (975), and sallying insectivores (931). Several guilds were dominated by one or a few species, omnivores being the most extreme case: 87% of captures were for a single species, the Ochre-bellied Flycatcher (*Mionectes oleagineus*). Frugivores were dominated by the Red-capped Manakin (*Ceratopipra mentalis*), with 53% of captures, and nectarivores were dominated by the Long-billed Hermit (*Phaethornis longirostris*), with

49% of captures. Bark insectivores were mostly represented by two species: Tawny-winged Woodcreeper (*Dendrocincla anabatina*) with 35% of captures, and the Wedge-billed Woodcreeper (*Glyphorynchus spirurus*), with

Table 3. Statistics for fitted models of bird abundance in Cerro San Gil Watershed Protection Reserve, eastern Guatemala. Predictive variables included in models are described in table 1.

	AICc	w _i
Total abundance		
Wet + Trend	215.3	0.60
Wet + Dry + Trend	216.1	0.40
Annual + Trend	295.9	0.00
Dry + Trend	379.7	0.00
Trend	379.8	0.00
Nectarivores		
Wet + Dry + Trend	236.4	0.99
Dry + Trend	246.4	0.01
Annual + Trend	277.7	0.00
Wet + Trend	336.3	0.00
Trend	346.2	0.00
Omnivores		
Wet + Dry + Trend	101.8	1.00
Wet + Trend	116.4	0.00
Dry + Trend	121.9	0.00
Trend	138.6	0.00
Annual + Trend	140.2	0.00
Frugivores		
Wet + Dry + Trend	141.4	0.99
Wet + Trend	150.3	0.01
Annual + Trend	180.1	0.00
Dry + Trend	215.3	0.00
Trend	226.3	0.00
Bark insectivores		
Wet + Trend	51.2	0.72
Wet + Dry + Trend	53.4	0.25
Annual + Trend	57.7	0.03
Trend	65.9	0.00
Dry + Trend	68.0	0.00
Foliage insectivores		
Annual + Trend	70.9	0.53
Trend	72.6	0.23
Dry + Trend	73.9	0.12
Wet + Trend	74.6	0.08
Wet + Dry + Trend	76.0	0.04
Sallying insectivores		
Wet + Trend	58.9	0.75
Wet + Dry + Trend	61.0	0.25
Trend	77.7	0.00
Annual + Trend	78.6	0.00
Dry + Trend	79.8	0.00




Figure 5. Temporal trends in total bird abundance and abundances for each foraging guild during the study period in Cerro San Gil Watershed Protection Reserve, eastern Guatemala. Pearson correlation coefficients between year and abundance are shown.

51% of captures. The other two guilds were more even in their distribution of individuals per species, although two or three species were significantly more abundant than other members of the guild.

When only considering the relationship between abundance and year (without considering rainfall effects), total abundance and the abundance of nectarivores, frugivores and omnivores showed significant, strong declines (Fig. 5). Fitted models of bird abundance with the highest support were those with wet-season rainfall, dry-season rainfall and temporal trend (average $w_i = 0.56$; Table 3), followed by the model containing wet-season rainfall and temporal trend (average $w_i = 0.31$). All other model types had average weights < 0.1 (Table 3). Upon consideration of weighted model averages, wet-season rainfall had the greatest positive effect on abundance, especially for total abundance and abundance of frugivores, followed by bark insectivores

and sallying insectivores (Fig. 6). In other words, total abundance and abundance for these guilds increased with increasing wet-season rainfall in the previous year. Foliage insectivore abundance was positively related to annual rainfall, but this effect was weak (the 95% confidence interval includes 0; Fig. 6). Nectarivore abundance was mainly affected by dry-season rainfall. Also, dry-season effects were negatively related to omnivore and frugivore abundance (greater abundance when the previous dry season had less rain), but these effects were weak. Finally, once rainfall effects had been taken into account, total abundance and the abundance of several guilds still showed decreases in abundance that were independent of rainfall. This was the case for total abundance, nectarivores, omnivores, frugivores, and to a lesser degree, foliage insectivores (Fig. 6).

DISCUSSION

In general, our results provide strong evidence of large, community-wide declines in bird abundance. More importantly, these declines were strongly associated with declines in rainfall amount. We hypothesized that precipitation has a direct effect on vegetation and insect productivity, and thus bird abundance is indirectly affected by rainfall amount, through its direct effects on bird resources (mainly food) (Powell et al. 2015). This hypothesis is consistent with the fact that most bird foraging guilds were strongly and negatively affected by decreases in rainfall amount, and thus the temporal decrease in rainfall is having an overall, community-wide effect. This result is also consistent with several, more species-focused studies in north temperate latitudes, which found direct evidence of a positive effect of rainfall amount on bird resources, and consequently on bird population dynamics (Silleet et al. 2000, Nott et al. 2002, Jones et al. 2003, Both and Visser 2005, Anders and Post 2006). In addition, in a Jamaican tropical dry forest, Brown and Sherry (2006) found that decreased precipitation associated with the dry season resulted in reduced arthropod activity at a time when migrants needed to accumulate fat in preparation for migration, causing a decrease in body condition of birds immediately prior to migration, which can in turn have demographic consequences on the breeding grounds




Figure 6. Average coefficients (weighted by Akaike weights) of fitted models of bird abundance in Cerro San Gil Watershed Protection Reserve, eastern Guatemala. Predictive variables included in models are described in table 1. Error bars show 95% confidence intervals.

(Morrissette et al. 2010, Norris 2005, Norris and Taylor 2006).

In our study site in eastern tropical Guatemala, total abundance of birds and the abundance of most foraging guilds (except for nectarivores and foliage insectivores) were mainly affected by wet-season precipitation. Because fruiting periods and insect biomass both peak during the rainy season, it is consistent that frugivore abundance, and bark and sallying insectivore abundance were related to wet-season rainfall. Omnivore abundance, which mainly consisted of one species that feeds on fruits and insects, the Ochre-bellied Flycatcher, was also strongly and positively associated to wet-season rainfall. Note that the fact that the omnivore guild was mainly represented by one species is clearly a caveat of the guild approach, if a guild is excessively dominated by a single species. It would probably not be an exaggeration to say that results related to this guild in fact reflect what is happening to this particular species. Species-specific analyses for this flycatcher are then called for.

Of the insectivores, only the foliage insectivores did not follow the pattern of association to wet-season rainfall, being only weakly associated to annual rainfall. Although using the guild approach might provide hypotheses on the observed declines (because of the logical causal chains between rainfall, plant and insect biomass and bird abundance), it might also clearly mask species-specific patterns. Several species for this guild had strong negative associations with "year" (a negative Pearson correlation coefficient; Table 2), indicating a negative population trend. It is thus a possibility that the grouping of species under the "foliage insectivore" denomination might be masking important species-specific patterns. A revision of the grouping strategy may be called for, and for those species which in principle indicate strong declines, specific analyses are necessary for a clearer picture regarding their population status.

The nectarivores were the only guild to be associated with dry-season precipitation. In a study carried out in Costa Rica, Stiles (1992) found severe negative effects on the local population of the Long-billed Hermit due to an unusual drought that caused a severe flower shortage during the peak of the breeding season, which occurs during the drier

months of the year. In our study site, this species is the most abundant understory hummingbird. It is thus plausible that the same mechanism that affected the Costa Rican population affected our local hummingbird population, given our result, thus providing some evidence for the bottle-neck hypothesis (Williams and Middleton 2008); i.e., that the amount of rainfall during the dry season, when flowering and hummingbird reproductive activity peaks, is a key driver of hummingbird populations.

Additional temporal decreases in abundance independent of rainfall amount in the previous year were detected for frugivores, nectarivores and omnivores. Evidently, the effects of the amount of rainfall may not only be limited to the rainy season immediately prior to captures, but to the amount of rainfall received in previous years. Time-lagged or carry-over effects are those effects of events that occur in one season but influence individual or population success in the following seasons, and recent empirical evidence suggests they may play an important role in bird population dynamics (Metcalf and Monaghan 2001, Norris 2005, Brown and Sherry 2006, Norris and Marra 2007, Morrissette et al. 2010). For example, the reproductive success of long-distance migratory birds is influenced by the quality of habitat in their wintering grounds in the season prior to breeding (Norris et al. 2004, Norris and Taylor 2006). Additionally, because of delayed parental care and reproduction in tropical birds (Stutchbury and Morton 2001, 2008, Russell et al. 2004), these effects may be evidenced much later during the life cycles of birds. Consequently, the total decline may be associated with different environmental effects (such as a reduction in rainfall) affecting different components (i.e., age classes) in bird populations, at different points in time.

Beyond those effects of precipitation amount which directly impact bird resources, other indirect pathways may exist of the effects of changes in climatic regimes. For example, Chase et al. (2005) found that for Song Sparrows (*Melospiza melodia*), nest predation rates were lower in wetter years, and Martin (2001) describes similar indirect effects of weather on reproductive success: changes in weather caused four bird species to shift to microhabitats that yielded decreased nesting

success. Biotic changes due to increasing temperature might also be expected: as climate warms, many types of vegetation and dependent organisms (including fruit trees and insects that many birds depend on) are expected to shift their distributions to track their preferred microclimates, possibly causing dispersal to higher altitudes (Sekercioglu et al. 2012). Thus, tropical lowland communities may experience net biotic attrition, as a result of the species moving to higher elevations not being replaced by other ones (Colwell et al. 2008, Harris et al. 2011, Sekercioglu et al. 2012). Lastly, synergies of climate change with other threats, such as habitat loss, emerging diseases, invasive species, and hunting may exacerbate the effects of climate change on tropical birds (Sekercioglu et al. 2012).

In conclusion, we detected strong, community-wide declines in species abundance, and these declines were strongly associated with declines in rainfall amount. Most importantly, the magnitude of the association between rainfall and bird abundance is large and not limited to a particular foraging guild, and we thus advocate a much greater effort to investigate current associations of animal abundance with weather, especially rainfall decline (and increase), in tropical regions. A more comprehensive understanding of the effects of climate change on the biotic abundance in tropical ecosystems is strongly needed to propose conservation actions for these diverse ecosystems.

ACKNOWLEDGEMENTS

We wish to thank the Foundation of Ecodevelopment and Conservation (FUNDAECO) of Guatemala, Neotropical Bird Conservation Act-USFWS, The Nature Conservancy-Tennessee Chapter, and the French Global Environmental Facility (Fond Français pour l'Environnement Mondial, FFEM), for financial support to carry out this research.

LITERATURE CITED

- AMERICAN ORNITHOLOGICAL SOCIETY (2016) *Checklist of North and Middle American birds*. American Ornithological Society, Chicago (URL: <http://checklist.aou.org/>)
- ANDERS AD AND POST E (2006) Distribution-wide effects of climate on population densities of a declining migratory bird. *Journal of Animal Ecology* 75:221–227
- ANDERSON DR, BURNHAM KP AND THOMPSON WL (2000) Null hypothesis testing: problems, prevalence and an alternative. *Journal of Wildlife Management* 64:912–923
- BEEVER EA, RAY C, WILKENING JL, BRUSSARD PF AND MOTE PW (2011) Contemporary climate change alters the pace and drivers of extinction. *Global Change Biology* 17:2054–2070
- BLAKE JG AND LOISELLE BA (2015) Enigmatic declines in bird numbers in lowland forest of eastern Ecuador may be a consequence of climate change. *PeerJ* 3:e1177
- BONEBRAKE TC, PONISIO LC, BOGGS CL AND EHRLICH PR (2010) More than just indicators: a review of tropical butterfly ecology and conservation. *Biological Conservation* 143:1831–1841
- BOTH C AND VISSER ME (2005) The effect of climate change on the correlation between avian life-history traits. *Global Change Biology* 11:1606–1613
- BROWN DR AND SHERRY TW (2006) Food supply controls the body condition of a migrant bird wintering in the tropics. *Oecologia* 149:22–32
- BURNHAM KP AND ANDERSON DR (1998) *Model selection and inference. A practical information-theoretic approach*. Springer-Verlag, New York
- BURNHAM KP AND ANDERSON DR (2001) Kullback-Leibler information as a basis for strong inference in ecological studies. *Wildlife Research* 28:111–119
- CAO M, PRINCE SD, SMALL J AND GOETZ SJ (2004) Remotely sensed interannual variations and trends in terrestrial net primary productivity 1981–2000. *Ecosystems* 7:233–242
- CATTANIO JH, ANDERSON AB, ROMBOLD JS AND NEPSTAD DC (2004) Phenology, litterfall, growth, and root biomass in a tidal floodplain forest in the Amazon estuary. *Brazilian Journal of Botany* 27:703–712
- CEREZO A, RAMÍREZ M, LÓPEZ A AND JAVIER O (2012) *The resident and migratory bird monitoring program of the Caribbean region of Guatemala: description and recent results*. Foundation for Ecodevelopment and Conservation, Guatemala
- CHASE MK, NUR N AND GEUPEL GR (2005) Effects of weather and population density on reproductive success and population dynamics in a Song Sparrow (*Melospiza melodia*) population: a long-term study. *Auk* 122:571–592
- CHEN I, SHIU H, BENEDICK S, HOLLOWAY JD, CHEY VK, BARLOW HS, HILL JK AND THOMAS CD (2004) Elevation increases in moth assemblages over 42 years on a tropical mountain. *Proceedings of the National Academy of Sciences* 106:1479–1483
- CLELAND EE, CHUINE I, MENZEL A, MOONEY HA AND SCHWARTZ MD (2007) Shifting plant phenology in response to global change. *Trends in Ecology and Evolution* 22:357–365
- COLWELL RK, BREHM G, CARDELÚS CL, GILMAN AC AND LONGINO JT (2008) Global warming, elevational range shifts, and lowland biotic attrition in the wet tropics. *Science* 322:258–261

- CRICK HQP (2004) The impact of climate change on birds. *Ibis* 146 (Suppl.):48–56
- DENLINGER DL (1980) Seasonal and annual variation of insect abundance in the Nairobi National Park, Kenya. *Biotropica* 12:100–106
- DORMANN CF, ELITH J, BACHER S, BUCHMANN C, CARL G, CARRÉ G, GARCÍA MARQUÉZ JR, GRUBER B, LAFOURCADE B, LEITAO PJ, MÜNDEMÜLLER T, MCCLEAN C, OSBORNE PE, REINEKING B, SCHRÖDER B, SKIDMORE AK, ZURELL D AND LAUTENBACH S (2013) Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 36:27–46
- FAABORG JW, ARENDT J, TOMS JD, DUGGER KM, COX WA AND MORA MC (2013) Long-term decline of a winter-resident bird community in Puerto Rico. *Biodiversity and Conservation* 22:63–75
- FODEN WB, BUTCHART SHM, STUART SN, VIÉ JC, AKCAKAYA HR, ANGULO A, DEVANTIER LM, GUTSCHE A, TURAK E, CAO L, DONNER SD, KATARIYA V, BERNARD R, HOLLAND RA, HUGHES AF, O'HANLON SE, GARNETT ST, SEKERCIOGLU CH AND MACE GM (2013) Identifying the world's most climate change vulnerable species: a systematic trait-based assessment of all birds, amphibians and corals. *PLoS One* 8:e65427
- FRANKIE GW, BAKER HG AND OPLER PA (1974) Comparative phenological studies of trees in tropical wet and dry forests in the lowlands of Costa Rica. *Journal of Ecology* 62:881–919
- FRITH CB AND FRITH DW (1985) Seasonality of insect abundance in an Australian upland tropical rainforest. *Australian Journal of Ecology* 10:237–248
- GORDO O AND SANZ JJ (2006) Climate change and bird phenology: a long-term study in the Iberian Peninsula. *Global Change Biology* 12:1993–2004
- GRAHAM MH (2003) Confronting multicollinearity in ecological multiple regression. *Ecology* 84:2809–2815
- HARRIS JB, SEKERCIOGLU CH, SODHI NS, FORDHAM DA, PATON DC AND BROOK BW (2011) The tropical frontier in avian climate impact research. *Ibis* 153:877–882
- HOLMGREN M, SCHEFFER M, EZCURRA E, GUTIÉRREZ JR AND MOHREN GM (2001) El Niño effects on the dynamics of terrestrial ecosystems. *Trends in Ecology and Evolution* 16:89–94
- INSIVUMEH (2012) *Climatic data for the 1993–2010 period*. Instituto Guatemalteco de Vulcanología, Meteorología e Hidrología, Guatemala
- IPCC (2014) *Climate change 2014. Impacts, adaptation and vulnerability*. Intergovernmental Panel on Climate Change, New York
- JETZ W, WILCOVE DS AND DOBSON AP (2007) Projected impacts of climate and land-use change on the global diversity of birds. *PLoS Biology* 5:1211–1219
- JONES J, DORAN PJ AND HOLMES RT (2003) Climate and food synchronize regional forest bird abundances. *Ecology* 84:3024–3032
- KARR JR AND BRAWN JD (1990) Food resources of understory birds in Central Panama: quantification and effects on avian populations. *Studies in Avian Biology* 13:58–64
- LOISELLE BA AND BLAKE JG (1991) Variation in resource abundance affects capture rates of birds in three lowland habitats in Costa Rica. *Auk* 108:114–130
- LOWMAN MD (1982) Seasonal variation in insect abundance among three Australian rain forests, with particular reference to phytophagous types. *Australian Journal of Ecology* 7:353–361
- MAHLI Y AND WRIGHT J (2004) Spatial patterns and recent trends in the climate of tropical rainforest regions. *Philosophical Transactions of the Royal Society of London, B* 359:311–329
- MARTIN TE (2001) Abiotic vs. biotic influences on habitat selection of coexisting species: climate change impacts? *Ecology* 82:175–188
- METCALFE NB AND MONAGHAN P (2001) Compensation for a bad start: grow now, pay later? *Trends in Ecology and Evolution* 16:254–260
- MORRISSETTE M, BÉTY J, GAUTHIER G, REED A AND LEFEBVRE J (2010) Climate, trophic interactions, density dependence and carry-over effects on the population productivity of a migratory Arctic herbivorous bird. *Oikos* 119:1181–1191
- NEELIN JD, MÜNNICH M, SU H, MEYERSON JE AND HOLLOWAY CE (2006) Tropical drying trends in global warming models and observations. *Proceedings of the National Academy of Sciences* 103:6110–6115
- NORRIS DR (2005) Carry-over effects and habitat quality in migratory populations. *Oikos* 109:178–186
- NORRIS DR AND MARRA PP (2007) Seasonal interactions, habitat quality, and population dynamics in migratory birds. *Condor* 109:535–547
- NORRIS DR, MARRA PP, KYSER TK, SHERRY TW AND RATCLIFFE LM (2004) Tropical winter habitat limits reproductive success on the temperate breeding grounds in a migratory bird. *Philosophical Transactions of the Royal Society, B* 271:59–64
- NORRIS DR AND TAYLOR CM (2006) Predicting the consequences of carry-over effects for migratory populations. *Biological Letters* 2:148–151
- NORTH AMERICAN BIRD BANDING COUNCIL (2001) *The North American Bird Banding Council's study guide*. North American Bird Banding Council, Point Reyes
- NOTT MP, DESANTE DE, SIEGEL RB AND PYLE P (2002) Influences of the El Niño/Southern Oscillation and the North Atlantic Oscillation on avian productivity in forests of the Pacific Northwest of North America. *Global Ecology and Biogeography* 11:333–342
- PARMESAN C (2006) Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution and Systematics* 37:637–669
- POLLOCK KH, NICHOLS JD, SIMONS TR, FARNSWORTH GL, BAILEY LL AND SAUER JR (2002) Large scale wildlife monitoring studies: statistical methods for design and analysis. *Environmetrics* 13:105–119
- POLLOCK KH AND OTTO MC (1983) Robust estimation of population size in closed animal populations from capture-recapture experiments. *Biometrics* 39:1035–1049

- POULIN B, LEFEBVRE G AND McNEIL R (1992) Tropical avian phenology in relation to abundance and exploitation of food resources. *Ecology* 73:2295–2309
- POWELL LL, CORDEIRO NJ AND STRATFORD JA (2015) Ecology and conservation of avian insectivores of the rainforest understory: a pantropical perspective. *Biological Conservation* 188:1–10
- R CORE TEAM (2014) *R: a language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna (URL: <http://www.R-project.org/>)
- RAUSCHER SA, GIORGI F, DIFFENBAUGH NS AND SETH A (2008) Extension and intensification of the Meso-American mid-summer drought in the twenty-first century. *Climate Dynamics* 31:551–571
- ROOT TL AND SCHNEIDER SH (2002) Climate change: overview and implications for wildlife. Pp. 1–56 in: SCHNEIDER SH AND ROOT TL (eds) *Wildlife responses to climate change: North American case studies*. Island Press, Washington DC
- RUSSELL EM, YOM-TOV Y AND GEFFEN E (2004) Extended parental care and delayed dispersal: northern, tropical, and southern passerines compared. *Behavioral Ecology* 15:831–838
- SAATCHI SS, HOUGHTON RA, DOS SANTOS ALVALÁ RC, SOARES JV AND YU Y (2007) Distribution of above-ground live biomass in the Amazon basin. *Global Change Biology* 13:816–837
- VAN SCHAİK CP, TERBORGH JW AND WRIGHT SJ (1993) The phenology of tropical forests: adaptive significance and consequences for primary consumers. *Annual Review of Ecology and Systematics* 24:353–377
- SCHLOSS AL, KICKLICHTER DW, KADUK J AND WITTENBERG U (1999) Comparing global models of terrestrial net primary productivity (NPP): comparison of NPP to climate and the Normalized Difference Vegetation Index (NDVI). *Global Change Biology* 5:25–34
- SCHUUR EA (2003) Productivity and global climate revisited: the sensitivity of tropical forest growth to precipitation. *Ecology* 84:1165–1170
- SEKERCIOGLU CH, PRIMACK RB AND WORMWORTH J (2012) The effects of climate change on tropical birds. *Biological Conservation* 148:1–18
- SILLETT TS, HOLMES RT AND SHERRY TW (2000) Impacts of a global climate cycle on population dynamics of a migratory songbird. *Science* 288:2040–2042
- STILES FG (1992) Effects of a severe drought on the population biology of a tropical hummingbird. *Ecology* 73:1375–1390
- STILES FG AND SKUTCH AF (1989) *A guide to the birds of Costa Rica*. Cornell University, Ithaca
- STUTCHBURY BJM AND MORTON ES (2001) *Behavioral ecology of tropical birds*. Academic Press, London
- STUTCHBURY BJM AND MORTON ES (2008) Recent advances in the behavioral ecology of tropical birds. *Wilson Journal of Ornithology* 120:26–37
- STYRSKY JN AND BRAUN JD (2011) Annual fecundity of a Neotropical bird during years of high and low rainfall. *Condor* 113:194–199
- TERBORGH J, ROBINSON SK, PARKER TA III, MUNN CA AND PIERPONT N (1990) Structure and organization of an Amazonian forest bird community. *Ecological Monographs* 60:213–238
- THOMAS CD, CAMERON A, GREEN RE, BAKKENES M, BEAUMONT LJ, COLLINGHAM YC, ERASMUS BFN, FERREIRA DE SIQUEIRA M, GRAINGER A, HANNAH L, HUGHES L, HUNTLEY B, VAN JAARSVELD AS, MIDGLEY GF, MILES L, ORTEGA-HUERTA MA, PETERSON AT, PHILLIPS OL AND WILLIAMS SE (2004) Extinction risk from climate change. *Nature* 427:145–148
- VINCENT PJ AND HAWORTH JM (1983) Poisson regression models of species abundance. *Journal of Biogeography* 10:153–160
- VISSER ME, BOTH C AND LAMBRECHTS MM (2004) Global climate change leads to mistimed avian reproduction. *Advances in Ecological Research* 35:89–110
- WALTHER G, POST E, CONVEY P, MENZEL A, PARMESAN C, BEEBEE TJC, FROMENTIN J, HOEGH-GULDBERG O AND BAIRLEIN F (2002) Ecological responses to recent climate change. *Nature* 416:389–395
- WILLIAMS BK, NICHOLS JD AND CONROY MJ (2002) *Analysis and management of animal populations*. Academic Press, New York
- WILLIAMS SE AND MIDDLETON J (2008) Climatic seasonality, resource bottlenecks, and abundance of rainforest birds: implications for global climate change. *Diversity and Distributions* 14:69–77
- WOLDA H (1978) Seasonal fluctuations in rainfall, food and abundance of tropical insects. *Journal of Animal Ecology* 47:369–381
- WOLFE JD, PYLE P AND RALPH CJ (2009) Breeding seasons, molt patterns, and gender and age criteria for selected northeastern Costa Rican resident landbirds. *Wilson Journal of Ornithology* 121:556–567
- WOLFE JD AND RALPH CJ (2009) Correlations between El Niño–Southern Oscillation and changes in Nearctic–Neotropical migrant condition in Central America. *Auk* 126:809–814
- WOLFE JD, RALPH CJ AND ELIZONDO P (2015) Changes in the apparent survival of a tropical bird in response to the El Niño Southern Oscillation in mature and young forest in Costa Rica. *Oecologia* 178:715–721
- ZUUR AF, IENO EN, WALKER NJ, SAVELIEV AA AND SMITH GM (2009) *Mixed effects models and extensions in ecology with R*. Springer, New York