

Classifying smooth lattice polytopes via toric fibrations

Alicia Dickenstein^{a,1}, Sandra Di Rocco^{b,2}, Ragni Piene^{c,*}

^a *Departamento de Matemática, FCEN, Universidad de Buenos Aires, Ciudad Universitaria – Pab. I, (1428) Buenos Aires, Argentina*

^b *Department of Mathematics, KTH, SE-10044 Stockholm, Sweden*

^c *CMA/Department of Mathematics, University of Oslo, PO Box 1053 Blindern, NO-0316 Oslo, Norway*

Received 17 September 2008; accepted 7 April 2009

Available online 22 April 2009

Communicated by Ravi Vakil

Abstract

We show that any smooth \mathbb{Q} -normal lattice polytope P of dimension n and degree d is a strict Cayley polytope if $n \geq 2d + 1$. This gives a sharp answer, for this class of polytopes, to a question raised by V.V. Batyrev and B. Nill.

© 2009 Elsevier Inc. All rights reserved.

MSC: 14M25; 52B20

Keywords: Lattice polytope; Toric variety; Toric fibration; Cayley polytope; Nef value

1. Introduction

Let P be an n -dimensional *lattice polytope* (i.e., a convex polytope with integer vertices) in \mathbb{R}^n . We represent it as an intersection of half spaces

* Corresponding author.

E-mail addresses: alidick@dm.uba.ar (A. Dickenstein), dirocco@math.kth.se (S. Di Rocco), ragnip@math.uio.no (R. Piene).

¹ A.D. was partially supported by UBACYT X042 and X064, CONICET PIP 5617 and ANPCyT PICT 20569, Argentina.

² S.D.R. was partially supported by Vetenskapsrådet's grant NT:2006-3539, and the CIAM (Center of Industrial and Applied Mathematics).




Fig. 1.

$$P = \bigcap_{i=1}^r H_{\rho_i, -a_i}^+,$$

where $H_{\rho_i, -a_i}^+ = \{x \in \mathbb{R}^n \mid \langle \rho_i, x \rangle \geq -a_i\}$ are the half spaces, $H_{\rho_i, -a_i} = \{x \in \mathbb{R}^n \mid \langle \rho_i, x \rangle = -a_i\}$ the supporting hyperplanes, ρ_i the corresponding primitive inner normals, and $a_i \in \mathbb{Z}$, $i = 1, \dots, r$. Recall that an n -dimensional lattice polytope P is *smooth* if every vertex is equal to the intersection of n of the hyperplanes $H_{\rho_i, -a_i}$, and if the corresponding n normal vectors ρ_i form a lattice basis for $\mathbb{Z}^n \subset \mathbb{R}^n$. Smooth polytopes are sometimes called Delzant polytopes or regular polytopes.

Definition 1.1. Let $P = \bigcap_{i=1}^r H_{\rho_i, -a_i}^+ \subset \mathbb{R}^n$ be an n -dimensional lattice polytope. Define $P^{(s)} := \bigcap_{i=1}^r H_{\rho_i, -a_i+s}^+$, for $s \geq 1$.

Note that the lattice points of $P^{(1)}$ are precisely the interior lattice points of P .

Definition 1.2. Let P be a smooth n -dimensional lattice polytope and $s \geq 1$ an integer. Let m be a vertex of P . Reorder the hyperplanes so that $\{m\} = \bigcap_{i=1}^n H_{\rho_i, -a_i}$. We say that P is s -spanned at m if the lattice point $m(s)$, defined by $\{m(s)\} = \bigcap_{i=1}^n H_{\rho_i, -a_i+s}$, lies in $P^{(s)}$. We say that P is s -spanned if P is s -spanned at every vertex.

If $\{m\} = \bigcap_{i=1}^n H_{\rho_i, -a_i}$, we can write $m = (-a_1, \dots, -a_n)$ in the dual basis of ρ_1, \dots, ρ_n , and similarly $m(s) = (-a_1 + s, \dots, -a_n + s)$. It follows from the definition that if P is s -spanned, then $P^{(s)} \cap \mathbb{Z}^n \neq \emptyset$.

Example 1.3. Let P be the smooth polytope obtained from the simplex $d\Delta_3$ by removing the simplex $\Delta_3 = \text{Conv}\{(0:0:0), (1:0:0), (0:1:0), (0:0:1)\}$, see Fig. 1. Assume $d \geq 4$. Then $P^{(1)} \neq \emptyset$, but P is not 1-spanned. In fact, consider the vertex m of P , given by $\{m\} = \{(1:0:0)\} = (y=0) \cap (z=0) \cap (x+y+z=1)$, then the lattice point $m(1)$, given by $\{m(1)\} = \{(0:1:1)\} = (y=1) \cap (z=1) \cap (x+y+z=2)$, is not a point in $P^{(1)}$. Similarly for the vertices $(0:1:0)$ and $(0:0:1)$.

Note that if we instead remove $2\Delta_3$, then the resulting polytope is 1-spanned. In this case, the vertices $(2:0:0)$, $(0:2:0)$, and $(0:0:2)$ all “go” to the same lattice point $(1:1:1)$, which is an interior point of the polytope.

We recall the definitions of degree and codegree of a lattice polytope introduced in [1].

Definition 1.4. Let $P \subset \mathbb{R}^n$ be an n -dimensional lattice polytope. The *codegree* of P is the natural number

$$\text{codeg}(P) := \min_{\mathbb{N}} \{k \mid (kP)^{(1)} \cap \mathbb{Z}^n \neq \emptyset\}.$$

The *degree* of P is

$$\deg(P) := n + 1 - \text{codeg}(P).$$

We further introduce a more “refined” notion of codegree.

Definition 1.5. Let $P \subset \mathbb{R}^n$ be an n -dimensional lattice polytope. The \mathbb{Q} -*codegree* of P is defined as

$$\text{codeg}_{\mathbb{Q}}(P) := \inf_{\mathbb{Q}} \left\{ \frac{a}{b} \mid (aP)^{(b)} \neq \emptyset \right\}.$$

The number $\text{codeg}_{\mathbb{Q}}(P)$ is well defined, since $m(aP^{(b)}) = (maP)^{(mb)}$ and, for any polytope P' , we have $P' \neq \emptyset$ if and only if $mP' \neq \emptyset$ for every integer $m \geq 1$. Moreover, it is clear that $\text{codeg}_{\mathbb{Q}}(P) \leq \text{codeg}(P)$ holds.

Example 1.6. Let Δ_n denote the n -dimensional simplex. Then we have $\text{codeg}_{\mathbb{Q}}(2\Delta_n) = \frac{n+1}{2}$ and $\text{codeg}(2\Delta_n) = \lceil \frac{n+1}{2} \rceil$.

As we shall see in Proposition 2.2, the following definition embodies the *polytope version* of the notion of *nef value* for projective varieties.

Definition 1.7. Let $P \subset \mathbb{R}^n$ be an n -dimensional smooth lattice polytope. The *nef value* of P is

$$\tau(P) = \inf_{\mathbb{Q}} \left\{ \frac{a}{b} \mid aP \text{ is } b\text{-spanned} \right\}.$$

Remark 1.8. Clearly, the inequality $\tau(P) \geq \text{codeg}_{\mathbb{Q}}(P)$ always holds. It can be strict, as in Example 1.3, where $\text{codeg}(P) = \text{codeg}_{\mathbb{Q}}(P) = 1$ and $\tau(P) = 2$.

Observe also that when $\tau(P)$ is an integer, it follows from Lemma 2.4 below that $\tau(P) \geq \text{codeg}(P)$.

Definition 1.9. An n -dimensional lattice polytope P in \mathbb{R}^n is called \mathbb{Q} -*normal* if $\text{codeg}_{\mathbb{Q}}(P) = \tau(P)$.

We shall now explain the notion of generalized Cayley polytopes — these are particular examples of the twisted Cayley polytopes defined in [5].

Definition 1.10. Let $P_0, \dots, P_k \subset \mathbb{R}^m$ be lattice polytopes in \mathbb{Z}^m , e_1, \dots, e_k a basis for \mathbb{Z}^k and $e_0 = 0 \in \mathbb{Z}^k$. If $\{m_i^j\}_i$ are the vertices of P_j , so that $P_j = \text{Conv}\{m_i^j\}$ is the convex hull, and s is a positive integer, consider the polytope $\text{Conv}\{(m_i^j, se_j)\}_{i,j} \subseteq \mathbb{R}^{m+k}$. Any polytope P which

is affinely equivalent to this polytope will be called an *sth order generalized Cayley polytope* associated to P_0, \dots, P_k , and it will be denoted by

$$P \cong [P_0 * P_1 * \dots * P_k]^s.$$

If all the polytopes P_0, \dots, P_k have the same normal fan Σ (equivalently, if they are strictly combinatorially equivalent), we call P *strict*, and denote it by

$$P \cong \text{Cayley}_\Sigma^s(P_0, \dots, P_k).$$

If in addition $s = 1$, we write $P \cong \text{Cayley}_\Sigma(P_0, \dots, P_k)$ and call P a *strict Cayley polytope*.

Smooth generalized strict Cayley polytopes are natural examples of \mathbb{Q} -normal polytopes. In Proposition 3.9 we give sufficient conditions for $P \cong \text{Cayley}_\Sigma^s(P_0, \dots, P_k)$ to be \mathbb{Q} -normal, and we compute the common value $\text{codeg}_{\mathbb{Q}}(P) = \tau(P)$ in this case.

In [1] Batyrev and Nill posed the following question:

Question. *Given an integer d , does there exist an integer $N(d)$ such that every lattice polytope of degree d and dimension $\geq N(d)$ is a Cayley polytope?*

A first general answer was given by C. Haase, B. Nill, and S. Payne in [9], where they prove the existence of a lower bound which is quadratic in d .

In this article we give an optimal linear bound in the case of smooth \mathbb{Q} -normal lattice polytopes, and show that polytopes of dimension greater than or equal to this bound are strict Cayley polytopes.

Answer. For n -dimensional smooth \mathbb{Q} -normal lattice polytope, we can take $N(d) = 2d + 1$. More precisely, if P is a smooth \mathbb{Q} -normal lattice polytope of dimension n and degree d such that $n \geq 2d + 1$, then P is a strict Cayley polytope.

Observe that the condition $n \geq 2\deg(P) + 1$ is equivalent to $\text{codeg}(P) \geq \frac{n+3}{2}$. Furthermore, note that our bound is sharp: consider the standard n -dimensional simplex Δ_n and let $P := 2\Delta_n$ as in Example 1.6. Then, $\tau(P) = \text{codeg}_{\mathbb{Q}}(P) = \frac{n+1}{2} = \frac{n+3}{2} - 1$ and P is not a Cayley polytope. It is, however, a generalized Cayley polytope, with $s = 2$. We conjecture that an even smaller linear bound, like $\frac{n+1}{s}$, should suffice for the polytope to be an *sth order generalized Cayley polytope*.

We shall deduce our answer from Theorem 1.12 below, which gives a characterization of \mathbb{Q} -normal smooth lattice polytopes with big codegree. Before stating our main theorem, we recall the notion of a defective projective variety.

Definition 1.11. Let $X \subset \mathbb{P}^N$ be a projective variety over an algebraically closed field and denote by $X^* \subset (\mathbb{P}^N)^\vee$ its dual variety in the dual projective space. Then X is *defective* if X^* is not a hypersurface, and its defect is the natural number $\delta = N - 1 - \dim(X^*)$.

Theorem 1.12. *Let $P \subset \mathbb{R}^n$ be a smooth lattice polytope of dimension n . Then the following statements are equivalent:*

- (1) P is \mathbb{Q} -normal and $\text{codeg}(P) \geq \frac{n+3}{2}$,
- (2) $P = \text{Cayley}_\Sigma(P_0, \dots, P_k)$ is a smooth strict Cayley polytope, where $k+1 = \text{codeg}(P)$ and $k > \frac{n}{2}$,
- (3) the (complex) toric polarized variety (X, L) corresponding to P is defective, with defect $\delta = 2 \text{codeg}(P) - 2 - n$.

We conjecture that the assumption $\tau(P) = \text{codeg}_{\mathbb{Q}}(P)$ always holds for smooth lattice polytopes satisfying $\text{codeg}(P) \geq \frac{n+3}{2}$, and we therefore expect that the above classification holds for all smooth polytopes.

Our proof of Theorem 1.12 relies on the study of the nef value of nonsingular toric polarized varieties. This is developed in Section 2. Section 3 contains the study of generalized Cayley polytopes in terms of fibrations. In particular, for smooth generalized strict Cayley polytopes $P^s = \text{Cayley}_\Sigma^s(P_0, \dots, P_k)$ such that $s < k+1$ and $\dim P_i + 1 < \frac{k+1}{s}$ for all i , Proposition 3.9 shows that $\text{codeg}(P^s) > 1$, providing a family of lattice polytopes without interior lattice points. Section 4 contains the proof of the classification Theorem 1.12, together with some final comments.

2. The codegree and the nef value

Let X be a nonsingular projective variety over an algebraically closed field, and let L be an ample line bundle (or divisor) on X .

Definition 2.1. Assume that the canonical divisor K_X is not nef. The *nef value* of (X, L) is defined as

$$\tau_L := \min_{\mathbb{R}} \{t \mid K_X + tL \text{ is nef}\}.$$

By Kawamata's rationality theorem [10, Prop. 3.1, p. 619], the nef value τ_L is a positive rational number.

Proposition 2.2. Let X be a nonsingular projective toric variety of dimension n , and let L be an ample line bundle on X . Let $P \subset \mathbb{R}^n$ be the associated n -dimensional smooth lattice polytope. Then

$$\tau_L = \tau(P).$$

Proof. If we write the polytope P as $\bigcap_{i=1}^r H_{\rho_i, -a_i}^+$, then $L = \sum_{i=1}^r a_i D_i$, where the D_i are the invariant divisors on X . Since $K_X = -\sum_{i=1}^r D_i$, the polytope associated to X and the adjoint line bundle $bK_X + aL$ is

$$P_{bK_X + aL} = \bigcap_{i=1}^r H_{\rho_i, -a \cdot a_i + b}^+ = (aP)^{(b)}.$$

The lattice points of $(aP)^{(b)}$ form a basis for the vector space of global sections of $bK_X + aL$,

$$H^0(X, bK_X + aL) = \bigoplus_{m \in (aP)^{(b)} \cap \mathbb{Z}^n} \mathbb{C} \chi^m$$

(see [13, Lemma 2.3, p. 72]).

Denote by Σ the fan of X , let $x(\sigma)$ be the fixed point associated to the n -dimensional cone $\sigma = \langle \rho_1, \dots, \rho_n \rangle \in \Sigma$, and let $U_\sigma = X \setminus (\bigcup_{\tau \not\supset \sigma} V(\tau))$ be the affine patch containing $x(\sigma)$. The restriction of a generator $\chi^m \in H^0(X, bK_X + aL)$ to U_σ is

$$\chi^m|_{U_\sigma} = \prod_1^n \chi_i^{(m, \rho_i) - (-a \cdot a_i + b)},$$

where χ_1, \dots, χ_n is a system of local coordinates such that $x(\sigma) = (0, \dots, 0)$. It follows that the line bundle $bK_X + aL$ is spanned, or globally generated, at $x(\sigma)$ (i.e., it has at least one nonvanishing section at $x(\sigma)$) if and only if the lattice point $(-a \cdot a_1 + b, \dots, -a \cdot a_n + b)$, written with respect to the dual basis of ρ_1, \dots, ρ_n , is in $(aP)^{(b)}$.

Because the base locus of a line bundle is invariant under the torus action, if it is nonempty, it must be the union of invariant subspaces. Hence it has to contain fixed points. It follows that $bK_X + aL$ is spanned if and only if it is spanned at each fixed point, hence if and only if the polytope aP is b -spanned. \square

Corollary 2.3. *Let $P \subset \mathbb{R}^n$ be an n -dimensional smooth lattice polytope. Then*

- (1) $\text{codeg}(P) = \text{codeg}_{\mathbb{Q}}(P) = \tau(P) = n + 1$ if and only if $P = \Delta_n$.
- (2) If $P \neq \Delta_n$, then $\text{codeg}_{\mathbb{Q}}(P) \leq \tau(P) \leq n$.

Proof. Let (X, L) be the polarized nonsingular toric variety corresponding to P . In [12, Cor. 4.2] it is proven that $K_X + H$ is spanned for any line bundle H on X such that $H \cdot C \geq n$ for all invariant curves C on X , unless $X = \mathbb{P}^n$ and $H = \mathcal{O}_{\mathbb{P}^n}(n)$.

Because the line bundle L is ample, we have $L \cdot C \geq 1$ and thus $nL \cdot C \geq n$, for all invariant curves C . It follows that if $X \neq \mathbb{P}^n$ or $nL \neq \mathcal{O}_{\mathbb{P}^n}(n)$, then $\tau_L = \tau(P) \leq n$. This proves (2). If $(X, L) = (\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$, then $\tau_L = \tau(P) = n + 1$, because $K_{\mathbb{P}^n} = \mathcal{O}_{\mathbb{P}^n}(-n - 1)$. The corresponding polytope is $P = \Delta_n$ and $\text{codeg}(P) = \text{codeg}_{\mathbb{Q}}(P) = \tau(P) = n + 1$, as stated in (1). \square

Lemma 2.4. *Let $P \subset \mathbb{R}^n$ be a smooth n -dimensional polytope with codegree c and nef value τ . Then $\tau \in \mathbb{Q}_{>0}$ and $\tau > c - 1$.*

Proof. Let (X, L) be the polarized projective toric variety associated to P . Then $\tau = \tau_L \in \mathbb{Q}_{>0}$. By [2, Lemma 0.8.3] $K_X + \tau L$ is nef (and not ample). For any $s \geq \tau$, $K_X + sL = (K_X + \tau L) + (s - \tau)L$ is also nef. When s is an integer, this implies that $K_X + sL$ is spanned, hence $(sP)^{(1)} \cap \mathbb{Z}^n \neq \emptyset$. Taking $s = c - 1$ and observing that $((c - 1)P)^{(1)}$ has no lattice points, we deduce that $c - 1 < \tau$, as claimed. \square

Let $\tau(P) = \frac{a}{b}$, where a, b are coprime. On complete toric varieties, nef line bundles with integer coefficients are spanned (see [12, Thm. 3.1]). It follows that the linear system $|bK_X + aL|$ defines a morphism

$$\varphi: X \rightarrow \mathbb{P}^N,$$

where $N = |(aP)^{(b)} \cap \mathbb{Z}^n| - 1$. The Remmert–Stein factorization gives $\varphi = f \circ \varphi_P$, where $\varphi_P: X \rightarrow Y$ is a morphism with connected fibers onto a normal toric variety Y such that $\dim Y = \dim(aP)^{(b)}$ and $f: Y \rightarrow \mathbb{P}^N$. Moreover, φ_P is the contraction of a face of the nef cone $\text{NE}(X)$ [3, Lemma 4.2.13, p. 94].

Remark 2.5. If the morphism φ_P contracts a line, i.e., if there is a curve C such that $L \cdot C = 1$ and $\varphi_P(C)$ is a point, then $\tau(P)$ is necessarily an integer. In fact $0 = (bK_X + aL) \cdot C$ implies $\frac{a}{b} = -K_X \cdot C \in \mathbb{Z}$.

Lemma 2.6. Let P be a smooth n -dimensional polytope and let (X, L) be the corresponding polarized toric variety. If $\tau(P) > \frac{n+1}{2}$, then there exists an invariant line on X contracted by the nef value morphism φ_P . In particular, $\tau(P) \in \mathbb{Z}$, and $\tau(P) \geq \text{codeg}(P)$. If, moreover, φ_P is not birational, then φ_P is the contraction of an extremal ray in the nef cone, unless n is even and $P = \Delta_{\frac{n}{2}} \times \Delta_{\frac{n}{2}}$.

Proof. Because the nef value morphism is the contraction of a face of the Mori cone, it contracts at least one extremal ray. Take C to be a generator of this ray. Recall that, by Mori's Cone theorem (see e.g. [6, p. 25]), $n+1 \geq -K_X \cdot C$. Because $(K_X + \tau(P)L) \cdot C = 0$, we have

$$n+1 \geq -K_X \cdot C = \tau(P)L \cdot C > \frac{n+1}{2}L \cdot C,$$

which gives $L \cdot C = 1$ and $\tau(P) = -K_X \cdot C \in \mathbb{Z}$. Lemma 2.4 gives $\tau(P) > \text{codeg}(P) - 1$ from which we deduce that $\tau(P) \geq \text{codeg}(P)$. If φ_P is not birational, the last assertion follows from [4, (3.1.1.1), p. 30]. \square

We will also need the following key lemma. This lemma, and its proof, is essentially the same as [3, Lemma 7.1.6, p. 157].

Lemma 2.7. Let (X, L) be the polarized nonsingular toric variety associated to a smooth \mathbb{Q} -normal lattice polytope P . Then the morphism φ_P is not birational.

Proof. Let $\frac{a}{b} = \tau(P)$. Assume the morphism φ_P is birational. By [3, Lemma 2.5.5, p. 60] there is an integer m and an effective divisor D on X such that

$$m(bK_X + aL) = L + D.$$

It follows that $D \in |mbK_X + (ma-1)L|$, and thus $\text{codeg}_{\mathbb{Q}}(P) \leq \frac{ma-1}{mb} < \frac{a}{b} = \tau(P)$, which contradicts the assumption that P is \mathbb{Q} -normal. \square

3. Generalized Cayley polytopes and toric fibrations

Strict generalized Cayley polytopes (recall Definition 1.10) correspond to particularly nice toric fibrations, namely projective bundles. We will compute their associated nef values and codegrees. We refer to [5, Section 3] for further details on toric fibrations.

Definition 3.1. A polarized toric fibration is a quintuple (f, X, Y, F, L) , where

1. X and Y are normal toric varieties with $\dim(Y) < \dim(X)$,
2. $f: X \rightarrow Y$ is an equivariant flat surjective morphism with connected fibers,
3. the general fiber of f is isomorphic to the (necessarily toric) variety F ,
4. L is an ample line bundle on X .




Fig. 2.

There is a 1–1 correspondence between polarized toric fibrations and fibrations of polytopes, making Definition 3.1 equivalent to the following.

Definition 3.2. Let $\pi : M \rightarrow \Lambda$ be a surjective map of lattices and let $P_0, \dots, P_k \subset M_{\mathbb{R}}$ be lattice polytopes. We call π a *fibration with fiber Δ* if

1. $\pi_{\mathbb{R}}(P_i) = m_i \in \Lambda$ for every $i = 0, \dots, k$,
2. m_0, \dots, m_k are all distinct and are the vertices of

$$\Delta := \text{Conv}\{m_0, \dots, m_k\} \subset \Lambda_{\mathbb{R}},$$

3. P_0, \dots, P_k have the same normal fan, Σ .

In [5, Lemma 3.6] it is proven that (f, X, Y, F, L) is a toric fibration if and only if the polytope $P \subset M_{\mathbb{R}}$ associated to (X, L) has the structure of a fibration. More precisely, (f, X, Y, F, L) is a toric fibration if and only if there is a sublattice $\Lambda^{\vee} \hookrightarrow M^{\vee}$ such that the dual map $\pi : M \rightarrow \Lambda$ is a fibration of polytopes with fiber $\text{Conv}\{\pi(P)\} \subset \Lambda \otimes_{\mathbb{Z}} \mathbb{R}$. Moreover, F is the toric variety defined by the inner normal fan of $\text{Conv}\{\pi(P)\}$, and every fiber of $f : X \rightarrow Y$, with the reduced scheme structure, is isomorphic to F .

Observe that by construction, $\text{Cayley}_{\Sigma}^s(P_0, \dots, P_k) \subset M_{\mathbb{R}} \cong \mathbb{R}^m \oplus \Lambda_{\mathbb{R}}$, where $\Lambda_{\mathbb{R}} = \mathbb{R}^k$. The projection

$$\pi : M \rightarrow \Lambda, \quad \pi(m, e) = e,$$

gives the polytope $\text{Cayley}_{\Sigma}^s(P_0, \dots, P_k)$ the structure of a fibration with fiber $s\Delta_k$.

It follows that $\text{Cayley}_{\Sigma}^s(P_0, \dots, P_k)$ defines a toric fibration $f : X \rightarrow Y$ with general fiber isomorphic to \mathbb{P}^k .

Example 3.3. The strict Cayley sum $\text{Cayley}_{\Sigma(\Delta_1)}^2(4\Delta_1, 2\Delta_1, 2\Delta_1)$ is associated to the toric fibration $(\pi, \mathbb{P}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(4) \oplus \mathcal{O}_{\mathbb{P}^1}(2) \oplus \mathcal{O}_{\mathbb{P}^1}(2)), \mathbb{P}^1, \mathbb{P}^2, \xi)$, where ξ is the tautological line bundle and π is the projection (see Fig. 2).

Remark 3.4. Even if all the P_i are smooth and the fiber Δ is smooth, the polytope $\text{Cayley}_{\Sigma}^s(P_0, \dots, P_k)$ is not necessarily smooth. Consider the polytope $\text{Cayley}_{\Sigma(\Delta_1)}^2(6\Delta_1, 5\Delta_1, 3\Delta_1)$ depicted in Fig. 3. At the vertex $(3, 0, 2)$, the first lattice points on the corresponding three edges give




Fig. 3.

the vectors $(2, 0, 2) - (3, 0, 2) = (-1, 0, 0)$, $(4, 1, 1) - (3, 0, 2) = (1, 1, -1)$, and $(6, 0, 0) - (3, 0, 2) = (3, 0, -2)$, which do not form a basis for the lattice.

Remark 3.5. When the polytopes P_i are not strictly combinatorially equivalent, the variety associated to the Cayley polytope $[P_0 \star \cdots \star P_k]^s$ is birationally equivalent to a toric variety associated to a strict Cayley polytope, in the following precise way.

The normal fan Σ defined by the Minkowski sum $P_0 + \cdots + P_k$ is a common refinement of the normal fans defined by the polytopes P_i (see e.g. [15, 7.12]). Let (X_{P_i}, L_i) be the polarized toric variety associated to the polytope P_i and let $\pi_i: Y \rightarrow X_{P_i}$ be the induced birational morphism, where Y is the toric variety defined by the fan Σ . Notice that the line bundle $\pi_i^* L_i$ on Y is spanned, and the associated polytope Q_i is affinely equivalent to P_i .

Set $P'_i = Q_i + \sum_{j=0}^k P_j$ for $i = 0, \dots, k$. Note that the P'_i are strictly combinatorially equivalent, since their inner normal fan is Σ . The normal fan of the polytope $\text{Cayley}_\Sigma^s(P'_0, \dots, P'_k)$ is then a refinement of the normal fan of the polytope $[P_0 \star \cdots \star P_k]^s$, and thus it defines a proper birational morphism

$$\pi: X_{\text{Cayley}_\Sigma^s(P'_0, \dots, P'_k)} \rightarrow X_{[P_0 \star \cdots \star P_k]^s}.$$

Lemma 3.6. Let $P_0, \dots, P_k \subset \mathbb{R}^m$ be strictly combinatorially equivalent polytopes such that the polytope $P^s = \text{Cayley}_\Sigma^s(P_0, \dots, P_k)$ is smooth. Let $f: X \rightarrow Y$ be the associated toric fibration. Then P_i is smooth for all $i = 0, \dots, k$, and all fibers are reduced and isomorphic to \mathbb{P}^k .

Proof. Let F be an invariant fiber of f , then, since it is not general, it must be the fiber over a fixed point of Y . Equivalently, there is a vertex m of $\text{Cayley}_\Sigma^s(P_0, \dots, P_k)$ which is the intersection of a k -dimensional face Q such that $\pi(Q) = s \Delta_k$ and an $(n - k)$ -dimensional face R such that $\pi(R) = e_i \in \mathbb{R}^k$. Hence $R = P_i$. Moreover, by construction, Q is a simplex (possibly non-standard) of edge lengths b_1, \dots, b_k , with $1 \leq b_i \leq s$. Because $\text{Cayley}_\Sigma^s(P_0, \dots, P_k)$ is smooth, the first lattice points m_1, \dots, m_n on the n edges meeting at m form a lattice basis. This is equivalent to asking that the $n \times n$ matrix formed by taking the integral vectors $m_1 - m, \dots, m_n - m$ as columns, has determinant ± 1 . After reordering we can assume that $e_0 = 0$. Then the corresponding matrix A has the shape featured in Fig. 4, where a_i , $1 \leq a_i \leq s$, corresponds to the coordinate of the first lattice point on the i th edge of the simplex Q . The matrix A_Y is the matrix given by the first lattice points through the (corresponding) vertex of P_0 . A standard computation in linear algebra gives $\det(A) = \det(A_Y) a_1 \cdots a_k$. It follows that $\det(A_Y) = 1$ and $a_1 = \cdots = a_k = 1$.

Because each vertex of P_i is the intersection of a smooth fiber with P_i , we conclude that P_i is smooth for all $i = 0, \dots, k$. The equalities $a_1 = \cdots = a_k = 1$ show that all invariant fibers have edge length s . Every special fiber (as a cycle) is a combination of invariant fibers. If there were a nonreduced fiber tF , then it should contain an invariant curve such that tC is numerically

$$\left[\begin{array}{c|cccc} & a_1 & 0 & \dots & 0 \\ & 0 & a_2 & \dots & 0 \\ & \dots & \dots & \dots & \dots \\ & 0 & 0 & \dots & a_k \\ \hline A_Y & & & & B \end{array} \right]$$

Fig. 4.

equivalent to C' , where C' is an invariant curve in $s\Delta$. Then, because $L \cdot C \geq s$ and $L \cdot C' = s$, we would have $t = 1$. Lemma 3.6 implies that the P_i are smooth, and hence Y is smooth. One can see this also from the standard fact that for every morphism with connected fibers between two normal toric varieties, the general fiber is necessarily toric [8, Lemma 1.2]. \square

Observe that the hypothesis that P^s is smooth in Lemma 3.6 above, is essential in order to prove that all fibers are reduced and embedded as Veronese varieties.

Toric projective bundles are isomorphic to projectivized bundles of a vector bundle, which necessarily splits as a sum of line bundles [8, Lemma 1.1]. We will denote the projectivized bundle by $\mathbb{P}_Y(L_0 \oplus \dots \oplus L_k)$, where Y is the toric variety associated to Σ .

Proposition 3.7. *Let $P_0, \dots, P_k \subset \mathbb{R}^m$ be strictly combinatorially equivalent polytopes such that*

$$P^s := \text{Cayley}_{\Sigma}^s(P_0, \dots, P_k)$$

is smooth. Then there are line bundles L_0, \dots, L_k on $Y = X(\Sigma)$ such that the toric variety $X(\Sigma_{P^s})$, defined by the inner normal fan Σ_{P^s} of P^s , is isomorphic to $\mathbb{P}_Y(L_0 \oplus \dots \oplus L_k)$.

Proof. Denote by L the ample line bundle on $X(\Sigma_{P^s})$ associated to the given polytope P^s . By Lemma 3.6, all fibers are isomorphic to \mathbb{P}^k and thus $X(\Sigma_{P^s})$ has the structure of a projective bundle over Y . Equivalently, $f_*(L) = L_0 \oplus \dots \oplus L_k$ and $X(\Sigma_{P^s}) \cong \mathbb{P}_Y(L_0 \oplus \dots \oplus L_k)$, where $f: X(\Sigma_{P^s}) \rightarrow Y$. \square

A complete description of the geometry of such varieties when $s = 1$ is contained in [7, Section 3] and [13, 1.1].

Throughout the rest of the section we will always assume that $P^s = \text{Cayley}_{\Sigma}^s(P_0, \dots, P_k)$ is a smooth polytope. Denote as before by (X, L) the associated polarized toric variety. Let $f: X \rightarrow Y$ be as above. The invariant curves on X are of two types:

1. pullbacks $f^*V(\alpha_i)$ of invariant curves from Y , corresponding to the edges of P_i ,
2. curves $V(\alpha_F)$ contained in a fiber F , corresponding to the edges on simplices $s\Delta^k$.

Line bundles on X can be written as $L = f^*(M) + a\xi$, where M is a line bundle on Y and ξ is the tautological line bundle on $\mathbb{P}_Y(L_0 \oplus \dots \oplus L_k)$.

Because a line bundle on a nonsingular toric variety is spanned, respectively (very) ample, if and only if the intersection with all the invariant curves is nonnegative, respectively positive, there is a well understood spannedness and ampleness criterion, see [7, Prop. 2].

Lemma 3.8. Let $L = f^*(M) + a\xi$ be a line bundle on $\mathbb{P}_Y(L_0 \oplus \cdots \oplus L_k)$. Then for every curve of type $f^*V(\alpha_i)$, we have $L \cdot f^*V(\alpha_i) = (M + sL_i) \cdot V(\alpha_i)$, and for every curve $V(\alpha_F)$ we have $V(\alpha_F) \cdot L = a$. Consequently

- (1) L is ample if and only if $a \geq 1$ and $M + sL_i$ is ample for all i ,
- (2) L is spanned if and only if $a \geq 0$ and $M + sL_i$ are spanned for all i .

Because the fibers of π correspond to simplices $s\Delta_k$, and thus are embedded as s -Veronese varieties, and because the line bundles L_i are ample, we see that $P^s = \text{Cayley}_\Sigma^s(P_0, \dots, P_k)$ corresponds to the toric embedding

$$(\mathbb{P}_Y(L_0 \oplus \cdots \oplus L_k), s\xi).$$

Proposition 3.9. Let $P^s = \text{Cayley}_\Sigma^s(P_0, \dots, P_k)$ be a smooth generalized strict Cayley polytope. Assume that $\dim P_i + 1 < \frac{k+1}{s}$ for all i . Then P^s is \mathbb{Q} -normal, and

$$\text{codeg}_{\mathbb{Q}}(P^s) = \tau(P^s) = \frac{k+1}{s}.$$

Proof. Recall that

$$X := X(P^s) = \mathbb{P}_Y(L_0 \oplus \cdots \oplus L_k),$$

where L_i is ample on $Y := X(\Sigma)$, for $i = 0, \dots, k$, and P^s is the polytope defined by the line bundle $s\xi$, where $\xi := \xi_{L_0 \oplus \cdots \oplus L_k}$.

Recall also that the canonical line bundle on X is $K_X = \pi^*(K_Y + L_0 + \cdots + L_k) - (k+1)\xi$, where $\pi : X \rightarrow Y$. It follows that

$$H := bK_X + as\xi = \pi^*(b(K_Y + L_0 + \cdots + L_k)) + (as - b(k+1))\xi.$$

By Lemma 3.8, H is spanned (resp. ample) if and only if $as - b(k+1) \geq 0$ (resp. ≥ 1) and $b(K_Y + L_0 + \cdots + L_k) + sL_i$ is spanned (resp. ample).

Observe that, because L_i is ample for each i , $K_Y + L_0 + \cdots + L_k$ is ample if $k+1 > \dim P_i + 1$ [12, Cor. 4.2 (ii)], which holds by assumption. It follows that, with the given hypotheses, we have

- (i) H is spanned if and only if $\frac{a}{b} \geq \frac{k+1}{s}$,
- (ii) H is ample if and only if $\frac{a}{b} > \frac{k+1}{s}$,

and thus $\tau(P) = \frac{k+1}{s}$.

Consider the projection map $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^k$ such that $\pi(P) = s\Delta_k$. Clearly, if a, b are such that $(aP)^{(b)} \neq \emptyset$, then for all points $m \in (aP)^{(b)}$, $\pi(m) \in (as\Delta_k)^{(b)}$. This implies that $\frac{a}{b} \geq \text{codeg}_{\mathbb{Q}}(s\Delta_k) = \frac{k+1}{s}$, and hence $\text{codeg}_{\mathbb{Q}}(P) \geq \frac{k+1}{s}$. Together with the inequality $\text{codeg}_{\mathbb{Q}}(P) \leq \tau(P) = \frac{k+1}{s}$, this proves

$$\text{codeg}_{\mathbb{Q}}(P) = \tau(P) = \frac{k+1}{s}. \quad \square$$

Remark 3.10. Christian Haase showed us the following beautiful geometric argument to prove Proposition 3.9. As before, denote by Σ the common normal fan of P_0, \dots, P_k and by $\Sigma(P^s)$ the normal fan of P^s . Consider the smooth toric varieties $X = X(\Sigma(P^s))$ and $Y = X(\Sigma)$, and let L denote the ample line bundle on X with associated polytope P^s and L_0, \dots, L_k the ample line bundles on Y with associated polytopes P_0, \dots, P_k .

As $\tau(s\Delta_k) = \frac{k+1}{s} > \tau(P_i)$, we have that an integer multiple of $L_i + \frac{s}{k+1}K_Y$ is ample on Y for all $i = 0, \dots, k$. Let ρ_1, \dots, ρ_ℓ be the primitive generators of the one-dimensional cones in Σ , so that

$$P_i = \bigcap_{j=1}^{\ell} H_{\rho_j, -a_{ij}}^+.$$

Then the polytope

$$\bigcap_{j=1}^{\ell} H_{\rho_j, -a_{ij} + \frac{s}{k+1}}^+,$$

corresponding to the line bundle (with coefficients in \mathbb{Q}) $L_i + \frac{s}{k+1}K_Y$, is combinatorially equivalent to P_i , for all $i = 0, \dots, k$. On the other hand, the polytope

$$\left\{ x = (x_1, \dots, x_k) \in s\Delta_k \mid \sum_{i=1}^k x_i \leq s - \frac{s}{k+1}, x_i \geq \frac{s}{k+1}, \forall i = 1, \dots, k \right\}$$

equals the barycenter v of the simplex $s\Delta_k$. Hence the polytope associated to $L + \frac{s}{k+1}K_X$ reduces to the fiber over v in P^s , which can be identified with $\frac{1}{k+1}(P_0 + \dots + P_k)$, and no multiple of the corresponding line bundle is ample.

For any rational number $\frac{a}{b} > \frac{k+1}{s}$, we have again that the polytope

$$\bigcap_{j=1}^{\ell} H_{\rho_j, -a_{ij} + \frac{b}{a}}^+$$

is combinatorially equivalent to P_i for all $i = 0, \dots, k$. But now the polytope given by the points in $s\Delta_k$ at lattice distance $\frac{b}{a}$ from each of its facets is also combinatorially equivalent to $s\Delta_k$, and therefore the polytope corresponding to $L + \frac{b}{a}K_X$ is combinatorially equivalent to P^s . We conclude that $\tau(P^s) = \frac{k+1}{s}$, as wanted.

4. Classifying smooth lattice polytopes with high codegree

We now use the results in the previous sections to give the proof of Theorem 1.12.

Proof of Theorem 1.12. Let (X, L) be the nonsingular toric variety and ample line bundle associated to P .

Assume (1) holds. Because P is \mathbb{Q} -normal, Lemma 2.7 implies that the nef value map $\varphi := \varphi_P$ is not birational. Set $\tau := \tau(P)$. Lemma 2.4 gives that $\tau > \text{codeg}(P) - 1 \geq \frac{n+1}{2}$. If n is even and

$P = \Delta_{\frac{n}{2}} \times \Delta_{\frac{n}{2}}$, then $\text{codeg}(P) = \frac{n}{2} + 1 < \frac{n+3}{2}$. Therefore, Lemma 2.6 implies that τ is an integer and $\tau \geq \text{codeg}(P) \geq \frac{n+3}{2}$, and that $\varphi: X \rightarrow Y$ is a (nonbirational) contraction of an extremal ray in the nef cone $NE(X)$ of X .

By [14, Cor. 2.5, p. 404], we know that φ is flat, Y is a smooth toric variety, and, since X is smooth, a general fiber F is isomorphic to \mathbb{P}^k , where $k = \dim X - \dim Y$. Under this isomorphism, $L|_F = \mathcal{O}_{\mathbb{P}^k}(s)$, for some positive integer s . Let $\ell \subseteq F \cong \mathbb{P}^k$ be a line. Then, since F , and hence ℓ , is contracted by φ , we have

$$0 = (K_X + \tau L) \cdot \ell = K_X \cdot \ell + \tau L \cdot \ell = K_F \cdot \ell + \tau s = -(k+1) + \tau s.$$

We therefore get

$$\frac{n+1}{2} < \tau = \frac{k+1}{s} \leq \frac{n+1}{s},$$

which gives $s = 1$ and $\tau = \text{codeg}_{\mathbb{Q}}(P) = k+1 \in \mathbb{Z}$. By Lemma 2.4, we have $\tau \geq \text{codeg}(P)$. As $\text{codeg}(P) \geq \text{codeg}_{\mathbb{Q}}(P)$, we get $\text{codeg}(P) = k+1$. Hence $k+1 \geq \frac{n+3}{2}$, so that $k > \frac{n}{2}$.

Since $L^k \cdot F = 1$ for a general fiber of φ and φ is flat, $L^k \cdot Z = 1$ for every fiber Z . Therefore all fibers are irreducible, reduced, and of degree one in the corresponding embedding. It follows that for every fiber Z , $(Z, L|_Z) \cong (\mathbb{P}^k, \mathcal{O}_{\mathbb{P}^k}(1))$. Therefore φ is a fiber bundle: $X = \mathbb{P}_Y(\varphi_* L)$, where $\varphi_* L$ is a rank $k+1$ vector bundle. Since Y is toric, this bundle splits as a sum of line bundles $\varphi_* L = L_0 \oplus \cdots \oplus L_k$, and therefore P is a strict Cayley polytope. Hence (1) implies (2).

Assume (2) holds. By Proposition 3.9 (with $s = 1$), P is \mathbb{Q} -normal, and $\text{codeg}_{\mathbb{Q}}(P) = \tau = k+1$. Since $\text{codeg}_{\mathbb{Q}}(P)$ is an integer, $\text{codeg}(P) = \text{codeg}_{\mathbb{Q}}(P)$, hence $\text{codeg}(P) = k+1 > \frac{n}{2} + 1$, so $\text{codeg}(P) \geq \frac{n+3}{2}$. Therefore (1) holds.

The equivalence of (2) and (3) is essentially contained in [7]; here is a brief sketch of the proof.

Assume (2) holds. Since $k > \frac{n}{2}$, (X, L) is defective with defect $\delta = 2k - n$ [11, Prop. 5.12, p. 369] hence $\delta = 2 \text{codeg}(P) - 2 - n$, since $\text{codeg}(P) = k+1$.

Assume (3) holds, so that P is defective with defect $\delta = 2 \text{codeg}(P) - 2 - n \geq 1$. By [7, Thm. 2], then $P = \text{Cayley}_{\Sigma}(P_0, \dots, P_k)$ is a smooth strict Cayley polytope, with $k = \frac{n+\delta}{2} > \frac{n}{2}$ and $\text{codeg}(P) = \frac{n+\delta}{2} + 1 = k+1$. \square

We isolate the following result from the statement and proof of the previous theorem.

Corollary 4.1. *Let P be a smooth \mathbb{Q} -normal lattice polytope of dimension n , and assume $\text{codeg}(P) \geq \frac{n+3}{2}$. Then, $\tau := \tau(P) = \text{codeg}(P)$ is an integer and the associated polarized toric variety (X, L) is defective with defect $\delta = 2\tau - 2 - n$.*

The classification of smooth \mathbb{Q} -normal lattice polytopes of degree 0 and 1 follows from Theorem 1.12 (cf. [1] for the general case).

Corollary 4.2. *Assume P is a smooth, \mathbb{Q} -normal n -dimensional lattice polytope. Then*

- (1) $\deg(P) = 0$ if and only if $P = \Delta_n$,
- (2) $\deg(P) = 1$ if and only if $P = \text{Cayley}(P_0, \dots, P_{n-1})$ is a Lawrence prism (the P_i are segments) or $P = 2\Delta_2$.

Proof. (1) Because $\tau(P) > \text{codeg}(P) - 1 = n \geq \frac{n+1}{2}$, Lemma 2.6 implies that φ_P contracts a line. It follows that $\tau(P)$ is an integer, $\tau(P) \geq n + 1$ and thus $\tau(P) = \text{codeg}(P) = n + 1$. Theorem 1.12 implies that P is a strict Cayley polytope with $k = n$. This is equivalent to $P = \Delta_n$.

(2) For $n \geq 3$ the same argument as in (1) applies and gives that P is Cayley with $k = n - 1$. Assume $n = 2$. Since $\text{codeg}(P) = 2$, $P \neq \Delta_2$ and P has no interior lattice points. Let m be a vertex of P . Because P is smooth, the first lattice points on the two edges containing $m(\sigma)$ form a lattice basis. In this basis $m(\sigma) = (0, 0)$, and the lattice point $(1, 1)$ is not an interior point of P . Therefore there are only three possibilities: (i) One of the edges has length > 1 , the other edge has length 1, and the point $(1, 1)$ is on a third edge parallel to ρ_1 , which gives a strict Cayley polytope. (ii) Both edges have length 1, and $(1, 1)$ is the fourth vertex, hence $P = \Delta_1 \times \Delta_1 = \text{Cayley}_{\Sigma(\Delta_1)}(\Delta_1, \Delta_1)$. (iii) Both edges have length 2, and there is only one other edge containing $(1, 1)$. In this case $P = 2\Delta_2$. \square

We close the paper with some loose ends. In the Introduction we conjectured that any smooth lattice polytope P with $\text{codeg}(P) \geq \frac{n+3}{2}$ is in fact \mathbb{Q} -normal. This conjecture is supported by [3, 7.1.8], which suggests that for $n \leq 7$, if P is a smooth lattice polytope and $\text{codeg}_{\mathbb{Q}}(P) > \frac{n}{2}$, then P is \mathbb{Q} -normal. We also expect that without any smoothness assumptions, all lattice polytopes P with $\text{codeg}(P) \geq \frac{n+3}{2}$ are indeed Cayley polytopes.

Acknowledgments

We thank the Institute for Mathematics and Its Applications (IMA), Minneapolis, where this project began. We also thank the Department of Mathematics of KTH and the Göran Gustafsson Foundation, as well as the Centre of Mathematics for Applications (CMA) and Department of Mathematics of the University of Oslo, for sponsoring several visits, during which this paper was completed. We are grateful to Eva Maria Feichtner and Bernd Sturmfels for calling our attention to the paper [1], and to Eduardo Cattani for useful discussions. We would also like to thank Christian Haase and Benjamin Nill for very interesting discussions, and for showing us the alternative proof in Remark 3.10. Finally, we would like to thank the referee for thoughtful and most helpful comments.

References

- [1] Victor V. Batyrev, Benjamin Nill, Multiples of lattice polytopes without interior lattice points, *Mosc. Math. J.* 7 (2) (2007) 195–207.
- [2] Mauro C. Beltrametti, Andrew J. Sommese, On the adjunction theoretic classification of polarized varieties, *J. Reine Angew. Math.* 427 (1992) 157–192.
- [3] Mauro C. Beltrametti, Andrew J. Sommese, *The Adjunction Theory of Complex Projective Varieties*, de Gruyter Exp. Math., vol. 16, Walter de Gruyter & Co., Berlin, 1995.
- [4] Mauro C. Beltrametti, Andrew J. Sommese, Jarosław A. Wiśniewski, Results on varieties with many lines and their applications to adjunction theory, in: *Complex Algebraic Varieties*, Bayreuth, 1990, in: *Lecture Notes in Math.*, vol. 1507, Springer-Verlag, Berlin, 1992, pp. 16–38.
- [5] Cinzia Casagrande, Sandra Di Rocco, Projective \mathbb{Q} -factorial toric varieties covered by lines, *Commun. Contemp. Math.* 10 (3) (2008) 363–389.
- [6] Herbert Clemens, János Kollár, Shigefumi Mori, Higher-dimensional complex geometry, *Astérisque* 166 (1988).
- [7] Sandra Di Rocco, Projective duality of toric manifolds and defect polytopes, *Proc. London Math. Soc.* (3) 93 (1) (2006) 85–104.
- [8] Sandra Di Rocco, Andrew J. Sommese, Chern numbers of ample vector bundles on toric surfaces, *Trans. Amer. Math. Soc.* 356 (2) (2004) 587–598.

- [9] Christian Haase, Benjamin Nill, Sam Payne, Cayley decompositions of lattice polytopes and upper bounds for h^* -polynomials, preprint, arXiv:0804.3667, 2008, *J. Reine Angew. Math.*, in press.
- [10] Yujiro Kawamata, The cone of curves of algebraic varieties, *Ann. of Math.* (2) 119 (3) (1984) 603–633.
- [11] Antonio Lanteri, Daniele Struppa, Projective manifolds whose topology is strongly reflected in their hyperplane sections, *Geom. Dedicata* 21 (3) (1986) 357–374.
- [12] Mircea Mustață, Vanishing theorems on toric varieties, *Tohoku Math. J.* (2) 54 (3) (2002) 451–470.
- [13] Tadao Oda, *Convex Bodies and Algebraic Geometry*, *Ergeb. Math. Grenzgeb.* (3), vol. 15, Springer-Verlag, Berlin, 1988.
- [14] Miles Reid, Decomposition of toric morphisms, in: *Arithmetic and Geometry*, vol. II, in: *Progr. Math.*, vol. 36, Birkhäuser Boston, Boston, MA, 1983, pp. 395–418.
- [15] Günter M. Ziegler, *Lectures on Polytopes*, *Grad. Texts in Math.*, vol. 152, Springer-Verlag, New York, 1995.