



Contents lists available at ScienceDirect

# Journal of Differential Equations

[www.elsevier.com/locate/jde](http://www.elsevier.com/locate/jde)



## Optimal boundary holes for the Sobolev trace constant

Leandro Del Pezzo<sup>a</sup>, Julián Fernández Bonder<sup>a,\*</sup>, Wladimir Neves<sup>b</sup>

<sup>a</sup> IMAS-CONICET and Departamento de Matemática, FCEyN, Universidad de Buenos Aires, Pabellón 1, Ciudad Universitaria (1428), Buenos Aires, Argentina

<sup>b</sup> Instituto de Matemática, Universidade Federal do Rio de Janeiro, C.P. 68530, Cidade Universitária, 21945-970, Rio de Janeiro, Brazil

### ARTICLE INFO

*Article history:*

Received 12 November 2010

Available online 21 July 2011

MSC:

35J66

47R05

*Keywords:*

Steklov eigenvalues

$p$ -Laplace operator

Shape optimization

### ABSTRACT

In this paper we study the problem of minimizing the Sobolev trace Rayleigh quotient  $\|u\|_{W^{1,p}(\Omega)}^p / \|u\|_{L^q(\partial\Omega)}^p$  among functions that vanish in a set contained on the boundary  $\partial\Omega$  of given boundary measure.

We prove existence of extremals for this problem, and analyze some particular cases where information about the location of the optimal boundary set can be given. Moreover, we further study the shape derivative of the Sobolev trace constant under regular perturbations of the boundary set.

© 2011 Elsevier Inc. All rights reserved.

## 1. Introduction

Sobolev inequalities have proved to be a fundamental tool in order to study differential equations. Among Sobolev inequalities, one that has captured a great deal of attention in recent years is the Sobolev trace inequality that states

$$S \left( \int_{\partial\Omega} |u|^q d\mathcal{H}^{N-1} \right)^{p/q} \leq \int_{\Omega} |\nabla u|^p + |u|^p dx,$$

for every  $u \in W^{1,p}(\Omega)$  for some constant  $S > 0$ ,  $1 \leq q \leq p_*$ , where  $p_*$  is the critical exponent in the Sobolev trace immersion, i.e.  $p_* = p(N - 1)/(N - p)$  if  $1 < p < N$  and  $p_* = \infty$  if  $p \geq N$  (the equality

\* Corresponding author.

E-mail addresses: [ldpezzo@dm.uba.ar](mailto:ldpezzo@dm.uba.ar) (L. Del Pezzo), [jfbonder@dm.uba.ar](mailto:jfbonder@dm.uba.ar) (J. Fernández Bonder), [wladimir@im.ufrj.br](mailto:wladimir@im.ufrj.br) (W. Neves).

URL: <http://mate.dm.uba.ar/~jfbonder> (J. Fernández Bonder).

$q = p_*$  does not hold in the limit case  $p = N$ ). Here  $\mathcal{H}^s$  denotes, as usual, the  $s$ -dimensional Hausdorff measure,  $\Omega \subset \mathbb{R}^N$  is a smooth bounded domain (Lipschitz will be enough for most of our arguments).

In these inequalities, a fundamental role is played by the *optimal constants* and their associated *extremals*. That is, respectively, the largest possible constant  $S$  in the above inequality defined as

$$S = S_{p,q}(\Omega) := \inf_{u \in \mathcal{X}} \frac{\int_{\Omega} |\nabla u|^p + |u|^p \, dx}{\left(\int_{\partial\Omega} |u|^q \, d\mathcal{H}^{N-1}\right)^{p/q}}$$

and extremals, which are functions  $w \in \mathcal{X}$  where the above infimum is attained. Here  $\mathcal{X}$  is the space of *admissible functions*,  $\mathcal{X} := W^{1,p}(\Omega) \setminus W_0^{1,p}(\Omega)$ .

It is a well-known fact that if  $1 < p < N$  and  $1 \leq q \leq p_*$  or  $p \geq N$  and  $1 \leq q < \infty$  then the constant  $S$  is positive. For the existence of extremals, the only case which is nontrivial is the *critical* one,  $1 < p < N$  and  $q = p_*$  where the immersion  $W^{1,p}(\Omega) \subset L^{p_*}(\partial\Omega)$  is no longer compact (see, for instance [10,11]).

The critical case (i.e.  $1 < p < N$  and  $q = p_*$ ) was analyzed in [12] and [16]. In those papers the authors show that, under very mild assumptions on the domain  $\Omega$  (e.g. the existence of a boundary point of positive mean curvature) there exist extremals for  $S$ .

Motivated by some problems in shape optimization for stored energies under prescribed loadings, in [15] the authors study a variant of the trace inequality (see [15] for further discussion on the problem): Given a set  $A \subset \Omega$ , minimize the *Rayleigh quotient* over the class of functions that vanishes on  $A$ , i.e.

$$S(A) := \inf_{u \in \mathcal{X}_A} \frac{\int_{\Omega} |\nabla u|^p + |u|^p \, dx}{\left(\int_{\partial\Omega} |u|^q \, d\mathcal{H}^{N-1}\right)^{p/q}}$$

where

$$\mathcal{X}_A := \{u \in \mathcal{X} : u = 0 \text{ a.e. on } A\}.$$

In the above mentioned paper [15], existence of extremals for  $S(A)$  is proved in the subcritical case  $q < p_*$  (see [16] for the critical case) and moreover the following shape optimization problem is studied: Minimize  $S(A)$  among measurable sets  $A \subset \Omega$  such that  $\mathcal{H}^N(A) = \alpha \mathcal{H}^N(\Omega)$  for some fixed  $0 < \alpha < 1$ . A set  $A^*$  that minimizes  $S(A)$  is called an optimal set.

In [15] the existence of optimal sets is established and some geometric properties of optimal sets are analyzed. Moreover, in the case  $p = 2$  the interior regularity of optimal sets is studied in [14]. See [13], where some asymptotic behaviors of optimal sets are studied (see also, Section 4). Further, in [8] and in [4] the so-called *shape derivative* for  $S(A)$  is computed with respect to regular deformations on the set  $A$ .

One observes that, in all the above mentioned works, the sets where the test functions are forced to vanish are *interior* sets, i.e.  $A \subset \Omega$  of positive Lebesgue measure. However, the important case of boundary sets, i.e.  $\Gamma \subset \partial\Omega$  was not treated previously. Hence, the main objective of this work is to fill this gap.

So, in this paper we study the best Sobolev trace constant from  $W^{1,p}(\Omega)$  into  $L^q(\partial\Omega)$  for functions that vanish on a subset  $\Gamma$  of  $\partial\Omega$ , i.e.

$$S(\Gamma) := \inf_{u \in \mathcal{X}_{\Gamma}} \frac{\int_{\Omega} |\nabla u|^p + |u|^p \, dx}{\left(\int_{\partial\Omega} |u|^q \, d\mathcal{H}^{N-1}\right)^{p/q}} \tag{1.1}$$

where

$$\mathcal{X}_{\Gamma} := \{u \in \mathcal{X} : u = 0 \text{ } \mathcal{H}^{N-1}\text{-a.e. } \Gamma\}.$$

Here, we consider exponents  $1 \leq q < p_*$ , so that the immersion  $W^{1,p}(\Omega) \subset L^q(\partial\Omega)$  turns out to be compact. Therefore, the existence of extremals for  $S(\Gamma)$  follows by direct minimization.

The critical case could be treated by the same method employed in [16]. However, we will not do it in this article.

Next, we study the following optimization problem: Given  $0 < \alpha < 1$ , we look for the value

$$S(\alpha) := \inf\{S(\Gamma) : \Gamma \subset \partial\Omega, \mathcal{H}^{N-1}(\Gamma) = \alpha\mathcal{H}^{N-1}(\partial\Omega)\}. \tag{1.2}$$

A set  $\Gamma^* \subset \partial\Omega$  is called an *optimal boundary hole*, when it realizes the above infimum, i.e.  $S(\Gamma^*) = S(\alpha)$  and  $\mathcal{H}^{N-1}(\Gamma^*) = \alpha\mathcal{H}^{N-1}(\partial\Omega)$ .

One of the main issues of this paper is to show the existence and geometric properties of optimal boundary holes.

*Organization of the paper*

The rest of the paper is organized as follows. After a short Section 2 where we collect some preliminary remarks, in Section 3 we establish the existence of optimal boundary holes. In Section 4, we analyze the simpler case where the domain  $\Omega$  is a Euclidean ball given a complete characterization of optimal boundary holes for this simpler geometry. In order to have a better understanding of more complex geometries, in Section 5 we use a dimension reduction technique to deal with domains that are stretched in some directions. Finally, in Section 6, we compute the so-called *shape derivative* of  $S(\Gamma)$  for regular deformations of a fixed boundary hole  $\Gamma$ .

**2. Preliminary remarks**

In this very short section, we give some preliminary observations that will be helpful in the remaining of the paper.

First, observe that if  $u$  is an extremal for  $S(\Gamma)$  then  $u$  turns out to be a weak solution to the following Euler–Lagrange equation

$$\begin{cases} -\Delta_p u + |u|^{p-2}u = 0 & \text{in } \Omega, \\ |\nabla u|^{p-2} \frac{\partial u}{\partial \nu} = \lambda |u|^{q-2}u & \text{on } \partial\Omega \setminus \Gamma, \\ u = 0 & \text{on } \Gamma, \end{cases} \tag{2.1}$$

where  $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2}\nabla u)$  is the usual  $p$ -Laplacian,  $\frac{\partial}{\partial \nu}$  is the outer unit normal derivative and  $\lambda$  is a positive constant that depends on the normalization of  $u$ . This is  $u \in \mathcal{X}_\Gamma$  and

$$\int_{\Omega} |\nabla u|^{p-2}\nabla u \nabla \phi + |u|^{p-2}u\phi \, dx = \lambda \int_{\partial\Omega} |u|^{q-2}u\phi \, d\mathcal{H}^{N-1},$$

for every  $\phi \in \mathcal{X}_\Gamma$ . Observe that, if  $\|u\|_{L^q(\partial\Omega)} = 1$ , then  $\lambda = S(\Gamma)$ .

As a consequence of (2.1), we have the following remarks:

**Remark 2.1.** By the regularity results of [21], an extremal  $u$  of  $S(\Gamma)$ , verify that  $u \in C^{1,\delta}_{loc}(\Omega)$  for some  $0 < \delta < 1$ .

Moreover, by [20], if  $\partial\Omega \setminus \bar{\Gamma} \in C^{1,\eta}$ , then the regularity up to the boundary is  $u \in C^{1,\gamma}_{loc}(\bar{\Omega} \setminus \bar{\Gamma})$  for some  $0 < \gamma < 1$ .

**Remark 2.2.** If  $u$  is an extremal of  $S(\Gamma)$ , then we have that  $|u|$  is also an extremal of  $S(\Gamma)$ . Thus, using that  $|u|$  is a weak solution of (2.1) and the maximum principle (see [24]), we have that  $u$  has constant sign. Therefore, we can always assume that

$$u > 0 \text{ in } \Omega \text{ and } u \geq 0 \text{ on } \partial\Omega.$$

Moreover, by Hopf’s Lemma (see [24]) and the boundary regularity we obtain that nonnegative solutions  $u$  to (2.1) verify

$$u > 0 \text{ in } \overline{\Omega} \setminus \overline{\Gamma}.$$

Finally, we need the following lemma on pointwise convergence for Sobolev functions. We believe that this result is well-known but we were unable to find it in the literature.

**Lemma 2.3.** Let  $\{f_n\}_{n \in \mathbb{N}} \subset W^{1,p}(\Omega)$  with  $1 < p < N$  be such that  $f_n \rightarrow 0$  as  $n \rightarrow \infty$  in  $W^{1,p}(\Omega)$ . Then, there exist a subsequence  $\{f_{n_j}\}_{j \in \mathbb{N}} \subset \{f_n\}_{n \in \mathbb{N}}$  and a set  $B \subset \overline{\Omega}$  such that  $\text{cap}_p(B) = 0$  and

$$f_{n_j}(x) \rightarrow 0, \text{ as } j \rightarrow \infty \text{ for } x \in \overline{\Omega} \setminus B.$$

**Proof.** The lemma is a consequence of Lemma 1 and Theorem 1 in Section 4.8 of [6]. In fact, by Lemma 1 in Section 4.8 of [6], we have, for  $\alpha > 0$ , the Tchebyshev-type inequality

$$\text{cap}_p(Mf > \alpha) \leq \frac{C}{\alpha^p} \|f\|_{W^{1,p}(\Omega)}^p,$$

where  $C$  is a positive constant that depends only on  $N, p$  and  $Mf$  is the Hardy–Littlewood maximal function. So, if  $f_n \rightarrow 0$  in  $W^{1,p}(\Omega)$ , there exists a subsequence,  $\{f_{n_j}\}_{j \in \mathbb{N}}$  such that

$$\text{cap}_p(Mf_{n_j} > 1/j) < \frac{C}{2^j}.$$

Let us define  $A_j := \{Mf_{n_j} > 1/j\}$  and let  $B_m := \bigcup_{j=m}^\infty A_j$ . Therefore,

$$\text{cap}_p(B_m) \leq \sum_{j=m}^\infty \text{cap}_p(A_j) < C \sum_{j=m}^\infty \frac{1}{2^j}.$$

Now, if  $x \in \Omega \setminus B_m$ ,  $Mf_{n_j}(x) < 1/j$  and by Theorem 1, Section 4.8 of [6], it follows that  $|f_{n_j}(x)| < 1/j$ , so  $f_{n_j} \rightarrow 0$  as  $j \rightarrow \infty$  in  $\Omega \setminus B_m$  for all  $m \in \mathbb{N}$ .

Since  $\text{cap}_p(B_m) \rightarrow 0$  as  $m \rightarrow \infty$  the result follows.  $\square$

### 3. The existence of an optimal boundary hole

In this section, following ideas from [15], we first prove that  $S(\Gamma)$  is lower semi-continuous with respect to the hole (Theorem 3.1). Then, we prove the existence of an optimal boundary hole.

**Theorem 3.1.** Let  $\{\Gamma_\varepsilon\}_{\varepsilon > 0}$  be a family of positive  $\mathcal{H}^{N-1}$ -measurable subsets of  $\partial\Omega$  and  $\Gamma_0 \subset \partial\Omega$  be a positive  $\mathcal{H}^{N-1}$ -measurable set, such that

$$\chi_{\Gamma_\varepsilon} \xrightarrow{*} \chi_{\Gamma_0} \text{ *-weakly in } L^\infty(\partial\Omega),$$

where  $\chi_A$  is the characteristic function of the set  $A$ . Then,

$$S(\Gamma_0) \leq \liminf_{\varepsilon \rightarrow 0^+} S(\Gamma_\varepsilon).$$

**Proof.** Let  $\{\Gamma_n\}_{n \in \mathbb{N}}$  be a subsequence of  $\{\Gamma_\varepsilon\}_{\varepsilon > 0}$  such that

$$\mathcal{L} = \liminf_{\varepsilon \rightarrow 0} S(\Gamma_\varepsilon) = \lim_{n \rightarrow \infty} S(\Gamma_n).$$

For each  $n \in \mathbb{N}$ , we consider  $u_n \in \mathcal{X}_{\Gamma_n}$  to be an extremal of  $S(\Gamma_n)$ , such that

$$u_n \geq 0 \quad \text{and} \quad \|u_n\|_{L^q(\partial\Omega)} = 1.$$

Therefore, the sequence  $\{u_n\}_{n \in \mathbb{N}}$  is bounded in  $W^{1,p}(\Omega)$  and hence there exists a function  $u \in W^{1,p}(\Omega)$ , such that, for a subsequence still denoted by  $\{u_n\}_{n \in \mathbb{N}}$ ,

$$u_n \rightharpoonup u \quad \text{weakly in } W^{1,p}(\Omega), \tag{3.1}$$

$$u_n \rightarrow u \quad \text{strongly in } L^p(\Omega), \tag{3.2}$$

$$u_n \rightarrow u \quad \text{strongly in } L^q(\partial\Omega). \tag{3.3}$$

In particular, we have that  $u \geq 0$ ,  $\|u\|_{L^q(\partial\Omega)} = 1$  and

$$\|u\|_{W^{1,p}(\Omega)} \leq \liminf_{n \rightarrow \infty} \|u_n\|_{W^{1,p}(\Omega)}.$$

Moreover, for each  $n \in \mathbb{N}$ ,  $u_n = 0$   $\mathcal{H}^{N-1}$ -a.e. on  $\Gamma_n$ . Thus, as

$$\chi_{\Gamma_n} \xrightarrow{*} \chi_{\Gamma_0} \quad \text{* -weakly in } L^\infty(\partial\Omega)$$

and by (3.3), we have

$$0 = \lim_{n \rightarrow \infty} \int_{\Gamma_n} u_n \, d\mathcal{H}^{N-1} = \int_{\Gamma_0} u \, d\mathcal{H}^{N-1}.$$

Therefore, since  $u \geq 0$ , we have that  $u = 0$   $\mathcal{H}^{N-1}$ -a.e. on  $\Gamma_0$ . Thus  $u$  is an admissible function in the characterization of  $S(\Gamma_0)$  and

$$S(\Gamma_0) \leq \|u\|_{W^{1,p}(\Omega)}^p \leq \liminf_{n \rightarrow \infty} \|u_n\|_{W^{1,p}(\Omega)}^p = \mathcal{L}.$$

This finishes the proof.  $\square$

**Remark 3.2.** There isn't any monotonicity assumption on the family  $\{\Gamma_\varepsilon\}_{\varepsilon > 0}$ .

The continuity of  $S(\Gamma)$  with respect to the topology of Theorem 3.1 does not hold, as is shown in the following example:

**Example 3.3.** We take  $1 < p \leq N$ . The case for  $p > N$  is easier by the compact embedding of  $W^{1,p}(\Omega)$  into continuous functions.

Let  $\Omega$  be a bounded domain in  $\mathbb{R}^n$  that satisfies the interior ball condition for all  $x \in \partial\Omega$ . Let  $x_0 \in \partial\Omega$  and let  $E \subset \partial\Omega$  be set of zero  $\mathcal{H}^{N-1}$ -measure such that  $cap_p(E) > 0$  and there exists  $r > 0$  such that  $B(x_0, r) \cap E = \emptyset$ . Then, we take  $\Gamma = B_{\frac{r}{2}}(x_0) \cap \partial\Omega$  and  $\Gamma_n = \Gamma \cup E_n$  where  $E_n = \bigcup_{x \in E} B(x, \frac{1}{n}) \cap \partial\Omega$  for all  $n \in \mathbb{N}$ . Observe that

$$\chi_{\Gamma_n} \xrightarrow{*} \chi_{\Gamma} \quad \text{* -weakly in } L^\infty(\partial\Omega).$$

Let  $u_n$  be a positive normalized extremal for  $S(\Gamma_n)$ . If we assume that  $S(\Gamma_n) \rightarrow S(\Gamma)$  as  $n \rightarrow +\infty$ , we have that there exist  $u \in W^{1,p}(\Omega)$  such that, for a subsequence still denote  $\{u_n\}_{n \in \mathbb{N}}$ ,  $u_n \rightarrow u$  strongly in  $W^{1,p}(\Omega)$  and  $u_n \rightarrow u$  strongly in  $L^q(\partial\Omega)$ . Therefore  $u$  is a positive normalized extremal for  $S(\Gamma)$ . Moreover, by the Hopf’s Lemma,  $u_n > 0$  on  $\partial\Omega \setminus \Gamma_n$  and  $u > 0$  on  $\partial\Omega \setminus \Gamma$ .

On the other hand, by Lemma 2.3, there exist a subsequence  $\{u_{n_j}\}_{j \in \mathbb{N}}$  of  $\{u_n\}_{n \in \mathbb{N}}$  and a set  $B \subset \overline{\Omega}$  such that  $cap_p(B) = 0$  and  $u_{n_j}(x) \rightarrow u$  as  $j \rightarrow \infty$  for  $x \in \overline{\Omega} \setminus B$ . Then, as  $u_{n_j}(x) = 0$  for all  $x \in E$  and  $j \in \mathbb{N}$ , and  $cap_p(E) > 0$ , we have that  $u(x) = 0$  for all  $x \in E$ , contrary to  $u > 0$  on  $\partial\Omega \setminus \Gamma$ .

Next we prove the existence of an optimal boundary hole. For this, we first need to show the following lemma:

**Lemma 3.4.** For each  $\alpha \in (0, 1)$ ,  $S(\alpha)$  has also the following characterization:

$$S(\alpha) := \inf \left\{ \frac{\|v\|_{W^{1,p}(\Omega)}^p}{\|v\|_{L^q(\partial\Omega)}^p} : v \in \mathcal{X}, \mathcal{H}^{N-1}(\{v = 0\}) \geq \alpha \mathcal{H}^{N-1}(\partial\Omega) \right\}.$$

**Proof.** Let  $\alpha \in (0, 1)$  and

$$\tilde{S}(\alpha) := \inf \left\{ \frac{\|v\|_{W^{1,p}(\Omega)}^p}{\|v\|_{L^q(\partial\Omega)}^p} : v \in \mathcal{X}, \mathcal{H}^{N-1}(\{v = 0\}) \geq \alpha \mathcal{H}^{N-1}(\partial\Omega) \right\}.$$

We want to prove that  $S(\alpha) = \tilde{S}(\alpha)$ . For this, we proceed in two steps.

**Step 1.** First, we show that  $\tilde{S}(\alpha) \leq S(\alpha)$ .

Let  $\Gamma$  be a subset of  $\partial\Omega$  such that  $\mathcal{H}^{N-1}(\Gamma) = \alpha \mathcal{H}^{N-1}(\partial\Omega)$ . Let  $u \in \mathcal{X}_\Gamma$  be a nonnegative extremal for  $S(\Gamma)$ .

Observe that,  $u$  is an admissible function in the characterization of  $\tilde{S}(\alpha)$  and

$$\tilde{S}(\alpha) \leq \frac{\|u\|_{W^{1,p}(\Omega)}^p}{\|u\|_{L^q(\partial\Omega)}^p} = S(\Gamma).$$

Consequently, we have that  $\tilde{S}(\alpha) \leq S(\alpha)$ .

**Step 2.** Now, we show that  $S(\alpha) \leq \tilde{S}(\alpha)$ .

Let  $\{v_n\}_{n \in \mathbb{N}}$  be a minimizing sequence of  $\tilde{S}(\alpha)$ , i.e.  $v_n \in \mathcal{X}$ ,

$$\tilde{S}(\alpha) = \lim_{n \rightarrow \infty} \frac{\|v_n\|_{W^{1,p}(\Omega)}^p}{\|v_n\|_{L^q(\partial\Omega)}^p} \quad \text{and} \quad \mathcal{H}^{N-1}(\{v_n = 0\}) \geq \alpha \mathcal{H}^{N-1}(\partial\Omega) \quad \forall n \in \mathbb{N}.$$

Thus, for each  $n \geq 1$ , we take

$$\Gamma_n \subset \{v_n = 0\}$$

such that  $\Gamma_n$  is  $\mathcal{H}^{N-1}$ -measurable and  $\mathcal{H}^{N-1}(\Gamma_n) = \alpha \mathcal{H}^{N-1}(\partial\Omega)$ . Thus, we have

$$S(\alpha) \leq S(\Gamma_n) \leq \frac{\|v_n\|_{W^{1,p}(\Omega)}^p}{\|v_n\|_{L^q(\Omega)}^p} \quad \forall n \in \mathbb{N}.$$

Then, passing to the limit in the above inequality when  $n \rightarrow \infty$ , we have

$$S(\alpha) \leq \lim_{n \rightarrow \infty} S(\Gamma_n) = \lim_{n \rightarrow \infty} \frac{\|v_n\|_{W^{1,p}(\Omega)}^p}{\|v_n\|_{L^q(\Omega)}^p} = \tilde{S}(\alpha).$$

The proof is complete.  $\square$

Now, we establish the main results of this section.

**Theorem 3.5.** *Let  $0 < \alpha < 1$ . Then, there exist:*

- (a) *A set  $\Gamma_0 \subset \partial\Omega$ , such that  $\mathcal{H}^{N-1}(\Gamma_0) = \alpha \mathcal{H}^{N-1}(\partial\Omega)$  and  $S(\alpha) = S(\Gamma_0)$ ;*
- (b) *A function  $u \in \mathcal{X}$  with  $\mathcal{H}^{N-1}(\{u = 0\}) \geq \alpha \mathcal{H}^{N-1}(\partial\Omega)$ , such that*

$$S(\alpha) = \frac{\|u\|_{W^{1,p}(\Omega)}^p}{\|u\|_{L^q(\partial\Omega)}^p}.$$

**Proof.** We divide the proof into two steps.

**Step 1.** First, we prove (b).

Let  $\{v_n\}_{n \in \mathbb{N}}$  be a nonnegative normalized minimizing sequence for  $S(\alpha)$ , i.e. for each  $n \geq 1$ ,

$$0 \leq v_n \in \mathcal{X}, \quad \|v_n\|_{L^q(\partial\Omega)} = 1, \quad \mathcal{H}^{N-1}(\{v_n = 0\}) \geq \alpha \mathcal{H}^{N-1}(\partial\Omega),$$

and

$$\lim_{n \rightarrow \infty} \|v_n\|_{W^{1,p}(\Omega)}^p = S(\alpha).$$

Thus the sequence  $\{v_n\}_{n \in \mathbb{N}}$  is bounded in  $W^{1,p}(\Omega)$  and, therefore there exists a function  $u \in W^{1,p}(\Omega)$  and a subsequence still denote  $\{v_n\}_{n \in \mathbb{N}}$  such that

$$v_n \rightharpoonup u \quad \text{weakly in } W^{1,p}(\Omega), \tag{3.4}$$

$$v_n \rightarrow u \quad \text{strongly in } L^p(\Omega), \tag{3.5}$$

$$v_n \rightarrow u \quad \text{strongly in } L^q(\partial\Omega), \tag{3.6}$$

$$v_n \rightarrow u \quad \mathcal{H}^{N-1}\text{-a.e. in } (\partial\Omega). \tag{3.7}$$

From (3.6) and (3.7), we have that  $\|u\|_{L^q(\partial\Omega)} = 1$  and

$$\mathcal{H}^{N-1}(\{u = 0\}) \geq \limsup_{n \rightarrow \infty} \mathcal{H}^{N-1}(\{v_n = 0\}) \geq \alpha \mathcal{H}^{N-1}(\partial\Omega).$$

Thus,  $u$  is an admissible function in the definition of  $\mathcal{S}(\alpha)$ , and therefore

$$\mathcal{S}(\alpha) \leq \|u\|_{W^{1,p}(\Omega)}^p.$$

The reverse inequality is clear, since from (3.4)

$$\|u\|_{W^{1,p}(\Omega)}^p \leq \lim_{n \rightarrow \infty} \|v_n\|_{W^{1,p}(\Omega)}^p = \mathcal{S}(\alpha).$$

**Step 2.** We show that (b) implies (a).

By (b), there exists  $u \in \mathcal{X}$  such that  $\mathcal{H}^{N-1}(\{u = 0\}) \geq \alpha \mathcal{H}^{N-1}(\partial\Omega)$  and

$$\mathcal{S}(\alpha) = \frac{\|u\|_{W^{1,p}(\Omega)}^p}{\|u\|_{L^q(\partial\Omega)}^p}.$$

Thus, there exists a set  $\Gamma_0 \subset \{x \in \partial\Omega : u(x) = 0\}$   $\mathcal{H}^{N-1}$ -measurable such that

$$\mathcal{H}^{N-1}(\Gamma_0) = \alpha \mathcal{H}^{N-1}(\partial\Omega).$$

Then we have that

$$\mathcal{S}(\Gamma_0) \leq \frac{\|u\|_{W^{1,p}(\Omega)}^p}{\|u\|_{L^q(\partial\Omega)}^p} = \mathcal{S}(\alpha),$$

and  $\mathcal{H}^{N-1}(\Gamma_0) = \alpha \mathcal{H}^{N-1}(\partial\Omega)$ . Therefore

$$\mathcal{S}(\alpha) = \mathcal{S}(\Gamma_0).$$

This finishes the proof.  $\square$

In the next theorem we make a refinement of Theorem 3.5 and prove, under further regularity assumptions on  $\partial\Omega$ , that for any extremal  $u \in \mathcal{X}$ , it holds that  $\mathcal{H}^{N-1}(\{u = 0\}) = \alpha \mathcal{H}^{N-1}(\partial\Omega)$  (i.e.  $\Gamma_0 = \{u = 0\}$  with the notation of the above proof).

**Theorem 3.6.** *Let  $u \in \mathcal{X}$  be an extremal of  $\mathcal{S}(\alpha)$ . Then, if  $\Omega$  satisfies the interior ball condition, we have that*

$$\mathcal{H}^{N-1}(\{u = 0\}) = \alpha \mathcal{H}^{N-1}(\partial\Omega).$$

**Proof.** Let  $u \in \mathcal{X}$  be an extremal of  $\mathcal{S}(\alpha)$ , i.e.  $\mathcal{H}^{N-1}(\{u = 0\}) \geq \alpha \mathcal{H}^{N-1}(\partial\Omega)$  and

$$\mathcal{S}(\alpha) = \frac{\|u\|_{W^{1,p}(\Omega)}^p}{\|u\|_{L^q(\partial\Omega)}^p}.$$

By contradiction, suppose the thesis were false, then

$$\mathcal{H}^{N-1}(\{u = 0\}) > \alpha \mathcal{H}^{N-1}(\partial\Omega).$$

Since  $\mathcal{H}^s$  is a Borel regular measure ( $0 \leq s < \infty$ ), see [6], there exists a closed set  $\Gamma_0 \subset \{x \in \partial\Omega : u(x) = 0\}$  such that

$$\mathcal{H}^{N-1}(\{u = 0\}) > \mathcal{H}^{N-1}(\Gamma_0) > \alpha \mathcal{H}^{N-1}(\partial\Omega).$$



Consequently, it follows that

$$S(\alpha) \leq S(\Gamma_0).$$

On the other hand, the function  $u$  is admissible in the characterization of  $S(\Gamma_0)$ , hence

$$S(\Gamma_0) \leq \frac{\|u\|_{W^{1,p}(\Omega)}^p}{\|u\|_{L^q(\partial\Omega)}^p} = S(\alpha).$$

Therefore,  $S(\alpha) = S(\Gamma_0)$  and so  $u$  is also an extremal of  $S(\Gamma_0)$ . Thus  $u$  is a weak solution of the following problem

$$\begin{cases} -\Delta_p u + |u|^{p-2}u = 0 & \text{in } \Omega, \\ |\nabla u|^{p-2} \frac{\partial u}{\partial \nu} = \lambda |u|^{q-2}u & \text{on } \partial\Omega \setminus \Gamma_0, \\ u = 0 & \text{on } \Gamma_0, \end{cases} \tag{3.8}$$

where  $\lambda$  depends on the normalization of  $u$ . Moreover, by Remark 2.1,  $u \in C^{1,\gamma}_{loc}(\Omega \cup (\partial\Omega \setminus \Gamma_0))$  for some  $0 < \gamma < 1$  and we can assume that  $u > 0$  in  $\Omega$ .

Now, by our assumption on  $\Omega$  we can apply Hopf’s Lemma (cf. Remark 2.2), to get

$$\frac{\partial u}{\partial \nu} > 0 \quad \text{on } \{x \in \partial\Omega : u(x) = 0\} \setminus \Gamma_0.$$

That is a contradiction.  $\square$

**Corollary 3.7.** *The set function  $S$  is strictly increasing with respect to  $\alpha$ .*

**Proof.** It is clear that  $S(\alpha)$  is nondecreasing. Now, if we suppose that there exists  $0 < \alpha < \beta < 1$ , such that  $S(\alpha) = S(\beta)$ , then an extremal for  $S(\beta)$  is also an extremal for  $S(\alpha)$ . But, if  $u$  is an extremal for  $S(\beta)$ , then

$$\mathcal{H}^{N-1}(\{u = 0\}) = \beta \mathcal{H}^{N-1}(\partial\Omega) > \alpha \mathcal{H}^{N-1}(\partial\Omega),$$

which is a contradiction to Theorem 3.6. Thus,  $S$  is strictly increasing.  $\square$

#### 4. Example: the unit ball

Now, we study symmetry properties of optimal holes in the special case where  $\Omega$  is the unit ball,  $\Omega = B(0, 1)$ . First, we recall some of the definitions and results concerning spherical caps. We address the reader to [19,23].

##### Spherical symmetrization

Given a measurable set  $A \subset \mathbb{R}^N$ , the spherical symmetrization  $A^*$  of  $A$  is constructed as follows: for each positive  $r$ , take  $A \cap \partial B(0, r)$  and replace it by the spherical cap of the same  $\mathcal{H}^{N-1}$ -measure and center  $re_N$ . This can be done for almost all  $r$ . The union of these caps is  $A^*$ . Now, the spherical symmetrization  $u^*$  of a given measurable function  $u \geq 0$  defined on  $\Omega$  is constructed by symmetrizing the super-level sets so that, for all  $t$ ,  $\{u^* \geq t\} = \{u \geq t\}^*$ . See [19,23].

The following theorem is proved in [23] (see also [19]).

**Theorem 4.1.** (See [23].) Let  $u \in W^{1,p}(B(0, 1))$  and let  $u^*$  be its spherical symmetrization. Then  $u^* \in W^{1,p}(B(0, 1))$  and

$$\begin{aligned} \int_{B(0,1)} |\nabla u^*|^p \, dx &\leq \int_{B(0,1)} |\nabla u|^p \, dx, \\ \int_{B(0,1)} |u^*|^p \, dx &= \int_{B(0,1)} |u|^p \, dx, \\ \int_{\partial B(0,1)} |u^*|^q \, d\mathcal{H}^{N-1} &= \int_{\partial B(0,1)} |u|^q \, d\mathcal{H}^{N-1}. \end{aligned} \tag{4.1}$$

In this case we can prove the following

**Theorem 4.2.** Let  $\Omega = B(0, 1)$  and let  $0 < \alpha < 1$ . Then, there exists an optimal boundary hole which is a spherical cap. Moreover, when  $p = 2$ ,  $\Gamma$  is an optimal boundary hole if, and only if  $\Gamma$  is a spherical cap (up to sets of zero  $\mathcal{H}^{N-1}$ -measure).

**Proof.** Fix  $\alpha \in (0, 1)$ , by Theorem 3.5, there exists a function  $u \in \mathcal{X}$  such that  $\mathcal{H}^{N-1}(\{u = 0\}) = \alpha \mathcal{H}^{N-1}(\partial B(0, 1))$  and

$$S(\alpha) = \frac{\|u\|_{W^{1,p}(B(0,1))}^p}{\|u\|_{L^q(\partial B(0,1))}^p}.$$

Let  $u^*$  be the spherical symmetrization of  $u$ . Then  $u^*$  is an admissible function in the definition of  $S(\alpha)$  and, by Theorem 4.1,

$$S(\alpha) \leq \frac{\|u^*\|_{W^{1,p}(B(0,1))}^p}{\|u^*\|_{L^q(\partial B(0,1))}^p} \leq \frac{\|u\|_{W^{1,p}(B(0,1))}^p}{\|u\|_{L^q(\partial B(0,1))}^p} = S(\alpha).$$

Therefore

$$S(\alpha) = \frac{\|u^*\|_{W^{1,p}(B(0,1))}^p}{\|u^*\|_{L^q(\partial B(0,1))}^p}. \tag{4.2}$$

Moreover,  $\Gamma := \{x \in \partial B(0, 1) : u^*(x) = 0\}$  is a spherical cap and, since  $\mathcal{H}^{N-1}(\{u = 0\}) = \alpha \mathcal{H}^{N-1}(\partial B(0, 1))$ , we have that  $\mathcal{H}^{N-1}(\Gamma) = \alpha \mathcal{H}^{N-1}(\partial B(0, 1))$ . Then, using (4.2), we get that

$$S(\alpha) = S(\Gamma).$$

Now consider  $p = 2$ . Let  $\Gamma$  be an optimal boundary hole and let  $u$  be an extremal of  $S(\Gamma)$ . In this case, it is proved in [5] that if equality holds in (4.1) then for each  $0 < r \leq 1$  there exists a rotation  $R_r$  such that

$$u|_{\partial B(0,r)} = (u^* \circ R_r)|_{\partial B(0,r)}. \tag{4.3}$$

We can assume that the axis of symmetry  $e_N$  was taken so that  $R_1 = Id$ . Therefore  $u$  and  $u^*$  coincide on  $\partial B(0, 1)$ . Then the set  $\{x \in \partial B(0, 1) : u(x) = 0\}$  is a spherical cap and, by Theorem 3.6,  $\mathcal{H}^{N-1}(\{u = 0\}) = \alpha \mathcal{H}^{N-1}(\partial B(0, 1))$ .  $\square$

### 5. Dimension reduction

In this section, we are interested in the characterization of optimal boundary holes, when we shrink some of the dimensions of the set  $\Omega$ . This procedure of dimension reduction is interesting for such domains  $\Omega$ , where one of the directions is smaller than other ones. We begin with a fundamental case when the set  $\Omega$  is given by a Cartesian product, then we extend our results for more general domains.

The ideas in this section follow closely the ones in [9] where the behavior of the best Sobolev trace constant for shrinking domains was analyzed and [13] where the interior set problem was studied.

#### 5.1. The product case

Let  $\Omega_1$  and  $\Omega_2$  be bounded domains respectively in  $\mathbb{R}^n$  and  $\mathbb{R}^k$ , which are connected and have smooth boundaries. Set  $\Omega = \Omega_1 \times \Omega_2$  and for some  $0 < \mu < 1$ , define

$$\Omega_\mu = \Omega_1 \times \mu\Omega_2 = \{(x, \mu y) : (x, y) \in \Omega\}. \tag{5.1}$$

It is easy to see that  $\partial\Omega_\mu = \overline{\Omega}_1 \times \mu\partial\Omega_2 \cup \partial\Omega_1 \times \mu\overline{\Omega}_2$  and

$$\mathcal{H}^{N-1}(\partial\Omega_\mu) = \mu^{k-1}\mathcal{H}^n(\Omega_1)\mathcal{H}^{k-1}(\partial\Omega_2) + \mu^k\mathcal{H}^{n-1}(\partial\Omega_1)\mathcal{H}^k(\Omega_2), \tag{5.2}$$

where we recall that  $N = n + k$ . Moreover we see that, formally,  $\Omega_1$  represents the boundary of  $\Omega_\mu$  in the limiting process. This fact will be made clear a posteriori.

Now let  $u_\mu$  be a function defined in  $\Omega_\mu$ . We define, for each  $(x, y) \in \Omega$ ,

$$v_\mu(x, y) = u_\mu(x, \mu y).$$

Then,  $v_\mu$  is defined in  $\Omega$  and enjoys the same regularity than  $u_\mu$ . More precisely, we have the following

**Lemma 5.1.** *If  $u_\mu \in W^{1,p}(\Omega_\mu)$ , then  $v_\mu \in W^{1,p}(\Omega)$ . Moreover,*

$$\begin{aligned} &\mathcal{H}^{N-1}(\{u_\mu = 0\} \cap \partial\Omega_\mu) \\ &= \mu^{k-1}\mathcal{H}^{N-1}(\{v_\mu = 0\} \cap (\Omega_1 \times \partial\Omega_2)) + \mu^k\mathcal{H}^{N-1}(\{v_\mu = 0\} \cap (\partial\Omega_1 \times \Omega_2)). \end{aligned}$$

**Proof.** The regularity of  $v_\mu$  is clear. On the other hand, since  $\chi_B \equiv \chi_A \circ T_\mu$ , where

$$A = \{(x, \zeta) \in \Omega_\mu; u_\mu(x, \zeta) = 0\}, \quad B = \{(x, y) \in \Omega; v_\mu(x, y) = 0\},$$

and  $T_\mu : \Omega \rightarrow \Omega_\mu$   $T_\mu(x, y) = (x, \mu y)$ . We have that,

$$\begin{aligned} \mathcal{H}^{N-1}(A) &= \int_{\partial\Omega_\mu} \chi_A \, d\mathcal{H}^{N-1} \\ &= \iint_{\Omega_1 \times \mu\partial\Omega_2} \chi_A \, d\mathcal{H}^{k-1} \, dx + \iint_{\partial\Omega_1 \times \mu\Omega_2} \chi_A \, d\mathcal{H}^{n-1} \, dy \\ &= \mu^{k-1} \iint_{\Omega_1 \times \partial\Omega_2} \chi_B \, d\mathcal{H}^{k-1} \, dx + \mu^k \iint_{\partial\Omega_1 \times \Omega_2} \chi_B \, d\mathcal{H}^{n-1} \, dy \\ &= \mu^{k-1}\mathcal{H}^{N-1}(B \cap (\Omega_1 \times \partial\Omega_2)) + \mu^k\mathcal{H}^{N-1}(B \cap (\partial\Omega_1 \times \Omega_2)). \end{aligned}$$

The proof is now complete.  $\square$

In the remainder of this section, we consider subcritical exponents  $1 \leq q < p^*$ , where  $p^*$  is the critical exponent for the Sobolev embedding  $W^{1,p}(\Omega_1) \hookrightarrow L^q(\Omega_1)$ , given by

$$p^* = \frac{pn}{n-p} \quad \text{if } 1 \leq p < n \quad \text{or} \quad p^* = \infty \quad \text{if } p \geq n.$$

Given  $\alpha, \mu \in (0, 1)$ , we define

$$\mathcal{S}_\mu(\alpha) := \inf\{S(\Gamma) : \Gamma \subset \partial\Omega_\mu, \mathcal{H}^{N-1}(\Gamma) \geq \alpha \mathcal{H}^{N-1}(\partial\Omega_\mu)\}$$

and

$$\mathbb{S}(\alpha) := \inf\left\{ \frac{\|v\|_{W^{1,p}(\Omega_1)}^p}{\|v\|_{L^q(\Omega_1)}^p} : v \in W^{1,p}(\Omega_1), \mathcal{H}^n(\{x \in \Omega_1 : v(x) = 0\}) \geq \alpha \mathcal{H}^n(\Omega_1) \right\}.$$

Observe that  $\mathbb{S}(\alpha)$  is the best Sobolev constant of the embedding  $W^{1,p}(\Omega_1) \subset L^q(\Omega_1)$  for functions that vanish on a subset of  $\Omega_1$  of a given positive measure greater than or equal to  $\alpha \mathcal{H}^n(\Omega_1)$ .

**Remark 5.2.** Arguing as in Section 2 (cf. with [15] where the interior set case is studied), we can prove that for every  $0 < \alpha < 1$  there exists  $v_\alpha \in W^{1,p}(\Omega_1)$  such that

$$\mathcal{H}^n(\{x \in \Omega_1 : v_\alpha(x) = 0\}) = \alpha \mathcal{H}^n(\Omega_1) \quad \text{and} \quad \mathbb{S}(\alpha) = \frac{\|v_\alpha\|_{W^{1,p}(\Omega_1)}^p}{\|v_\alpha\|_{L^q(\Omega_1)}^p}.$$

Moreover,  $\mathbb{S}(\alpha)$  is strictly increasing as a function of  $\alpha$ .

Next, we give a characterization of the asymptotic, as  $\mu \rightarrow 0^+$ , behavior of  $\mathcal{S}_\mu(\alpha)$ . In fact, we see that, properly rescaled, the limit behavior is given by  $\mathbb{S}(\alpha)$ .

In order to do this, we need a couple of lemmas. The first one is easy and was proved in [8].

**Lemma 5.3.** (See [8, Lemma 3.1].) Let  $\Omega_1 \subset \mathbb{R}^n$  be a domain and let  $f_j, f : \Omega_1 \rightarrow \mathbb{R}$  be nonnegative measurable functions ( $j = 1, 2, \dots$ ) such that  $f_j \rightarrow f$  a.e. in  $\Omega_1$ . Set  $A_j = \{x \in \Omega_1 : f_j(x) = 0\}$  and  $A = \{x \in \Omega_1 : f(x) = 0\}$  and suppose that  $\mathcal{H}^n(A_j) \rightarrow \mathcal{H}^n(A)$  as  $j \rightarrow +\infty$ . Then

$$\lim_{j \rightarrow +\infty} \mathcal{H}^{N-1}(A_j \Delta A) = 0.$$

The second lemma gives the right continuity of  $\mathbb{S}(\alpha)$  with respect to  $\alpha$ .

**Lemma 5.4.** Let  $1 \leq p < n, 1 \leq q < p^*$  and  $0 < \alpha_0 < 1$ . Then,

$$\lim_{\alpha \rightarrow \alpha_0^+} \mathbb{S}(\alpha) = \mathbb{S}(\alpha_0).$$

Moreover, if we denote by  $v_\alpha$  a nonnegative extremal for  $\mathbb{S}(\alpha)$  normalized such that  $\|v_\alpha\|_{L^q(\Omega_1)} = 1$ , then there exists a sequence  $\{\alpha_j\}_{j \in \mathbb{N}}, \alpha_j > 0$  for every  $j \in \mathbb{N}$ , such that  $\alpha_j \rightarrow \alpha_0^+$  as  $j \rightarrow +\infty$  and

$$\lim_{j \rightarrow +\infty} v_{\alpha_j} = v \quad \text{strongly in } W^{1,p}(\Omega_1), \tag{5.3}$$

where  $v$  is a nonnegative extremal for  $\mathbb{S}(\alpha_0)$ .

Lastly, if  $A_j = \{x \in \Omega_1: v_{\alpha_j}(x) = 0\}$  and  $A = \{x \in \Omega_1: v(x) = 0\}$ , we have that

$$\lim_{j \rightarrow +\infty} \mathcal{H}^n(A_j \Delta A) = 0. \tag{5.4}$$

**Proof.** For this, we proceed in three steps.

**Step 1.** First, we prove that  $\mathbb{S}(\alpha) \rightarrow \mathbb{S}(\alpha_0)$  as  $\alpha \searrow \alpha_0$ .

We begin by observing that, since  $\mathbb{S}(\cdot)$  is increasing by Remark 5.2, there exist

$$\mathcal{L} = \lim_{\alpha \rightarrow \alpha_0^+} \mathbb{S}(\alpha) \quad \text{and} \quad \mathcal{L} \geq \mathbb{S}(\alpha_0). \tag{5.5}$$

On the other hand, by Remark 5.2, there exists  $v_{\alpha_0} \in W^{1,p}(\Omega_1)$  an extremal of  $\mathbb{S}(\alpha_0)$  such that  $\|v_{\alpha_0}\|_{L^q(\Omega_1)} = 1$  and

$$\mathcal{H}^n(A_{\alpha_0}) = \alpha_0 \mathcal{H}^n(\Omega_1),$$

where  $A_{\alpha_0} = \{x \in \Omega_1: v_{\alpha_0}(x) = 0\}$ .

Now we choose a smooth function  $\eta$  satisfying

$$\begin{cases} \eta = 0 & \text{in } B(0, 1), \\ \eta = 1 & \text{in } \mathbb{R}^n \setminus B(0, 2), \\ 0 \leq \eta \leq 1 & \text{and } \|\nabla \eta\|_{L^\infty(\mathbb{R}^n)} \leq 2. \end{cases}$$

Take  $x_0 \in \Omega_1 \setminus A_{\alpha_0}$  a point of density one (see definition in Chapter 1.7 of [6]) and for each  $\varepsilon > 0$ , set  $\eta_\varepsilon(x) = \eta(\frac{x-x_0}{\varepsilon})$  and  $w_\varepsilon = \eta_\varepsilon v_{\alpha_0} \in W^{1,p}(\Omega)$ . Observe that

$$\mathcal{H}^n(\{x \in \Omega_1: w_\varepsilon(x) = 0\}) > \alpha_0 \mathcal{H}^n(\Omega_1), \tag{5.6}$$

for  $\varepsilon$  sufficiently small and

$$\lim_{\varepsilon \rightarrow 0^+} \|w_\varepsilon\|_{L^q(\Omega_1)} = \|v_{\alpha_0}\|_{L^q(\Omega_1)} \quad \forall q \in [1, p^*]. \tag{5.7}$$

Moreover

$$\begin{aligned} \|\nabla w_\varepsilon\|_{L^p(\Omega_1)} &\leq \|\nabla \eta_\varepsilon v_{\alpha_0} + \eta_\varepsilon \nabla v_{\alpha_0}\|_{L^p(\Omega_1)} \\ &\leq \|\nabla \eta_\varepsilon v_{\alpha_0}\|_{L^p(\Omega_1)} + \|\nabla v_{\alpha_0}\|_{L^p(\Omega_1)} \\ &\leq \frac{C}{\varepsilon} \|v_{\alpha_0}\|_{L^p(B(x_0, 2\varepsilon) \setminus B(x_0, \varepsilon))} + \|\nabla v_{\alpha_0}\|_{L^p(\Omega_1)} \end{aligned}$$

and, by Hölder’s inequality, we get that

$$\|\nabla w_\varepsilon\|_{L^p(\Omega_1)} \leq C \|v_{\alpha_0}\|_{L^{p^*}(B(x_0, 2\varepsilon) \setminus B(x_0, \varepsilon))} + \|\nabla v_{\alpha_0}\|_{L^p(\Omega_1)}, \tag{5.8}$$

where  $C$  is a constant independent of  $\varepsilon$ . Then, by (5.6), there exist  $\delta > 0$  such that

$$\mathcal{H}^n(\{x \in \Omega_1: w_\varepsilon(x) = 0\}) > \alpha \mathcal{H}^n(\Omega_1) \quad \forall 0 < \alpha - \alpha_0 < \delta.$$

Therefore,  $w_\varepsilon$  is an admissible function in the definition of  $\mathbb{S}(\alpha)$  and, using (5.8), we have that

$$\begin{aligned}
 S(\alpha) &\leq \frac{\|w_\varepsilon\|_{W^{1,p}(\Omega_1)}^p}{\|w_\varepsilon\|_{L^q(\Omega_1)}^p} \\
 &\leq \frac{(C\|v_{\alpha_0}\|_{L^{p^*}(B(x_0,2\varepsilon)\setminus B(x_0,\varepsilon))} + \|\nabla v_{\alpha_0}\|_{L^p(\Omega_1)})^p + \|w_\varepsilon\|_{L^p(\Omega_1)}^p}{\|w_\varepsilon\|_{L^q(\Omega_1)}^p}
 \end{aligned}$$

for all  $\alpha > \alpha_0$ . Then, by (5.5),

$$\mathcal{L} \leq \frac{(C\|v_{\alpha_0}\|_{L^{p^*}(B(x_0,2\varepsilon)\setminus B(x_0,\varepsilon))} + \|\nabla v_{\alpha_0}\|_{L^p(\Omega_1)})^p + \|w_\varepsilon\|_{L^p(\Omega_1)}^p}{\|w_\varepsilon\|_{L^q(\Omega_1)}^p} \quad \forall \varepsilon > 0.$$

Lastly, taking limit as  $\varepsilon \rightarrow 0^+$  and using (5.7) and (5.5), we get that

$$\mathcal{L} \leq \frac{\|v_{\alpha_0}\|_{W^{1,p}(\Omega_1)}^p}{\|v_{\alpha_0}\|_{L^q(\Omega_1)}^p} = \mathbb{S}(\alpha_0) \leq \mathcal{L}.$$

Then, we have that

$$\lim_{\alpha \rightarrow \alpha_0^+} \mathbb{S}(\alpha) = \mathbb{S}(\alpha_0), \tag{5.9}$$

as we wanted to show.

**Step 2.** Now, we prove that (5.3) holds.

Let  $v_\alpha$  be a nonnegative extremal for  $\mathbb{S}(\alpha)$  normalized such that  $\|v_\alpha\|_{L^q(\Omega_1)} = 1$ . Thus, by (5.9), we have that

$$\mathbb{S}(\alpha_0) = \lim_{\alpha \rightarrow \alpha_0^+} \mathbb{S}(\alpha) = \lim_{\alpha \rightarrow \alpha_0^+} \|v_\alpha\|_{W^{1,p}(\Omega_1)}^p, \tag{5.10}$$

and therefore  $\{v_\alpha\}$  is bounded in  $W^{1,p}(\Omega_1)$ . Then, there exists a sequence  $\{\alpha_j\}$  such that  $\alpha_j \rightarrow \alpha_0^+$  as  $j \rightarrow +\infty$  and

$$v_{\alpha_j} \rightharpoonup v \quad \text{weakly in } W^{1,p}(\Omega_1), \tag{5.11}$$

$$v_{\alpha_j} \rightarrow v \quad \text{strongly in } L^p(\Omega_1), \tag{5.12}$$

$$v_{\alpha_j} \rightarrow v \quad \text{strongly in } L^q(\Omega_1), \tag{5.13}$$

$$v_{\alpha_j} \rightarrow v \quad \mathcal{H}^n\text{-a.e. in } (\Omega_1), \tag{5.14}$$

where  $v \in W^{1,p}(\Omega_1)$ . Since  $\|v_{\alpha_j}\|_{L^q(\Omega_1)} = 1$  for all  $j \in \mathbb{N}$ , using (5.13), we have that  $\|v\|_{L^q(\Omega_1)} = 1$  and by (5.14)  $v$  is nonnegative. By (5.10), (5.11) and (5.12), we get that

$$\mathbb{S}(\alpha_0) = \lim_{j \rightarrow +\infty} \|v_{\alpha_j}\|_{W^{1,p}(\Omega_1)}^p \geq \|v\|_{W^{1,p}(\Omega_1)}^p, \tag{5.15}$$

and using (5.14), we have that

$$\alpha_0 \mathcal{H}^n(\Omega_1) \leq \liminf_{j \rightarrow +\infty} \mathcal{H}^n(A_j) \leq \limsup_{j \rightarrow +\infty} \mathcal{H}^n(A_j) \leq \mathcal{H}^n(A), \tag{5.16}$$

where  $A_j = \{x \in \Omega_1: v_{\alpha_j}(x) = 0\}$  and  $A = \{x \in \Omega_1: v(x) = 0\}$ . Then,  $v$  is an admissible function in the definition of  $\mathbb{S}(\alpha_0)$ , and using (5.15), we get that

$$\mathbb{S}(\alpha_0) \leq \|v\|_{W^{1,p}(\Omega_1)}^p \leq \mathbb{S}(\alpha_0).$$

Therefore  $v$  is an extremal for  $\mathbb{S}(\alpha_0)$  and, by (5.10), we have

$$\lim_{j \rightarrow +\infty} \|v_{\alpha_j}\|_{W^{1,p}(\Omega_1)} = \|v\|_{W^{1,p}(\Omega_1)}. \tag{5.17}$$

Moreover, using (5.11) and (5.17), we can conclude that

$$\lim_{j \rightarrow +\infty} v_{\alpha_j} = v \text{ strongly in } W^{1,p}(\Omega_1).$$

**Step 3.** Lastly, we prove that (5.4) holds.

First, we prove that  $\mathcal{H}^n(A) = \alpha_0 \mathcal{H}^n(\Omega_1)$ . On the contrary, suppose that  $\mathcal{H}^n(A) > \alpha_0 \mathcal{H}^n(\Omega_1)$ , then there exists  $j_0$  such that  $\mathcal{H}^n(A) > \alpha_j \mathcal{H}^n(\Omega_1)$  for all  $j \geq j_0$  and therefore

$$\mathbb{S}(\alpha_0) = \|v\|_{W^{1,p}(\Omega_1)}^p > \mathbb{S}(\alpha_j) > \mathbb{S}(\alpha_0)$$

and we obtain a contradiction. Thus  $\mathcal{H}^n(A) = \alpha_0 \mathcal{H}^n(\Omega_1)$  and by (5.16)

$$\lim_{j \rightarrow +\infty} \mathcal{H}^n(A_j) = \mathcal{H}^n(A).$$

Then, by (5.14) and Lemma 5.3, we have that

$$\lim_{j \rightarrow +\infty} \mathcal{H}^n(A_j \Delta A) = 0.$$

This finishes the proof.  $\square$

We arrive now at the main result of this section.

**Theorem 5.5.** *Let  $0 < \alpha, \mu < 1, 1 \leq p < n$ , and  $1 \leq q < p^*$ , then*

$$\lim_{\mu \rightarrow 0^+} \frac{S_\mu(\alpha)}{\mu^{\frac{k(q-p)+p}{q}}} = \frac{\mathcal{H}^k(\Omega_2)}{\mathcal{H}^{k-1}(\partial\Omega_2)^{\frac{p}{q}}} \mathbb{S}(\alpha).$$

**Proof.** We begin by proving

$$\limsup_{\mu \rightarrow 0^+} \frac{S_\mu(\alpha)}{\mu^{\frac{k(q-p)+p}{q}}} \leq \frac{\mathcal{H}^k(\Omega_2)}{\mathcal{H}^{k-1}(\partial\Omega_2)^{\frac{p}{q}}} \mathbb{S}(\alpha).$$

Let

$$\alpha_\mu = \alpha \left( 1 + \mu \frac{\mathcal{H}^{n-1}(\partial\Omega_1) \mathcal{H}^k(\Omega_2)}{\mathcal{H}^n(\Omega_1) \mathcal{H}^{k-1}(\partial\Omega_2)} \right)$$

and take  $v \in W^{1,p}(\Omega_1)$  such that

$$\mathcal{H}^n(A) \geq \alpha_\mu \mathcal{H}^n(\Omega_1),$$

where

$$A = \{x \in \Omega_1: v(x) = 0\}.$$

Then, if we take  $u(x, y) = v(x)$  for all  $(x, y) \in \Omega_\mu$ , we have that

$$\begin{aligned} \mathcal{H}^{N-1}(\{w = 0\} \cap \partial\Omega_\mu) &\geq \mathcal{H}^{N-1}(\{w = 0\} \cap (\bar{\Omega}_1 \times \mu\partial\Omega_2)) \\ &\geq \mathcal{H}^{N-1}(A \times \mu\partial\Omega_2) \\ &= \mu^{k-1}\mathcal{H}^n(A)\mathcal{H}^{k-1}(\partial\Omega_2) \\ &\geq \mu^{k-1}\alpha_\mu\mathcal{H}^n(\Omega_1)\mathcal{H}^{k-1}(\partial\Omega_2) \\ &= \alpha\mathcal{H}^{N-1}(\partial\Omega_\mu). \end{aligned}$$

Therefore,  $u$  is an admissible function in the characterization of  $S_\mu(\alpha)$  (see Lemma 3.4), then

$$\begin{aligned} S_\mu(\alpha) &\leq \frac{\iint_{\Omega_\mu} |\nabla w|^p + |w|^p \, dx \, dy}{\left(\int_{\partial\Omega_\mu} |w|^q \, d\mathcal{H}^{N-1}\right)^{\frac{p}{q}}} \\ &= \frac{\mu^k\mathcal{H}^k(\Omega_2) \int_{\Omega_1} |\nabla v|^p + |v|^p \, dx}{\left(\mu^{k-1}\mathcal{H}^{k-1}(\partial\Omega_2) \int_{\Omega_1} |v|^q \, dx + \mu^k\mathcal{H}^k(\Omega_2) \int_{\partial\Omega_1} |v|^q \, d\mathcal{H}^{n-1}\right)^{\frac{p}{q}}} \\ &\leq \mu^{\frac{k(q-p)+p}{q}} \frac{\mathcal{H}^k(\Omega_2)}{\mathcal{H}^{k-1}(\partial\Omega_2)^{\frac{p}{q}}} \frac{\int_{\Omega_1} |\nabla v|^p + |v|^p \, dx}{\left(\int_{\Omega_1} |v|^q \, dx\right)^{\frac{p}{q}}}. \end{aligned}$$

Thus, taking infimum over all  $v \in W^{1,p}(\Omega_1)$  such that

$$\mathcal{H}^n(\{x \in \Omega_1: v(x) = 0\}) \geq \alpha_\mu\mathcal{H}^n(\Omega_1),$$

we get that

$$\frac{S_\mu(\alpha)}{\mu^{\frac{k(q-p)+p}{q}}} \leq \frac{\mathcal{H}^k(\Omega_2)}{\mathcal{H}^{k-1}(\partial\Omega_2)^{\frac{p}{q}}} \mathbb{S}(\alpha_\mu).$$

Therefore, using Lemma 5.4,

$$\limsup_{\mu \rightarrow 0^+} \frac{S_\mu(\alpha)}{\mu^{\frac{k(q-p)+p}{q}}} \leq \frac{\mathcal{H}^k(\Omega_2)}{\mathcal{H}^{k-1}(\partial\Omega_2)^{\frac{p}{q}}} \mathbb{S}(\alpha). \tag{5.18}$$

On the other hand, for each  $\mu$  there exists an extremal  $u_\mu \in W^{1,p}(\Omega_\mu)$  of  $S_\mu(\alpha)$  such that

$$\iint_{\Omega_1 \times \partial\Omega_2} |v_\mu|^q \, dx \, d\mathcal{H}^{k-1} + \mu \iint_{\partial\Omega_1 \times \Omega_2} |v_\mu|^q \, d\mathcal{H}^{n-1} \, dy = 1, \tag{5.19}$$

where  $v_\mu(x, y) = u_\mu(x, \mu y)$ .



Then,

$$\begin{aligned}
 S_\mu(\alpha) &= \frac{\iint_{\Omega_\mu} |\nabla u_\mu|^p + |u_\mu|^p \, dx \, dy}{(\int_{\partial\Omega_\mu} |u_\mu|^q \, d\mathcal{H}^{N-1})^{p/q}} \\
 &= \frac{\iint_{\Omega} (|\nabla_x v_\mu, \mu^{-1} \nabla_y v_\mu|^p + |v_\mu|^p) \mu^k \, dx \, dy}{(\mu^{k-1} \iint_{\Omega_1 \times \partial\Omega_2} |v_\mu|^q \, dx \, d\mathcal{H}^{k-1} + \mu^k \iint_{\partial\Omega_1 \times \Omega_2} |v_\mu|^q \, d\mathcal{H}^{n-1} \, dy)^{\frac{p}{q}}} \\
 &= \mu^{\frac{k(q-p)+p}{q}} \left( \iint_{\Omega} |\nabla_x v_\mu, \mu^{-1} \nabla_y v_\mu|^p + |v_\mu|^p \, dx \, dy \right).
 \end{aligned}$$

Thus,

$$\frac{S_\mu(\alpha)}{\mu^{\frac{k(q-p)+p}{q}}} = \iint_{\Omega} |\nabla_x v_\mu, \mu^{-1} \nabla_y v_\mu|^p + |v_\mu|^p \, dx \, dy \quad \forall \mu \in (0, 1). \tag{5.20}$$

Let  $\{\mu_j\}_{j \in \mathbb{N}}$  be a sequence such that  $\mu_j \rightarrow 0^+$  as  $j \rightarrow \infty$  and

$$\liminf_{\mu \rightarrow 0^+} \frac{S_\mu(\alpha)}{\mu^{\frac{k(q-p)+p}{q}}} = \lim_{j \rightarrow +\infty} \frac{S_{\mu_j}(\alpha)}{\mu_j^{\frac{k(q-p)+p}{q}}}.$$

To simplify the notation, we write  $v_j$  instead of  $v_{\mu_j}$  for all  $j \in \mathbb{N}$ .

Then, by (5.18), we have that  $\{v_j\}_{j \in \mathbb{N}}$  is bounded in  $W^{1,p}(\Omega)$ . Therefore, there exist a function  $v \in W^{1,p}(\Omega)$  and a subsequence of  $\{v_j\}_{j \in \mathbb{N}}$  (still denoted by  $\{v_j\}_{j \in \mathbb{N}}$ ) such that

$$v_j \rightharpoonup v \quad \text{weakly in } W^{1,p}(\Omega), \tag{5.21}$$

$$v_j \rightarrow v \quad \text{strongly in } L^p(\Omega), \tag{5.22}$$

$$v_j \rightarrow v \quad \text{strongly in } L^q(\partial\Omega). \tag{5.23}$$

Observe that, by (5.23), we have that

$$v_j \rightarrow v \quad \text{strongly in } L^q(\partial\Omega_1 \times \Omega_2), \tag{5.24}$$

$$v_j \rightarrow v \quad \text{strongly in } L^q(\Omega_1 \times \partial\Omega_2), \tag{5.25}$$

and, using (5.19), (5.24) and (5.25), we get

$$\iint_{\Omega_1 \times \partial\Omega_2} |v|^q \, dx \, d\mathcal{H}^{k-1} = 1,$$

from where we conclude that  $v \not\equiv 0$ .

Now, using again (5.18) and (5.20), we have that there exists a constant  $C$  such that

$$\iint_{\Omega} |\mu_j^{-1} \nabla_y v_j|^p \, dx \, dy \leq C \quad \forall j \in \mathbb{N},$$

then  $\{\mu_j^{-1}\nabla_y v_j\}_{j \in \mathbb{N}}$  is bounded in  $L^p(\Omega)$  and

$$\iint_{\Omega} |\nabla_y v_j|^p \, dx \, dy \leq C \mu_j^p \rightarrow 0 \quad \text{as } j \rightarrow \infty.$$

Therefore  $v$  does not depend on  $y$ , i.e.  $v = v(x)$  and

$$1 = \iint_{\Omega_1 \times \partial\Omega_2} |v|^q \, dx \, d\mathcal{H}^{k-1} = \mathcal{H}^{k-1}(\partial\Omega_2) \int_{\Omega_1} |v|^q \, dx. \tag{5.26}$$

On the other hand, using that  $\{\mu_j^{-1}\nabla_y v_j\}_{j \in \mathbb{N}}$  is bounded in  $L^p(\Omega)$ , there exist  $w \in L^p(\Omega)$  such that

$$\mu_j^{-1}\nabla_y v_j \rightharpoonup w \quad \text{weakly in } L^p(\Omega).$$

Then

$$\begin{aligned} \liminf_{\mu \rightarrow 0^+} \frac{S_{\mu}(\alpha)}{\mu^{\frac{k(q-p)+p}{q}}} &= \lim_{j \rightarrow +\infty} \frac{S_{\mu_j}(\alpha)}{\mu_j^{\frac{k(q-p)+p}{q}}} \\ &= \lim_{j \rightarrow +\infty} \iint_{\Omega} |(\nabla_x v_j, \mu_j^{-1}\nabla_y v_j)|^p + |v_j|^p \, dx \, dy \\ &\geq \iint_{\Omega} |(\nabla_x v, w)|^p + |v|^p \, dx \, dy \\ &\geq \mathcal{H}^k(\Omega_2) \|v\|_{W^{1,p}(\Omega_1)}^p, \end{aligned}$$

and, by (5.26), we get

$$\liminf_{\mu \rightarrow 0^+} \frac{S_{\mu}(\alpha)}{\mu^{\frac{k(q-p)+p}{q}}} \geq \frac{\mathcal{H}^k(\Omega_2)}{\mathcal{H}^{k-1}(\partial\Omega_2)^{\frac{p}{q}}} \frac{\|v\|_{W^{1,p}(\Omega_1)}^p}{\|v\|_{L^q(\Omega_1)}^{\frac{p}{q}}}. \tag{5.27}$$

Lastly, by (5.2), Lemma 5.1 and since  $u_{\mu_j}$  is an extremal for  $S_{\mu_j}(\alpha)$  for all  $j \in \mathbb{N}$ , we have that

$$\alpha \mathcal{H}^n(\Omega_1) \mathcal{H}^{k-1}(\partial\Omega_2) \leq \mathcal{H}^{N-1}(\{v_j = 0\} \cap (\Omega_1 \times \partial\Omega_2)) + \mu_j \mathcal{H}^{N-1}(\{v_j = 0\} \cap (\partial\Omega_1 \times \Omega_2))$$

for all  $j \in \mathbb{N}$ . Then, using (5.25), we get that

$$\begin{aligned} \alpha \mathcal{H}^n(\Omega_1) \mathcal{H}^{k-1}(\partial\Omega_2) &\leq \limsup_{j \rightarrow +\infty} \mathcal{H}^{N-1}(\{v_j = 0\} \cap (\Omega_1 \times \partial\Omega_2)) \\ &\leq \mathcal{H}^{N-1}(\{v = 0\} \cap (\Omega_1 \times \partial\Omega_2)) \\ &= \mathcal{H}^{N-1}(\{v = 0\} \cap \Omega_1 \times \partial\Omega_2) \\ &= \mathcal{H}^n(\{v = 0\} \cap \Omega_1) \mathcal{H}^{k-1}(\partial\Omega_2). \end{aligned}$$

Thus,

$$\alpha \mathcal{H}^n(\Omega_1) \leq \mathcal{H}^n(\{v = 0\} \cap \Omega_1),$$

and  $v$  is an admissible function in the characterization of  $\mathbb{S}(\alpha)$ . Then, using (5.18) and (5.27), we have that

$$\frac{\mathcal{H}^k(\Omega_2)}{\mathcal{H}^{k-1}(\partial\Omega_2)^{\frac{p}{q}}} \mathbb{S}(\alpha) \leq \liminf_{\mu \rightarrow 0^+} \frac{S_\mu(\alpha)}{\mu^{\frac{k(q-p)+p}{q}}} \leq \limsup_{\mu \rightarrow 0^+} \frac{S_\mu(\alpha)}{\mu^{\frac{k(q-p)+p}{q}}} \leq \frac{\mathcal{H}^k(\Omega_2)}{\mathcal{H}^{k-1}(\partial\Omega_2)^{\frac{p}{q}}} \mathbb{S}(\alpha).$$

The proof is now complete.  $\square$

### 5.2. The case $n = 1$

When the limit problem is one-dimensional we can give a more precise description of the situation. So in this subsection we consider the case  $\Omega_1 = (a, b) \subset \mathbb{R}$ , an interval.

In [13] the following theorem regarding the limit problem for  $n = 1$  is proved:

**Theorem 5.6.** (See [13, Theorem 1.2].) *The optimal limit constant  $\mathbb{S}(\alpha)$  is attained only for a hole  $A^* = (a, a + \alpha(b - a))$  or  $A^* = (b - \alpha(b - a), b)$ , that is the best hole is an interval concentrated on one side of the interval  $(a, b)$ . Moreover, the optimal limit constant is given by*

$$\mathbb{S}(\alpha) = \frac{(2\pi)^p(p - 1)}{(2\alpha(b - a)p \sin(\frac{\pi}{p}))^p} + 1.$$

As a consequence of this theorem, we have the following corollary on the approximate shape and location of optimal boundary holes:

**Corollary 5.7.** *For  $\mu$  small enough the best boundary hole  $\Gamma_\mu$  for the domain  $\Omega_\mu = (a, b) \times \mu\Omega_2$  with measure  $\mathcal{H}^{N-1}(\Gamma_\mu) = \alpha\mathcal{H}^{N-1}(\partial\Omega_\mu)$  looks like  $\Gamma_\mu \simeq (a, a + \alpha(b - a)) \times \partial\mu\Omega_2$  or like  $\Gamma_\mu \simeq (b - \alpha(b - a), b) \times \partial\mu\Omega_2$ .*

### 5.3. General geometries

We finish this section by observing that, once the product case is studied, the extension of our results to more general domains  $\Omega$  in  $\mathbb{R}^N$  than a product is done by a standard procedure. Cf. with [9,13].

So, in this case we let  $\Omega_\mu = \{(x, \mu y) : (x, y) \in \Omega\}$ .

We have the following

**Theorem 5.8.** *Let  $\Omega$  be a bounded and Lipschitz domain in  $\mathbb{R}^N$ . Let  $\Omega_x$  be the  $x$ -section of  $\Omega$  and  $P(\Omega)$  be the projection of  $\Omega$  onto the  $x$  variable, i.e.*

$$\Omega_x := \{y \in \mathbb{R}^k : (x, y) \in \Omega\} \quad \text{and} \quad P(\Omega) := \{x \in \mathbb{R}^n : \Omega_x \neq \emptyset\}.$$

Then, if we call  $\rho(x) = \mathcal{H}^k(\Omega_x)$  and  $\beta(x) = \mathcal{H}^{k-1}(\partial\Omega_x)$  we have that

$$\lim_{\mu \rightarrow 0^+} \frac{S_\mu(\alpha)}{\mu^{\frac{k(q-p)+p}{q}}} = \mathbb{S}(\alpha, \rho, \beta),$$

where

$$\mathbb{S}(\alpha, \rho, \beta) = \inf \left\{ \frac{\int_{P(\Omega)} (|\nabla v|^p + |v|^p) \rho(x) \, dx}{\left( \int_{P(\Omega)} |v|^q \beta(x) \, dx \right)^{\frac{p}{q}}} : v \in A(\alpha) \right\}$$

with

$$A(\alpha) = \{ v \in W^{1,p}(P(\Omega), \rho) : \mathcal{H}^n(\{x \in P(\Omega) : v(x) = 0\}) \geq \alpha \mathcal{H}^n(P(\Omega)) \}.$$

Here  $W^{1,p}(P(\Omega), \rho)$  is the weighted Sobolev space,

$$W^{1,p}(P(\Omega), \rho) = \left\{ v : P(\Omega) \rightarrow \mathbb{R} : \int_{P(\Omega)} (|\nabla v|^p + |v|^p) \rho(x) \, dx < +\infty \right\}.$$

**Proof.** Once the product case is studied, the extension to general geometries is analog to Theorem 1.1 in [9]. See also Theorem 1.3 in [13]. We omit the details.  $\square$

### 6. Shape derivative

In this section, we are interested in the computation of the derivative of the set function  $S(\cdot)$  with respect to regular deformations of the set. The formula obtained in this way could be used in the (numerical) computation of optimal boundary holes. This approach has been used with relevant success in similar problems. See [3,8,17,22] and references therein.

Since the domains of  $S(\cdot)$  are sets contained at the boundary  $\partial\Omega$  which is a manifold of codimension one, we must take deformations of sets, which stay in  $\partial\Omega$ .

We begin describing the kind of variations we are going to consider. Let  $V : \mathbb{R}^N \rightarrow \mathbb{R}^N$  be a Lipschitz field such that,  $V \cdot \nu = 0$  on  $\partial\Omega$ , where  $\nu$  is the outer unit normal vector to  $\partial\Omega$ , and

$$\text{spt}(V) \subset \Omega_\delta := \{x \in \mathbb{R}^N : \text{dist}(x, \partial\Omega) < \delta\}$$

for some  $\delta > 0$  small, where  $\text{spt}(V)$  is the support of  $V$ .

Now, we consider the flow associated to the field  $V$ . Let  $\Phi : [0, \infty) \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ , satisfying

$$\frac{d}{dt} \Phi_t(x) = V(\Phi_t(x)), \quad \Phi_0(x) = x,$$

where  $\Phi_t(\cdot) \equiv \Phi(t, \cdot)$ .

It is not difficult to see that, for each  $t$  fixed,  $\Phi_t$  is a diffeomorphism. Indeed, by construction of the flow,  $\Phi_t$  is invertible with inverse given by  $\Phi_{-t}$ . In [17], the following asymptotic formulas were proved

$$\begin{aligned} \Phi_t(x) &= x + tV(x) + o(t), \\ D\Phi_t(x) &= Id + tDV(x) + o(t), \\ D\Phi_t(x)^{-1} &= Id - tDV(x) + o(t), \\ J\Phi_t(x) &= 1 + t \operatorname{div} V(x) + o(t), \\ J_\tau \Phi_t(x) &= 1 + t \operatorname{div}_\tau V(x) + o(t), \end{aligned}$$

for all  $x \in \mathbb{R}^N$ , where  $J\Phi_t$  is the Jacobian of the flow and  $\operatorname{div}_\tau$  denotes the tangential component of the divergence operator.

So, given  $\Gamma \subset \partial\Omega$ , we are allowed to define

$$\Gamma_t := \Phi_t(\Gamma) \subset \partial\Omega, \tag{6.1}$$

and

$$s(t) := S(\Gamma_t). \tag{6.2}$$

Observe that  $s(0) = S(\Gamma)$ .

**Remark 6.1.** By construction, the flow preserves the topology of the initial domain. Therefore, if  $\Gamma$  is a connected set, then  $\Gamma_t$  will be also connected. In fact, this is one of the characteristics of the *shape derivative*, opposite, for instance, to the *topological derivative*, see [1,2,7,18], etc.

Our first result of this section shows that,  $s(t)$  is continuous with respect to  $t$  at  $t = 0$ .

**Theorem 6.2.** *With the previous notation,*

$$\lim_{t \rightarrow 0^+} s(t) = S(\Gamma).$$

**Proof.** Let  $u \in \mathcal{X}_\Gamma$  and we consider  $v = u \circ \Phi_t^{-1} \in \mathcal{X}_{\Gamma_t}$ . By the change of variables formula, we have

$$\int_{\Omega} |v|^p \, dx = \int_{\Omega} |u|^p \, dx + t \int_{\Omega} |u|^p \, dx + o(t),$$

and

$$\int_{\Omega} |\nabla v|^p \, dx = \int_{\Omega} |\nabla u|^p \, dx + t \int_{\Omega} (|\nabla u|^p \operatorname{div} V - p|\nabla u|^{p-2} \langle \nabla u, DV^T \nabla u^T \rangle) \, dx + o(t).$$

Then,

$$\int_{\Omega} |\nabla v|^p + |v|^p \, dx = \int_{\Omega} |\nabla u|^p + |u|^p \, dx + tR(u) + o(t),$$

where

$$R(u) = \int_{\Omega} (|u|^p + |\nabla u|^p) \operatorname{div} V \, dx - p \int_{\Omega} |\nabla u|^{p-2} \langle \nabla u, DV^T \nabla u^T \rangle \, dx.$$

On the other hand, by the change of variables formula on manifolds, see [17], we obtain

$$\int_{\partial\Omega} |v|^q \, d\mathcal{H}^{N-1} = \int_{\partial\Omega} |u|^q \, d\mathcal{H}^{N-1} + t \int_{\partial\Omega} |u|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + o(t).$$

Then,

$$\begin{aligned}
 s(t) &\leq \frac{\int_{\Omega} |\nabla v|^p + |v|^p \, dx}{\left(\int_{\partial\Omega} |v|^q \, d\mathcal{H}^{N-1}\right)^{\frac{p}{q}}} \\
 &= \frac{\int_{\Omega} |\nabla u|^p + |u|^p \, dx + tR(u) + o(t)}{\left(\int_{\partial\Omega} |u|^q \, d\mathcal{H}^{N-1} + t \int_{\partial\Omega} |u|^q \operatorname{div}_{\tau} V \, d\mathcal{H}^{N-1} + o(t)\right)^{\frac{p}{q}}}, \tag{6.3}
 \end{aligned}$$

and therefore

$$\limsup_{t \rightarrow 0^+} s(t) \leq \frac{\int_{\Omega} |\nabla u|^p + |u|^p \, dx}{\left(\int_{\partial\Omega} |u|^q \, d\mathcal{H}^{N-1}\right)^{\frac{p}{q}}} \quad \forall u \in \mathcal{X}_{\Gamma}.$$

Then

$$\limsup_{t \rightarrow 0^+} s(t) \leq S(\Gamma). \tag{6.4}$$

Now, let  $\{t_n\}_{n \in \mathbb{N}}$  such that  $t_n \rightarrow 0^+$  as  $n \rightarrow \infty$  and

$$\liminf_{t \rightarrow 0^+} s(t) = \lim_{n \rightarrow \infty} s(t_n). \tag{6.5}$$

For each  $n \in \mathbb{N}$ , let  $v_n$  be a positive normalized extremal of  $s(t_n)$ , i.e.  $v_n \in \mathcal{X}_{\Gamma_n}$ ,  $v_n > 0$  in  $\Omega$ ,  $\|v_n\|_{L^q(\partial\Omega)} = 1$  and

$$s(t_n) = \int_{\Omega} |\nabla v_n|^p + |v_n|^p \, dx. \tag{6.6}$$

Using (6.4) and (6.5), we have that  $\{v_n\}_{n \in \mathbb{N}}$  is bounded in  $W^{1,p}(\Omega)$  and therefore there exist  $u \in W^{1,p}(\Omega)$  and some subsequence of  $\{v_n\}_{n \in \mathbb{N}}$  (still denote  $\{v_n\}_{n \in \mathbb{N}}$ ) such that

$$v_n \rightharpoonup u \quad \text{weakly in } W^{1,p}(\Omega), \tag{6.7}$$

$$v_n \rightarrow u \quad \text{strongly in } L^p(\Omega), \tag{6.8}$$

$$v_n \rightarrow u \quad \text{strongly in } L^q(\partial\Omega). \tag{6.9}$$

Then,  $u \geq 0$  and  $\|u\|_{L^q(\partial\Omega)} = 1$  and

$$\liminf_{t \rightarrow 0^+} s(t) \geq \int_{\Omega} |\nabla u|^p + |u|^p \, dx.$$

On the other hand, since  $\Phi_{-t} \rightarrow Id$  in the  $C^1$  topology when  $t \rightarrow 0$  and using (6.9), we have

$$\int_{\partial\Omega} u \chi_{\Gamma} \, d\mathcal{H}^{N-1} = 0$$

and therefore  $u \in \mathcal{X}_\Gamma$ . Then, using (6.4)

$$S(\Gamma) \leq \int_{\Omega} |\nabla u|^p + |u|^p \, dx \leq \liminf_{t \rightarrow 0^+} s(t) \leq \limsup_{t \rightarrow 0^+} s(t) \leq S(\Gamma).$$

Thus,

$$\lim_{t \rightarrow 0^+} s(t) = S(\Gamma).$$

The proof is now completed.  $\square$

**Remark 6.3.** Observe that, in the above prove, we really have that  $v_n \rightarrow u$  strongly in  $W^{1,p}(\Omega)$  when  $n \rightarrow \infty$  because  $\|v_n\|_{W^{1,p}(\Omega)} \rightarrow \|u\|_{W^{1,p}(\Omega)}$  when  $n \rightarrow \infty$  and by (6.7).

Now we arrive at the main result of this section.

**Theorem 6.4.** *If  $\Gamma \subset \partial\Omega$  is a positive  $\mathcal{H}^{N-1}$ -measurable subset, we have that  $s(t)$  is differentiable at  $t = 0$  and*

$$\frac{ds}{dt}(0) = -\frac{p}{q} S(\Gamma) \int_{\partial\Omega} |u|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + R(u), \tag{6.10}$$

where

$$R(u) = \int_{\Omega} (|u|^p + |\nabla u|^p) \operatorname{div} V \, dx - p \int_{\Omega} |\nabla u|^{p-2} \langle \nabla u, DV^T \nabla u^T \rangle \, dx$$

and  $u$  is an extremal of  $S(\Gamma)$ .

**Proof.** Let  $u$  be a positive normalized extremal of  $S(\Gamma)$ . Then, using (6.3), we have that

$$s(t) \leq \frac{S(\Gamma) + tR(u) + o(t)}{(1 + t \int_{\partial\Omega} |u|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + o(t))^{\frac{p}{q}}}.$$

Thus, for all  $t > 0$

$$\begin{aligned} \frac{s(t) - S(\Gamma)}{t} &\leq \frac{S(\Gamma) \left( 1 - (1 + t \int_{\partial\Omega} |u|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + o(t))^{\frac{p}{q}} \right)}{t \left( 1 + t \int_{\partial\Omega} |u|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + o(t) \right)^{\frac{p}{q}}} \\ &\quad + \frac{R(u) + o(1)}{\left( 1 + t \int_{\partial\Omega} |u|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + o(t) \right)^{\frac{p}{q}}}. \end{aligned}$$

Therefore

$$\limsup_{t \rightarrow 0^+} \frac{s(t) - S(\Gamma)}{t} \leq -\frac{p}{q} S(\Gamma) \int_{\partial\Omega} |u|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + R(u). \tag{6.11}$$

On other hand, let  $\{t_n\}_{n \in \mathbb{N}}$  be a positive sequence such that  $t_n \rightarrow 0^+$  when  $n \rightarrow \infty$ , and

$$\liminf_{t \rightarrow 0^+} \frac{s(t) - S(\Gamma)}{t} = \lim_{n \rightarrow \infty} \frac{s(t_n) - S(\Gamma)}{t_n}.$$

Observe that, by Lemma 6.2, we have that  $s(t_n) \rightarrow S(\Gamma)$ . We can now proceed analogously to the proof of Lemma 6.2, and we find a subsequence of  $\{t_n\}_{n \in \mathbb{N}}$  (still denote  $\{t_n\}_{n \in \mathbb{N}}$ ) such that

$$v_n \rightarrow u \text{ strongly in } W^{1,p}(\Omega),$$

where  $v_n$  is a positive normalized extremal of  $s(t_n)$  for all  $n \in \mathbb{N}$  and  $u$  is a positive normalized extremal of  $S(\Gamma)$ , see also Remark 6.3.

Thus, taking  $u_n = v_n \circ \Phi_{t_n} \in W_r^{1,p}(\Omega)$ , we get

$$S(\Gamma) \leq \frac{s(t_n) - t_n R(v_n) + o(t_n)}{(1 - t_n \int_{\partial\Omega} |v_n|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + o(t_n))^{\frac{p}{q}}}.$$

Then

$$\begin{aligned} \frac{s(t_n) - S(\Gamma)}{t_n} &\geq \frac{s(t_n) (1 - t_n \int_{\partial\Omega} |c_n|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + o(t_n))^{\frac{p}{q}} - 1}{t_n (1 - t_n \int_{\partial\Omega} |v_n|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + o(t_n))^{\frac{p}{q}}} \\ &\quad + \frac{R(v_n) + o(1)}{(1 - t_n \int_{\partial\Omega} |v_n|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + o(t_n))^{\frac{p}{q}}}. \end{aligned}$$

Therefore

$$\begin{aligned} \liminf_{t \rightarrow 0^+} \frac{s(t) - S(\Gamma)}{t} &= \lim_{n \rightarrow \infty} \frac{s(t_n) - S(\Gamma)}{t_n} \\ &\geq -\frac{p}{q} S(\Gamma) \int_{\partial\Omega} |u|^q \operatorname{div}_\tau V \, d\mathcal{H}^{N-1} + R(u). \end{aligned} \tag{6.12}$$

Thus, by (6.11) and (6.12), we have that  $s(t)$  is differentiable at  $t = 0$  and (6.10) holds.  $\square$

**Remark 6.5.** One observes that, we do not need in our approach the derivative of the eigenfunctions.

**Remark 6.6.** It would be desirable to obtain a simplification of formula (6.10). In many problems (cf. [8,17,22], etc.) this can be done by using, in an appropriate way, the equation satisfied by  $u$ . In our case, the obstruction we have encountered in order to do that, is the lack of regularity of  $u$  at the boundary. A similar problem was found in [3] where the authors attempt to overcome this difficulty by working on a subset  $\Omega_\delta \subset \Omega$  and then passing to the limit (however, the results are not completely satisfactory). In our case, since we cannot control the normal derivative of  $u$  in  $\Omega_\delta$ , this approach does not seem to be feasible.

**Acknowledgments**

This work was partially supported by project PROSUL (CNPq-CONICET) No. 490329/2008-0. J. Fernández Bonder and Leandro Del Pezzo were also partially supported by Universidad de Buenos Aires under grant X078, by ANPCyT PICT No. 2006-290 and CONICET (Argentina) PIP 5478/1438.



Wladimir Neves was also partially supported by FAPERJ through the grant E-26/111.564/2008 entitled Analysis, Geometry and Applications and by Pronex–FAPERJ through the grant E-26/110.560/2010 entitled Nonlinear Partial Differential Equations.

## References

- [1] Grégoire Allaire, Frédéric de Gournay, François Jouve, Anca-Maria Toader, Structural optimization using topological and shape sensitivity via a level set method, *Control Cybernet.* 34 (1) (2005) 59–80. MR 2211063 (2006j:49073).
- [2] Cristian Barbarosie, Anca-Maria Toader, Shape and topology optimization for periodic problems. I. The shape and the topological derivative, *Struct. Multidiscip. Optim.* 40 (1–6) (2010) 381–391. MR 2565077.
- [3] Almut Burchard, Jochen Denzler, On the geometry of optimal windows, with special focus on the square, *SIAM J. Math. Anal.* 37 (6) (2006) 1800–1827 (electronic).
- [4] L.M. Del Pezzo, Optimization problem for extremals of the trace inequality in domains with holes, *Commun. Contemp. Math.* 12 (4) (2010) 569–586.
- [5] Jochen Denzler, Windows of given area with minimal heat diffusion, *Trans. Amer. Math. Soc.* 351 (2) (1999) 569–580.
- [6] Lawrence C. Evans, Ronald F. Gariepy, *Measure Theory and Fine Properties of Functions*, Stud. Adv. Math., CRC Press, Boca Raton, FL, 1992.
- [7] Raúl A. Feijóo, Antonio A. Novotny, Edgardo Taroco, Claudio Padra, The topological derivative for the Poisson's problem, *Math. Models Methods Appl. Sci.* 13 (12) (2003) 1825–1844. MR 2032213 (2004m:49105).
- [8] Julián Fernández Bonder, Pablo Groisman, Julio D. Rossi, Optimization of the first Steklov eigenvalue in domains with holes: a shape derivative approach, *Ann. Mat. Pura Appl.* (4) 186 (2) (2007) 341–358.
- [9] Julián Fernández Bonder, Sandra Martínez, Julio D. Rossi, The behavior of the best Sobolev trace constant and extremals in thin domains, *J. Differential Equations* 198 (1) (2004) 129–148. MR 2037752 (2005f:35060).
- [10] Julián Fernández Bonder, Julio D. Rossi, Existence results for the  $p$ -Laplacian with nonlinear boundary conditions, *J. Math. Anal. Appl.* 263 (1) (2001) 195–223.
- [11] Julián Fernández Bonder, Julio D. Rossi, Asymptotic behavior of the best Sobolev trace constant in expanding and contracting domains, *Commun. Pure Appl. Anal.* 1 (3) (2002) 359–378.
- [12] Julián Fernández Bonder, Julio D. Rossi, On the existence of extremals for the Sobolev trace embedding theorem with critical exponent, *Bull. Lond. Math. Soc.* 37 (1) (2005) 119–125. MR 2105826 (2006a:35079).
- [13] Julián Fernández Bonder, Julio D. Rossi, Carola-Bibiane Schönlieb, An optimization problem related to the best Sobolev trace constant in thin domains, *Commun. Contemp. Math.* 10 (5) (2008) 633–650. MR 2446893 (2010c:46080).
- [14] Julián Fernández Bonder, Julio D. Rossi, Noemi Wolanski, Regularity of the free boundary in an optimization problem related to the best Sobolev trace constant, *SIAM J. Control Optim.* 44 (5) (2005) 1614–1635 (electronic).
- [15] Julián Fernández Bonder, Julio D. Rossi, Noemi Wolanski, On the best Sobolev trace constant and extremals in domains with holes, *Bull. Sci. Math.* 130 (7) (2006) 565–579.
- [16] Julián Fernández Bonder, Nicolas Saintier, Estimates for the Sobolev trace constant with critical exponent and applications, *Ann. Mat. Pura Appl.* (4) 187 (4) (2008) 683–704. MR 2413374 (2009h:35075).
- [17] Antoine Henrot, Michel Pierre, *Variation et optimisation de formes : Une analyse géométrique*, Mathématiques & Applications (Mathematics & Applications), vol. 48, Springer, Berlin, 2005.
- [18] Mohamed Iguernane, Serguei A. Nazarov, Roche Jean-Rodolphe, Jan Sokolowski, Katarzyna Szulc, Topological derivatives for semilinear elliptic equations, *Int. J. Appl. Math. Comput. Sci.* 19 (2) (2009) 191–205. MR 2541192 (2010g:49073).
- [19] Bernhard Kawohl, *Rearrangements and Convexity of Level Sets in PDE*, Lect. Notes in Math., vol. 1150, Springer-Verlag, Berlin, 1985.
- [20] Gary M. Lieberman, Boundary regularity for solutions of degenerate elliptic equations, *Nonlinear Anal.* 12 (11) (1988) 1203–1219.
- [21] Gary M. Lieberman, The natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva for elliptic equations, *Comm. Partial Differential Equations* 16 (2–3) (1991) 311–361. MR 1104103 (92c:35041).
- [22] Édouard Oudet, Numerical minimization of eigenmodes of a membrane with respect to the domain, *ESAIM Control Optim. Calc. Var.* 10 (3) (2004) 315–330 (electronic).
- [23] Emanuel Sperner Jr., Spherical symmetrization and eigenvalue estimates, *Math. Z.* 176 (1) (1981) 75–86.
- [24] J.L. Vázquez, A strong maximum principle for some quasilinear elliptic equations, *Appl. Math. Optim.* 12 (3) (1984) 191–202.