

On p -compact mappings and the p -approximation property[☆]

Silvia Lassalle, Pablo Turco^{*}

Departamento de Matemática, Pab I, Facultad de Cs. Exactas y Naturales, Universidad de Buenos Aires, (1428) Buenos Aires, Argentina
IMAS, CONICET, Argentina

ARTICLE INFO

Article history:

Received 2 November 2011

Available online 4 January 2012

Submitted by Richard M. Aron

Keywords:

p -Compact sets

Holomorphic mappings

Approximation properties

ABSTRACT

The notion of p -compact sets arises naturally from Grothendieck's characterization of compact sets as those contained in the convex hull of a norm null sequence. The definition, due to Sinha and Karn (2002), leads to the concepts of p -approximation property and p -compact operators (which form an ideal with its ideal norm κ_p). This paper examines the interaction between the p -approximation property and certain space of holomorphic functions, the p -compact analytic functions. In order to understand these functions we define a p -compact radius of convergence which allows us to give a characterization of the functions in the class. We show that p -compact holomorphic functions behave more like nuclear than compact maps. We use the ϵ -product of Schwartz, to characterize the p -approximation property of a Banach space in terms of p -compact homogeneous polynomials and in terms of p -compact holomorphic functions with range on the space. Finally, we show that p -compact holomorphic functions fit into the framework of holomorphy types which allows us to inspect the κ_p -approximation property. Our approach also allows us to solve several questions posed by Aron, Maestre and Rueda (2010).

© 2012 Elsevier Inc. All rights reserved.

0. Introduction

In the theory of Banach spaces (or more precisely, of infinite dimensional locally convex spaces), three concepts appear systematically related since the foundational articles by Grothendieck [21] and Schwartz [29]. We are referring to compact sets, compact operators and the approximation property. A Banach space E has the approximation property whenever the identity map can be uniformly approximated by finite rank operators on compact sets. Equivalently, if $E' \otimes E$, the subspace of finite rank operators, is dense in $\mathcal{L}_c(E; E)$, the space of continuous linear operators considered with the uniform convergence on compact sets. Another classical reformulation states that E has the approximation property if $F' \otimes E$ is dense in $\mathcal{K}(F; E)$, the space of compact operators, for all Banach spaces F . It was not until 1972 that Enflo provided us with the first example of a Banach space without the approximation property [19]. In the quest of a better understanding of these concepts, important variants of the approximation property have emerged and were intensively studied in relation with different spaces of functions. For the main developments on the subject we refer the reader to [9,23] and the references therein.

Inspired by Grothendieck's result which characterizes relatively compact sets as those contained in the convex hull of a norm null sequence of vectors of the space, Sinha and Karn [30] introduced the concept of relatively p -compact sets. Loosely speaking, these sets are determined by norm p -summable sequences. Associated to relatively p -compact sets we have naturally defined the notions of p -compact operators and the p -approximation property (see definitions below). Since relatively p -compact sets are, in particular, relatively compact, the p -approximation property can be seen as a way to

[☆] This project was supported in part by UBACyT X218 and UBACyT X038.

^{*} Corresponding author.

E-mail addresses: slassall@dm.uba.ar (S. Lassalle), paturco@dm.uba.ar (P. Turco).

weaken the approximation property. These three concepts were first studied by Sinha and Karn [30,31] and, more recently, several authors continued the research on this subject [10,12–14,20].

This paper examines the interaction between the p -approximation property and the class of p -compact holomorphic functions. The connection between the approximation property and the space of holomorphic functions is not without precedent. The pioneer article on this topic is due to Aron and Schottenloher [5], who prove that a Banach space E has the approximation property if and only if $(\mathcal{H}(E), \tau_0)$, the space of the entire functions with the compact open topology, has the approximation property. Since then, many authors studied the approximation property for spaces of holomorphic functions in different contexts, see for instance [6,7,17,18,25]. Recently, Aron, Maestre and Rueda [2] prove that E has the p -approximation property if and only if $(\mathcal{H}(E), \tau_{0p})$ has the approximation property, here τ_{0p} denotes the topology of the uniform convergence on p -compact sets. The relation between the approximation property and holomorphic mappings was studied in detail in [5], where the class of compact holomorphic functions plays a crucial role.

The article is organized as follows. In the first section we fix the notation and state some basic results on p -compact mappings. In Section 2 we study the behavior of p -compact homogeneous polynomials which can be considered as a polynomial Banach ideal with a natural norm denoted by κ_p . We use the standard linearization of polynomials via the symmetric projective tensor product and show that any p -compact homogeneous polynomial P factors through a quotient of ℓ_1 and a quotient of ℓ_q , $\frac{1}{p} + \frac{1}{q} = 1$, and give a characterization of the κ_p -norm of P in terms of these factorizations. We also prove that the Aron–Berner extension preserves isometrically the class of p -compact polynomials and exhibit an isometric relationship between the adjoint of p -compact polynomials and quasi- p -nuclear operators.

Section 3 is devoted to the study of p -compact holomorphic mappings. Since p -compact functions are compact, we pay special attention to the results obtained by Aron and Schottenloher [5], where the authors prove that any holomorphic function is compact if and only if each polynomial of its Taylor series expansion at 0 is compact [5, Proposition 3.4]. Then, Aron, Maestre and Rueda [2, Proposition 3.5] show that each component of the Taylor series expansion of a p -compact holomorphic mapping has to be also p -compact and wonder if there is reciprocal result. We define a natural p -compact radius of convergence and, in Proposition 3.4, we give a characterization of this type of functions. Surprisingly, we found that p -compact holomorphic functions behave more like nuclear than compact mappings. We show this feature with two examples. Example 3.7 shows that Proposition 3.4 cannot be improved and also that it is possible to find an entire function whose polynomials at 0 are p -compact but the function fails to be p -compact at 0, which answers by the negative the question posed in [2, Problem 5.2]. In Example 3.8 we construct an entire function on ℓ_1 which is p -compact on the open unit ball, but it fails to be p -compact at the first element of the canonical basis of ℓ_1 , giving an answer to [2, Problem 5.1].

We apply the results of Sections 2 and 3 to study the p -approximation property in Section 4. We characterize the p -approximation property of a Banach space in terms of p -compact homogeneous polynomials with range on the space. Our proof requires the notion of the ϵ -product of Schwartz [29]. We show that a Banach space E has the p -approximation property if and only if p -compact homogeneous polynomials with range on E can be uniformly approximated by finite rank polynomials. We also give the analogous result for p -compact holomorphic mappings endowed with the Nachbin topology, Proposition 4.7.

The final section is dedicated to the p -compact holomorphic mappings within the framework of holomorphy types, concept introduced by Nachbin [26,27]. This allows us to inspect the κ_p -approximation property introduced, in [13], in the spirit of [5, Theorem 4.1].

1. Preliminaries

Throughout this paper E and F are Banach spaces. We denote by B_E the closed unit ball of E , by E' its topological dual, and by $\ell_p(E)$ the Banach space of the p -summable sequences of elements of E , endowed with its natural norm. Also, $c_0(E)$ denotes the space of null sequences of E endowed with the supremum norm. Following [30], we say that a subset $K \subset E$ is relatively p -compact, $1 \leq p \leq \infty$, if there exists a sequence $(x_n)_n \in \ell_p(E)$ so that K is contained in the closure of $\{\sum \alpha_n x_n : (\alpha_n)_n \in B_{\ell_q}\}$, where B_{ℓ_q} denotes the closed unit ball of ℓ_q , with $\frac{1}{p} + \frac{1}{q} = 1$. We denote this set by $p\text{-co}\{x_n\}$ and its closure by $\overline{p\text{-co}\{x_n\}}$. With $p = \infty$ the definition of compact sets is recovered. When $p = 1$, the 1-convex hull is obtained by considering coefficients in B_{ℓ_∞} or, if necessary, with some extra work by coefficients in B_{c_0} , see [14, Remark 3.3].

Since the sequence $(x_n)_n$ in the definition of a relatively p -compact set K converges to zero, any p -compact set is compact. Such a sequence is not unique, then we may consider

$$m_p(K; E) = \inf\{\|(x_n)_n\|_p : K \subset p\text{-co}\{x_n\}\}$$

which measures the size of K as a p -compact set of E . If $K \subset E$ is not p -compact, we write $m_p(K; E) = \infty$. For simplicity, along this work we write $m_p(K)$ instead of $m_p(K; E)$. When $K \subset p\text{-co}\{x_n\}$, $(x_n)_n \in \ell_p(E)$, any $x \in K$ has the form $x = \sum \alpha_n x_n$ for some $(\alpha_n)_n \in B_{\ell_q}$. By Hölder's inequality, $\|x\| \leq \|(x_n)_n\|_{\ell_p(E)}$ and $\|x\| \leq m_p(K)$, for all $x \in K$. We use without any further mention the following equalities: $m_p(K) = m_p(\overline{K}) = m_p(\Gamma(K))$, where $\Gamma(K)$ denotes the absolutely convex hull of K , a relatively p -compact set.

The space of linear bounded operators from E to F is denoted by $\mathcal{L}(E; F)$ and $E' \otimes F$ denotes its subspace of finite rank operators. As in [30], we say that an operator $T \in \mathcal{L}(E; F)$ is p -compact, $1 \leq p \leq \infty$, if $T(B_E)$ is a relatively p -compact set in F . The space of p -compact operators from E to F is denoted by $\mathcal{K}_p(E; F)$. In $\mathcal{K}_p(E; F)$ there is a natural norm defined as

$$\kappa_p(T) = \inf\{\|(y_n)_n\|_p : (y_n)_n \in \ell_p(F) \text{ and } T(B_E) \subset p\text{-co}\{y_n\}\},$$

for any $T \in \mathcal{K}_p(E; F)$, where κ_∞ coincides with the supremum norm. The pair $(\mathcal{K}_p, \kappa_p)$ is a Banach operator ideal [28] (see also [14]).

The Banach ideal \mathcal{K}_p is associated by duality with the ideal of quasi- p -nuclear operators, introduced and studied by Persson and Pietsch [28]. Recall that an operator $T \in \mathcal{L}(E; F)$ is quasi- p -nuclear if and only if there exists a sequence $(x'_n)_n \subset \ell_p(E')$, such that

$$\|Tx\| \leq \left(\sum_n |x'_n(x)|^p\right)^{\frac{1}{p}},$$

for all $x \in E$ and the quasi- p -nuclear norm of T is given by

$$\nu_p^Q(T) = \inf\left\{\|(x'_n)_n\|_p : \|Tx\|^p \leq \sum_n |x'_n(x)|^p, \forall x \in E\right\}.$$

The space of quasi- p -nuclear operators from E to F is denoted by $\mathcal{QN}_p(E; F)$. The dual relationship is as follows. Given $T \in \mathcal{L}(E; F)$, T is p -compact if and only if its adjoint is quasi- p -nuclear. Also, T is quasi- p -nuclear if and only if its adjoint is p -compact, see [14, Corollary 3.4] and [14, Proposition 3.8].

A mapping $P : E \rightarrow F$ is an m -homogeneous polynomial if there exists a (unique) symmetric m -linear form $\overset{\vee}{P} : \underbrace{E \times \dots \times E}_m \rightarrow F$ such that

$$P(x) = \overset{\vee}{P}(x, \dots, x),$$

for all $x \in E$. The space of m -homogeneous continuous polynomials from E to F is denoted by $\mathcal{P}^m(E; F)$, which is a Banach space considered with the supremum norm

$$\|P\| = \sup\{\|P(x)\| : x \in B_E\}.$$

Given a homogeneous polynomial $P \in \mathcal{P}^m(E; F)$, there are two natural mappings associated to it. The *linearization*, denoted by $L_P \in \mathcal{L}(\otimes_{\pi_s}^m E; F)$, where $\otimes_{\pi_s}^m E$ stands for the completion of the symmetric m -tensor product endowed with the symmetric projective norm. Also we have the polynomial $\bar{P} \in \mathcal{P}^m(E'', F'')$, known as *the Aron-Berner extension of P* [1], which is the extension of P from E to E'' obtained by weak-star density. We have $\|L_P\| \leq \frac{m^m}{m!} \|P\|$ and $\|\bar{P}\| = \|P\|$ [11].

A mapping $f : E \rightarrow F$ is holomorphic at $x_0 \in E$ if there exists a sequence of polynomials $P_m f(x_0) \in \mathcal{P}^m(E; F)$ such that

$$f(x) = \sum_{m=0}^{\infty} P_m f(x_0)(x - x_0),$$

uniformly for all x in some neighborhood of x_0 . We say that $\sum_{m=0}^{\infty} P_m f(x_0)$ is the Taylor series expansion of f at x_0 and that $P_m f(x_0)$ is its m -component of the series at x_0 . A function is said to be holomorphic or entire if it is holomorphic at x for all $x \in E$. The space of entire functions from E to F is denoted by $\mathcal{H}(E; F)$.

We refer the reader to [16,24] for general background on polynomials and holomorphic functions.

2. The p -compact polynomials

We want to understand the behavior of p -compact holomorphic mappings. The definition, due to Aron, Maestre and Rueda [2] was introduced as a natural extension of p -compact operators to the nonlinear case. In [2] the authors show that for any p -compact holomorphic function each m -homogeneous polynomial of its Taylor series expansion must be p -compact. Motivated by this fact we devote this section to the study of polynomials.

Recall that $P \in \mathcal{P}^m(E; F)$ is said to be p -compact, $1 \leq p \leq \infty$, if $P(B_E)$ is relatively p -compact in F . In particular, any p -compact polynomial is compact. We denote by $\mathcal{PK}_p^m(E; F)$ the space of p -compact m -homogeneous polynomials and by $\mathcal{PK}^m(E; F)$ the space of compact polynomials. On $\mathcal{PK}_p^m(E; F)$ we may define

$$\kappa_p(P) = m_p(P(B_E)),$$

which is a norm satisfying that $\|P\| \leq \kappa_p(P)$, for any p -compact homogeneous polynomial P . Also, $(\mathcal{PK}_p^m(E; F), \kappa_p)$ is a polynomial Banach ideal.

Lemma 2.1. *Let E and F be Banach spaces and let $P \in \mathcal{P}^m(E; F)$. The following statements are equivalent.*

- (i) P is p -compact.
- (ii) $L_P : \otimes_{\pi_s}^m E \rightarrow F$, the linearization of P , is a p -compact operator.

Moreover, we have $\kappa_p(P) = \kappa_p(L_P)$.

Proof. To show the equivalence, we appeal to the familiar diagram, where Λ is a norm one homogeneous polynomial ($\Lambda(x) = x^m$) and $P = L_P \Lambda$:

$$\begin{array}{ccc}
 E & \xrightarrow{P} & F \\
 \Lambda \downarrow & \nearrow L_P & \\
 \otimes_{\pi_s}^m E & &
 \end{array}
 \tag{1}$$

Note that the open unit ball of $\otimes_{\pi_s}^m E$ is the absolutely convex hull $\Gamma\{x^m : \|x\| < 1\}$. Then, we have that $P(B_E) \subset \Gamma(\{L_P(x^m) : \|x\| < 1\}) = \Gamma(P(B_E))$. Now, the equality $L_P(\Gamma\{x^m : \|x\| < 1\}) = \Gamma(P(B_E))$ shows that any sequence $(y_n)_n \in \ell_p(F)$ involved in the definition of $\kappa_p(P)$ is also involved in the definition of $\kappa_p(L_P)$ and vice versa, which completes the proof. \square

The behavior of p -compact polynomials is similar to that described for p -compact operators in [20, Proposition 2.9] (see also [10, Theorem 3.1]).

Proposition 2.2. *Let E and F be Banach spaces, $1 \leq p < \infty$, $P \in \mathcal{P}(^m E; F)$. Then, if $\frac{1}{p} + \frac{1}{q} = 1$, the following are equivalent.*

- (i) $P \in \mathcal{P}_{K_p}(^m E; F)$.
- (ii) *There exist subspaces $M \subset \ell_1$ and $N \subset \ell_q$, a compact polynomial $Q \in \mathcal{P}_K(^m E; \ell_q/N)$ and operators $T \in \mathcal{K}_p(\ell_q/N; \ell_1/M)$ and $S \in \mathcal{K}(\ell_1/M; F)$ such that $P = STQ$. In this case*

$$\kappa_p(P) = \inf\{\|S\|\kappa_p(T)\|Q\|\},$$

where the infimum is taken over all the factorizations as above.

Proof. We only have to prove (i) implies (ii). By Lemma 2.1, $P = L_P \Lambda$ where L_P is p -compact and $\kappa_p(P) = \kappa_p(L_P)$. Now, we merge diagram (1) with that of [20, Proposition 2.9] for the operator L_P and obtain

$$\begin{array}{ccccc}
 E & \xrightarrow{P} & & \xrightarrow{\quad} & F \\
 \Lambda \downarrow & & & \nearrow L_P & \\
 \otimes_{\pi_s}^m E & \xrightarrow{R} & \ell_q/N & \xrightarrow{T} & \ell_1/M \\
 & & & & \uparrow S
 \end{array}$$

where the operators R and S are compact and T is p -compact. Then, with $Q = R\Lambda$ we have the factorization desired. Note that $\kappa_p(P) \leq \|S\|\kappa_p(T)\|Q\| \leq \|S\|\kappa_p(T)\|R\|$. By [20, Proposition 2.9], $\kappa_p(L_P) = \inf\{\|S\|\kappa_p(T)\|R\|\}$. Using again Lemma 2.1, the proof is complete. \square

It is shown in [14, Corollary 3.6], that an operator $T : E \rightarrow F$ is p -compact if and only if its bitranspose $T'' : E'' \rightarrow F''$ is p -compact with $\kappa_p(T'') \leq \kappa_p(T)$. In [20, Corollary 2.6], it is proved that, in fact, $\kappa_p(T'') = \kappa_p(T)$ regardless T'' is considered as an operator on F'' or, thanks to the Gantmacher theorem, as an operator on F . This result, allows us to show that the Aron-Berner extension is a κ_p -isometric extension which preserves the ideal of p -compact homogeneous polynomials. Recall that \bar{P} denotes the Aron-Berner extension of P .

Proposition 2.3. *Let E and F be Banach spaces, $1 \leq p < \infty$, and $P \in \mathcal{P}(^m E; F)$. Then P is p -compact if and only if \bar{P} is p -compact. Moreover, $\kappa_p(P) = \kappa_p(\bar{P})$.*

Proof. Clearly, P is p -compact whenever \bar{P} is and also $\kappa_p(P) \leq \kappa_p(\bar{P})$. Now, suppose that P is p -compact. By Lemma 2.1, we can factorize P via its linearization as $P = L_P \Lambda$, with $\|\Lambda\| = 1$ and L_P a p -compact operator. Since $\bar{P} = L'_p \bar{\Lambda}$, applying [20, Corollary 2.6] and Lemma 2.1, we see that \bar{P} is p -compact and $\kappa_p(\bar{P}) \leq \kappa_p(L'_p) = \kappa_p(L_P) = \kappa_p(P)$, which gives the reverse inequality. \square

We finish this section by relating the transpose of p -compact polynomials with quasi- p -nuclear operators. Given an m -homogeneous polynomial $P : E \rightarrow F$ its adjoint is defined as the linear operator $P' : F' \rightarrow \mathcal{P}(^m E)$ given by $P'(y') = y' \circ P$. In [20, Corollary 2.7], it is shown that the transpose of a p -compact linear operator satisfies the equality $\kappa_p(T) = \nu_p^Q(T')$. Since $P' = L'_p$, where L_P is the linearization of P , using Lemma 2.1 we immediately have:

Corollary 2.4. *A homogeneous polynomial $P \in \mathcal{P}({}^m E; F)$ is p -compact if and only if its transpose $P' \in \mathcal{L}(F'; \mathcal{P}({}^m E))$ is quasi- p -nuclear, and $\kappa_p(P) = v_p^Q(P')$.*

When this manuscript was complete we learned that R. Aron and P. Rueda were also been working on p -compact polynomials [3]. They obtained Lemma 2.1 and a nonisometric version of the corollary above.

3. The p -compact holomorphic mappings


In this section we undertake a detailed study of p -compact holomorphic mappings, whose definition recovers the notion of compact holomorphic mappings for $p = \infty$ [2]. Recall that for E and F Banach spaces, $1 \leq p \leq \infty$, a holomorphic function $f : E \rightarrow F$ is said to be p -compact at x_0 if there is a neighborhood V_{x_0} of x_0 , such that $f(V_{x_0}) \subset F$ is a relatively p -compact set. Also, $f \in \mathcal{H}(E; F)$ is said to be p -compact if it is p -compact at x for all $x \in E$. We denote by $\mathcal{H}_{\kappa_p}(E; F)$ the space of p -compact entire functions and by $\mathcal{H}_K(E; F)$ the space of compact holomorphic mappings. For homogeneous polynomials, it is equivalent to be compact (p -compact) at some point of E or to be compact (p -compact) at every point of the space [2,5]. The same property remains valid for compact holomorphic mappings [5, Proposition 3.4] although the situation is very different for p -compact holomorphic functions, $1 \leq p < \infty$. Indeed, we show that p -compact holomorphic mappings, $1 \leq p < \infty$, behave more like nuclear than compact holomorphic functions.

Having in mind that $(\mathcal{P}_{\kappa_p}({}^m E; F), \kappa_p)$ is a polynomial Banach ideal with $\kappa_p(P) = m_p(P(B_E))$, and that all polynomials in the Taylor series expansion of a p -compact holomorphic function at x_0 are p -compact [2, Proposition 3.5], we propose to connect the concepts of p -compact holomorphic mappings and the size of p -compact sets measured by m_p . We start with a simple but useful lemma.

Lemma 3.1. *Let E be a Banach space and consider K_1, K_2, \dots a sequence of relatively p -compact sets in E , $1 \leq p < \infty$. If $\sum_{j=1}^{\infty} m_p(K_j) < \infty$, then the series $\sum_{j=1}^{\infty} x_j$ is absolutely convergent for any choice of $x_j \in K_j$ and the set $K = \{\sum_{j=1}^{\infty} x_j : x_j \in K_j\}$ is relatively p -compact with $m_p(K) \leq \sum_{j=1}^{\infty} m_p(K_j) < \infty$.*

Proof. Note that K is well defined since for $x_j \in K_j$, $\|x_j\| \leq m_p(K_j)$, for all j and $\sum_{j=1}^{\infty} m_p(K_j) < \infty$.

First, suppose that $p > 1$ and fix $\varepsilon > 0$. For each $j \in \mathbb{N}$, we may assume that K_j is nonempty and we may choose $(x_n^j)_n \in \ell_p(E)$ such that $K_j \subset p\text{-co}\{x_n^j : n \in \mathbb{N}\}$ with $\|(x_n^j)_n\|_p \leq m_p(K_j)(1 + \frac{\varepsilon}{2^j} m_p(K_j)^{-1})^{1/p}$. Now, take $\lambda_j = m_p(K_j)^{-1/q}$, where $\frac{1}{p} + \frac{1}{q} = 1$ and define the sequence $(z_k)_k \subset E$ such that each term is of the form $\lambda_j x_n^j$, following the standard order:



Then

$$\begin{aligned} \sum_{k=1}^{\infty} \|z_k\|^p &= \sum_{j=1}^{\infty} \sum_{n=1}^{\infty} \lambda_j^p \|x_n^j\|^p \\ &= \sum_{j=1}^{\infty} m_p(K_j)^{-p/q} \|(x_n^j)_n\|_{\ell_p(E)}^p \\ &\leq \sum_{j=1}^{\infty} m_p(K_j)^{-p/q} m_p(K_j)^p \left(1 + \frac{\varepsilon}{2^j} m_p(K_j)^{-1}\right) \\ &= \sum_{j=1}^{\infty} m_p(K_j) + \varepsilon. \end{aligned}$$

Hence, $(z_k)_k$ belongs to $\ell_p(E)$ and $\|(z_k)_k\|_{\ell_p(E)} \leq (\sum_{j=1}^{\infty} m_p(K_j) + \varepsilon)^{1/p}$.

Now, take $K = \{\sum_{j=1}^{\infty} x_j : x_j \in K_j\}$ and $x \in K$, $x = \sum_{j=1}^{\infty} x_j$ with $x_j \in K_j$. For each $j \in \mathbb{N}$, there exists $(\alpha_n^j)_n \in B_{\ell_q}$ such that $x_j = \sum_{n=1}^{\infty} \alpha_n^j x_n^j$. Then $x = \sum_{j=1}^{\infty} \sum_{n=1}^{\infty} \alpha_n^j x_n^j$, and the series converges absolutely as the partial sums of $|\alpha_n^j| \|x_n^j\|$ are convergent. We may write $x = \sum m_p(K_j)^{1/q} \alpha_n^j \lambda_j x_n^j$ with

$$\sum |m_p(K_j)^{1/q} \alpha_n^j|^q = \sum_{j=1}^{\infty} \sum_{n=1}^{\infty} |\alpha_n^j|^q m_p(K_j) \leq \sum_{j=1}^{\infty} m_p(K_j).$$

Then K is p -compact as $K \subset (\sum_{j=1}^{\infty} m_p(K_j))^{1/q} p\text{-co}\{z_k\}$. Also

$$m_p(K) \leq \left(\sum_{j=1}^{\infty} m_p(K_j) \right)^{1/q} \| (z_k)_k \|_{\ell_p(E)} \leq \left(\sum_{j=1}^{\infty} m_p(K_j) \right)^{1/q} \left(\sum_{j=1}^{\infty} m_p(K_j) + \varepsilon \right)^{1/p}.$$

Letting $\varepsilon \rightarrow 0$, we conclude that $m_p(K) \leq \sum_{j=1}^{\infty} m_p(K_j)$.

With the usual modifications, the case $p = 1$ follows from the above construction considering $\lambda_j = 1$, for all j . \square

Aron, Maestre and Rueda [2, Proposition 3.5] prove that if f is a p -compact holomorphic mapping at some $x_0 \in E$, every homogeneous polynomial of the Taylor series expansion of f at x_0 is p -compact. At the light of the existent characterization for compact holomorphic mappings [5], they also wonder if the converse is true [2, Problem 5.2]. To tackle this question we need to define the p -compact radius of convergence of a function f at $x_0 \in E$.

Definition 3.2. Let E and F be Banach spaces, $f \in \mathcal{H}(E; F)$ and $x_0 \in E$. If $\sum_{m=0}^{\infty} P_m f(x_0)$ is the Taylor series expansion of f at x_0 , we say that

$$r_p(f, x_0) = 1 / \limsup \kappa_p(P_m f(x_0))^{1/m}$$

is the radius of p -compact convergence of f at x_0 , for $1 \leq p < \infty$.

As usual, we understand that whenever $\limsup \kappa_p(P_m f(x_0))^{1/m} = 0$, the radius of p -compact convergence is infinite. Also, if $P_m f(x_0)$ fails to be p -compact for some m , f fails to be p -compact and $r_p(f, x_0) = 0$.

The following lemma is obtained by a slight modification of the generalized Cauchy formula given in the proof of [2, Proposition 3.5], which asserts that if $f \in \mathcal{H}(E; F)$, $x_0 \in E$ and $\varepsilon > 0$, $P_m f(x_0)(B_\varepsilon(0)) \subset \overline{\text{co}}\{f(B_\varepsilon(x_0))\}$, where $B_\varepsilon(x_0)$ stands for the open ball of center x_0 and radius ε . We state the result as it is used in Section 4, also we are interested in measuring the m_p -size of $P_m f(x_0)(V)$ in terms of the m_p -size of $f(x_0 + V)$ for certain absolutely convex open sets $V \subset E$.

Lemma 3.3. Let E and F be Banach spaces, let $x_0 \in E$ and let $V \subset E$ be an absolutely convex open set. Let $f \in \mathcal{H}(E; F)$ whose Taylor series expansion at x_0 is given by $\sum_{m=0}^{\infty} P_m f(x_0)$. Then

- (a) $P_m f(x_0)(V) \subset \overline{\text{co}}\{f(x_0 + V)\}$, for all m .
- (b) If $f(x_0 + V)$ is relatively p -compact then $m_p(P_m f(x_0)(V)) \leq m_p(f(x_0 + V))$, for all m .

Now we are ready to give a characterization of a p -compact analytic function in terms of the polynomials in its Taylor series expansion and the p -compact radius of convergence.

Proposition 3.4. Let E and F be Banach spaces and let $f \in \mathcal{H}(E; F)$ whose Taylor series expansion at x_0 is given by $\sum_{m=0}^{\infty} P_m f(x_0)$. For $1 \leq p < \infty$, the following statements are equivalent.

- (i) f is p -compact at x_0 .
- (ii) $P_m f(x_0) \in \mathcal{P}_{\kappa_p}(^m E; F)$, for all m and $\limsup \kappa_p(P_m f(x_0))^{1/m} < \infty$.

Proof. To prove that (i) implies (ii), take $\varepsilon > 0$ such that $f(B_\varepsilon(x_0))$ is relatively p -compact and $f(x) = \sum_{m=1}^{\infty} P_m f(x_0)(x - x_0)^m$, with uniform convergence in $B_\varepsilon(x_0)$. By [2, Proposition 3.5], $P_m f(x_0)(\varepsilon B_E) \subset \overline{\text{co}}\{f(B_\varepsilon(x_0))\}$ and $P_m f(x_0)$ is p -compact, for all m . Moreover, by the lemma above,

$$\kappa_p(P_m f(x_0)) = m_p(P_m f(x_0)(B_E)) = \frac{1}{\varepsilon^m} m_p(P_m f(x_0)(\varepsilon B_E)) \leq \frac{1}{\varepsilon^m} m_p(\overline{\text{co}}\{f(B_\varepsilon(x_0))\}).$$

It follows that $\limsup \kappa_p(P_m f(x_0))^{1/m} \leq \frac{1}{\varepsilon}$, as we wanted to prove.

Conversely, suppose that $\limsup \kappa_p(P_m f(x_0))^{1/m} = C > 0$ and choose $0 < \varepsilon < r_p(f, x_0)$ such that, for all $x \in B_\varepsilon(x_0)$, $f(x) = \sum_{m=1}^\infty P_m f(x_0)(x - x_0)$, with uniform convergence. Now we have

$$f(B_\varepsilon(x_0)) \subset \left\{ \sum_{m=1}^\infty x_m : x_m \in P_m f(x_0)(\varepsilon B_E) \right\}.$$

By Lemma 3.1, we obtain the result if we prove that $\sum_{m=1}^\infty m_p(P_m f(x_0)(\varepsilon B_E)) < \infty$, which follows from the equality

$$\sum_{m=1}^\infty m_p(P_m f(x_0)(\varepsilon B_E)) = \sum_{m=1}^\infty \varepsilon^m \kappa_p(P_m f(x_0)),$$

and the choice of ε . \square

Remark 3.5. Let f be a p -compact holomorphic mapping at x_0 and let $\sum_{m=0}^\infty P_m f(x_0)$ be its Taylor series expansion at x_0 . Then, if $\varepsilon < r_p(f, x_0)$,

$$m_p(f(B_\varepsilon(x_0))) \leq \sum_{m=1}^\infty m_p(P_m f(x_0)(\varepsilon B_E)),$$

where the right hand series is convergent.

The p -compact radius has the following natural property.

Proposition 3.6. Let E and F be Banach spaces, $1 \leq p < \infty$, and $f \in \mathcal{H}(E; F)$. Suppose that f is p -compact at x_0 with positive p -compact radius $r = r_p(f, x_0)$. Then f is p -compact for all $x \in B_r(x_0)$. Also, if f is p -compact at x_0 with infinite p -compact radius, then f is p -compact at x , for all $x \in E$.

Proof. Without loss of generality, we can assume that $x_0 = 0$. For $r = r_p(f, 0)$, take $x \in E$, $\|x\| < r$. By [26, Proposition 1, p. 26], there exists $\varepsilon > 0$ such that $f(y) = \sum_{m=1}^\infty P_m f(0)(y)$ converges uniformly for all $y \in B_\varepsilon(x)$. We also may assume that $\|x\| + \varepsilon < r$.

As in Proposition 3.4, we have that $f(B_\varepsilon(x)) \subset \{\sum_{m=1}^\infty x_m : x_m \in P_m f(0)(B_\varepsilon(x))\}$. Now, if we prove that $\sum_{m=1}^\infty m_p(P_m f(0)(B_\varepsilon(x))) < \infty$, the result follows from Lemma 3.1. Indeed,

$$\begin{aligned} \sum_{m=1}^\infty m_p(P_m f(0)(B_\varepsilon(x))) &= \sum_{m=1}^\infty (\|x\| + \varepsilon)^m m_p\left(P_m f(0)\left(\frac{1}{\|x\| + \varepsilon} B_\varepsilon(x)\right)\right) \\ &\leq \sum_{m=1}^\infty (\|x\| + \varepsilon)^m m_p(P_m f(0)(B_E)) \\ &= \sum_{m=1}^\infty ((\|x\| + \varepsilon) \kappa_p(P_m f(0))^{1/m})^m. \end{aligned}$$

Since $(\|x\| + \varepsilon)r^{-1} < 1$, the last series is convergent and the claim is proved. \square

We recently learned that R. Aron and P. Rueda defined, in the context of ideals of holomorphic functions [4], a radius of \mathcal{I} -boundedness which for p -compact holomorphic functions coincides with Definition 3.2. With the radius of \mathcal{I} -boundedness they obtained a partial version of Proposition 3.4.

Thanks to the Josefson–Nissenzweig theorem we have, for any Banach spaces E and F , a p -compact holomorphic mapping, $f \in \mathcal{H}_{\kappa_p}(E; F)$, whose p -compact radius of convergence at the origin is finite. It is enough to consider a sequence $(x'_m)_m \subset E'$ with $\|x'_m\| = 1 \forall m \in \mathbb{N}$, and $(x'_m)_m$ point-wise convergent to 0. Then, $f(x) = \sum_{m=1}^\infty x'_m(x)^m$ belongs to $\mathcal{H}(E)$, is 1-compact (hence, p -compact for any $p > 1$) and $r_p(f, 0) = 1$ since $\kappa_p((x'_m)^m) = \|x'_m\| = 1$. The example can be modified to obtain a vector valued holomorphic function with similar properties.

There are two main questions related to p -compact holomorphic functions which were stated as Problem 5.1 and Problem 5.2 by Aron, Maestre and Rueda [2]. Both arise from properties that compact holomorphic functions satisfy. Recall that we may consider compact sets as ∞ -compact sets and compact mappings as ∞ -compact functions, where $\kappa_\infty(P) = \|P\|$, for any compact m -homogeneous polynomial P . Let us consider $f \in \mathcal{H}(E; F)$, by [5, Proposition 3.4] it is known that if f is compact at one point, say at the origin, then f is compact at x for all $x \in E$. Also, if $\sum_{m=0}^\infty P_m f(0)$ is the Taylor series expansion of f at 0, and for each m the homogeneous polynomial $P_m f(0) : E \rightarrow F$ is compact, then f is compact. With Example 3.7 we show that this later result is no longer true for $1 \leq p < \infty$. Note that $\limsup \|P_m\|^{1/m} < \infty$ is fulfilled by the Cauchy inequalities whenever f is compact. Example 3.7 also shows that, in Proposition 3.4, the hypothesis $\limsup \kappa_p(P_m f(x_0))^{1/m} < \infty$ cannot be ruled out. For our purposes, we adapt [15, Example 10].

Example 3.7. For every $1 \leq p < \infty$, there exists a holomorphic function $f \in \mathcal{H}(\ell_1; \ell_p)$ such that for all $m \in \mathbb{N}$, $P_m f(0)$ is p -compact, but f is not p -compact at 0.

Furthermore, every polynomial $P_m f(y)$ in the Taylor series expansion of f at any $y \in \ell_1$ is 1-compact, and therefore p -compact for all $1 \leq p < \infty$, but f is not p -compact at any y .

Proof. Consider $\{\sigma_m\}_m$ the partition of the natural numbers such that each σ_m is a finite set of $m!$ consecutive elements:

$$\sigma_1 = \{1\}; \quad \sigma_2 = \underbrace{\{2, 3\}}_{2!}; \quad \sigma_3 = \underbrace{\{4, 5, 6, 7, 8, 9\}}_{3!}; \quad \sigma_4 = \underbrace{\{\dots\}}_{4!}; \quad \dots$$

Let $(e_j)_j$ be the canonical basis of ℓ_p and denote by $(e'_j)_j$ the sequence of coordinate functionals on ℓ_1 . Fixing $m \geq 1$, consider the polynomial $P_m \in \mathcal{P}({}^m\ell_1; \ell_p)$, defined by

$$P_m(x) = \left(\frac{m^{m/2}}{m!}\right)^{1/p} \sum_{j \in \sigma_m} e'_j(x)^m e_j.$$

Then

$$\|P_m\| = \left(\frac{m^{m/2}}{m!}\right)^{1/p} \sup_{x \in B_{\ell_1}} \left\| \sum_{j \in \sigma_m} e'_j(x)^m e_j \right\|_{\ell_p} \leq \left(\frac{m^{m/2}}{m!}\right)^{1/p} \sup_{x \in B_{\ell_1}} \|x\|_1^{1/p} = \left(\frac{m^{m/2}}{m!}\right)^{1/p}.$$

First, note that P_m is p -compact since it is of finite rank. Now, as $\limsup \|P_m\|^{1/m} \leq \lim \left(\frac{m^{1/2}}{m^{1/m}}\right)^{1/p} = 0$, we may define f as the series $\sum_{m=1}^\infty P_m$, and $f \in \mathcal{H}(\ell_1; \ell_p)$.

In order to show that f fails to be p -compact at 0, we show that $\limsup \kappa_p(P_m)^{1/m} = \infty$. Fix $m \in \mathbb{N}$ and take $(x_n)_n \in \ell_p(\ell_p)$, such that $P_m(B_{\ell_1}) \subset p\text{-co}\{x_n\}$. Each x_n may be written by $x_n = \sum_{k=1}^\infty x_k^n e_k$. For each $j \in \sigma_m$, there is a sequence $(\alpha_n^j)_n \in B_{\ell_q}$ such that

$$P_m(e_j) = \left(\frac{m^{m/2}}{m!}\right)^{1/p} e_j = \sum_{n=1}^\infty \alpha_n^j x_n = \sum_{n=1}^\infty \sum_{k=1}^\infty \alpha_n^j x_k^n e_k = \sum_{k=1}^\infty \left(\sum_{n=1}^\infty \alpha_n^j x_k^n\right) e_k.$$

Therefore, we have that $\left(\frac{m^{m/2}}{m!}\right)^{1/p} = \sum_{n=1}^\infty \alpha_n^j x_n^j$, for each $j \in \sigma_m$. Then

$$\begin{aligned} m^{m/2} &= \sum_{j \in \sigma_m} \left| \left(\frac{m^{m/2}}{m!}\right)^{1/p} \right|^p = \sum_{j \in \sigma_m} \left| \sum_{n=1}^\infty \alpha_n^j x_n^j \right|^p \\ &\leq \sum_{j \in \sigma_m} \left(\sum_{n=1}^\infty |\alpha_n^j x_n^j| \right)^p \\ &\leq \sum_{j \in \sigma_m} \left(\sum_{n=1}^\infty |\alpha_n^j|^q \right)^{p/q} \sum_{n=1}^\infty |x_n^j|^p \\ &\leq \sum_{j \in \sigma_m} \sum_{n=1}^\infty |x_n^j|^p \leq \|(x_n)_n\|_{\ell_p(\ell_p)}^p. \end{aligned}$$

We have shown that for any sequence $(x_n)_n \in \ell_p(\ell_p)$ such that $P_m(B_{\ell_1}) \subset p\text{-co}\{x_n\}$, the inequality $\|(x_n)_n\|_{\ell_p(\ell_p)} \geq m^{m/2p}$ holds. Then, $\kappa_p(P_m) \geq m^{m/2p}$ for all $m \in \mathbb{N}$ and $\limsup \kappa_p(P_m)^{1/m} = \infty$. By Proposition 3.4, f cannot be p -compact at 0, which proves the first statement of the example.

To show the second assertion, take any nonzero element $y \in \ell_1$ and fix $m_0 \in \mathbb{N}$. For all $x \in B_{\ell_1}$,

$$\begin{aligned} P_{m_0} f(y)(x) &= \sum_{m=m_0}^\infty \binom{m}{m_0} \check{P}_m(y^{m-m_0}, x^{m_0}) \\ &= \sum_{m=m_0}^\infty \binom{m}{m_0} \left(\frac{m^{m/2}}{m!}\right)^{1/p} \sum_{j \in \sigma_m} e'_j(y)^{m-m_0} e'_j(x)^{m_0} e_j. \end{aligned}$$

We claim that the sequence $\left(\binom{m}{m_0} \left(\frac{m^{m/2}}{m!}\right)^{1/p} e'_j(y)^{m-m_0} e_j\right)_{\substack{j \in \sigma_m \\ m > m_0}}$ belongs to $\ell_1(\ell_p)$. In fact,

$$\sum_{m > m_0} \binom{m}{m_0} \left(\frac{m^{m/2}}{m!}\right)^{1/p} \sum_{j \in \sigma_m} |e'_j(y)|^{m-m_0} \leq \sum_{m > m_0} \binom{m}{m_0} \left(\frac{m^{m/2}}{m!}\right)^{1/p} \|y\|_1^{m-m_0} < \infty.$$

Then, since $(e'_j(x)^{m_0})_{\substack{j \in \sigma_m \\ m \geq m_0}}$ belongs to B_{C_0} , the set $P_{m_0}f(y)(B_{\ell_1})$ is included in the 1-convex hull of

$$\left\{ \binom{m}{m_0} \left(\frac{m^{m/2}}{m!} \right)^{1/p} e'_j(y)^{m-m_0} e_j : m \geq m_0, j \in \sigma_m \right\},$$

which proves that $P_{m_0}f(y)$ is 1-compact and, therefore, p -compact for every $1 \leq p$, for any m_0 .

To show that f is not p -compact at y , note that fixed m , it is enough to choose $j \in \sigma_m$, to obtain that $P_m f(y)(e_j) = \left(\frac{m^{m/2}}{m!}\right)^{1/p} e_j$. Now, we can proceed as in the first part of the example to show that $\limsup \kappa_p(P_m f(y))^{1/m} = \infty$. And, again by Proposition 3.4, f cannot be p -compact at y . \square

The following example gives a negative answer to [2, Problem 5.1]. We show an entire function which is p -compact at 0, but this property does not extend beyond the ball $B_{r_p(f,0)}(0)$. Example 3.8 proves, in addition, that Proposition 3.6 cannot be improved. We base our construction in [15, Example 11].

Example 3.8. For every $1 \leq p < \infty$, there exists a holomorphic function $f \in \mathcal{H}(\ell_1; \ell_p)$ such that f is p -compact at 0, with $\limsup \kappa_p(P_m f(0))^{1/m} = 1$, but f is not p -compact at e_1 .

Proof. Consider $\{\sigma_m\}_m$, the partition of the natural numbers given in Example 3.7. Let $(e_j)_j$ be the canonical basis of ℓ_p and denote by $(e'_j)_j$ the sequence of coordinate functionals on ℓ_1 .

Fixing $m \geq 2$, define $P_m \in \mathcal{P}(^m \ell_1; \ell_p)$, the m -homogeneous polynomial

$$P_m(x) = \left(\frac{1}{m!}\right)^{1/p} e'_1(x)^{m-2} \sum_{j \in \sigma_m} e'_j(x)^2 e_j.$$

Then

$$\begin{aligned} \|P_m\| &= \left(\frac{1}{m!}\right)^{1/p} \sup_{x \in B_{\ell_1}} \left(\sum_{j \in \sigma_m} |e'_1(x)^{m-2} e'_j(x)^2|^p \right)^{1/p} \\ &\leq \left(\frac{1}{m!}\right)^{1/p} \sup_{x \in B_{\ell_1}} \left(\sum_{j \in \sigma_m} |e'_j(x)|^{2p} \right)^{1/p} \leq \left(\frac{1}{m!}\right)^{1/p}. \end{aligned}$$

Since $\lim \|P_m\|^{1/m} \leq \lim \left(\frac{1}{m!}\right)^{1/pm} = 0$, we may define f as $f(x) = \sum_{m \geq 2} P_m(x)$, which belongs to $\mathcal{H}(\ell_1; \ell_p)$ and $\sum_{m \geq 2} P_m$ is its Taylor series expansion at 0.

Note that each P_m is p -compact, as it is of finite rank, for all $m \geq 2$. Moreover, when computing $\|P_m\|$, we showed that $\alpha(x) = (e'_1(x)^{m-2} e'_j(x)^2)_j \in B_{\ell_q}$ for all $x \in B_{\ell_1}$. Then $P_m(B_{\ell_1}) \subset \left(\frac{1}{m!}\right)^{1/p} p\text{-co}\{e_j : j \in \sigma_m\}$ and since $\|(e_j)_{j \in \sigma_m}\|_{\ell_p(\ell_p)} = \left(\sum_{j \in \sigma_m} 1\right)^{1/p} = (m!)^{1/p}$, we have that $\kappa_p(P_m) \leq \left(\frac{1}{m!}\right)^{1/p} (m!)^{1/p} = 1$. Then, $\limsup \kappa_p(P_m)^{1/m} \leq 1$ and, by Proposition 3.4, f is p -compact at 0.

To show that $r_p(f, 0) = 1$, fix $m \geq 2$ and $\varepsilon > 0$. Take $x_j \in B_{\ell_1}$ such that $e'_1(x_j) = 1 - \varepsilon$, $e'_j(x_j) = \varepsilon$ and $e'_k(x_j) = 0$ for $j \in \sigma_m$ and $k \neq j$.

Now, fix any sequence $(y_n)_n \in \ell_p(\ell_p)$ such that $P_m(B_{\ell_1}) \subset p\text{-co}\{y_n\}$ and write $y_n = \sum_{k=1}^{\infty} \alpha_n^k e_k$.

Then, for each $j \in \sigma_m$ there exists $(\alpha_n^j)_n \in B_{\ell_q}$ so that

$$P_m(x_j) = \left(\frac{1}{m!}\right)^{1/p} (1 - \varepsilon)^{m-2} \varepsilon^2 e_j = \sum_{n=1}^{\infty} \alpha_n^j y_n.$$

Thus, for each $j \in \sigma_m$, the equality $\left(\frac{1}{m!}\right)^{1/p} (1 - \varepsilon)^{m-2} \varepsilon^2 = \sum_{n=1}^{\infty} \alpha_n^j y_n^j$ holds.

In consequence

$$\begin{aligned} ((1 - \varepsilon)^{m-2} \varepsilon^2)^p &= \sum_{j \in \sigma_m} \frac{1}{m!} ((1 - \varepsilon)^{m-2} \varepsilon^2)^p \\ &= \sum_{j \in \sigma_m} \left| \left(\frac{1}{m!}\right)^{1/p} (1 - \varepsilon)^{m-2} \varepsilon^2 \right|^p \\ &= \sum_{j \in \sigma_m} \left| \sum_{n=1}^{\infty} \alpha_n^j y_n^j \right|^p \end{aligned}$$

$$\begin{aligned} &\leq \sum_{j \in \sigma_m} \left(\sum_{n=1}^{\infty} |\alpha_n^j y_j^n| \right)^p \\ &\leq \sum_{j \in \sigma_m} \sum_{n=1}^{\infty} |y_j^n|^p \leq \|(y_n)_n\|_{\ell_p(\ell_p)}^p. \end{aligned}$$

Finally, we get that $\kappa_p(P_m) \geq (1 - \varepsilon)^{m-2} \varepsilon^2$ which implies that $\limsup \kappa_p(P_m)^{1/m} \geq 1 - \varepsilon$. Since $\varepsilon > 0$ was arbitrary, we obtain that $r_p(f, 0) = 1$.

Now, to prove that f is not p -compact at e_1 we show that the 2-homogeneous polynomial $P_2 f(e_1) : \ell_1 \rightarrow \ell_p$ is not p -compact. We have

$$P_2 f(e_1)(x) = \sum_{m=2}^{\infty} \binom{m}{2} \check{P}_m(e_1^{m-2}, x^2) \tag{2}$$

where \check{P}_m is the symmetric m -linear mapping associated to P_m .

By the definition of P_m we easily obtain a multilinear mapping $A_m \in \mathcal{L}^m(\ell_1; \ell_p)$ satisfying $P_m(x) = A_m(x, \dots, x)$, defined by

$$A_m(x_1, \dots, x_m) = \left(\frac{1}{m!} \right)^{1/p} e'_1(x_1) \cdots e'_1(x_{m-2}) \sum_{j \in \sigma_m} e'_j(x_{m-1}) e'_j(x_m) e_j.$$

Let S_m be the symmetric group on $\{1, \dots, m\}$ and denote by A_m^ξ , the multilinear mapping given by $A_m^\xi(x_1, \dots, x_m) = A_m(x_{\xi(1)}, \dots, x_{\xi(m)})$, where $\xi \in S_m$. Then we have

$$\check{P}_m(e_1^{m-2}, x^2) = \frac{1}{m!} \sum_{\xi \in S_m} A_m^\xi(e_1^{m-2}, x^2).$$

Since $A_m(x_1, \dots, x_{m-2}, e_1, x_{m-1}) = A_m(x_1, \dots, x_{m-1}, e_1) = 0$, for all $x_1, \dots, x_{m-1} \in \ell_1$, and $A_m(e_1^{m-2}, x^2) = (\frac{1}{m!})^{1/p} \times \sum_{j \in \sigma_m} e'_j(x)^2 e_j$, we obtain

$$\check{P}_m(e_1^{m-2}, x^2) = \frac{1}{m!} 2(m-2)! \left(\frac{1}{m!} \right)^{1/p} \sum_{j \in \sigma_m} e'_j(x)^2 e_j. \tag{3}$$

Combining (2) and (3) we get that

$$P_2 f(e_1)(x) = \sum_{m \geq 2} \left(\frac{1}{m!} \right)^{1/p} \sum_{j \in \sigma_m} e'_j(x)^2 e_j.$$

Suppose that $P_2 f(e_1)$ is p -compact. Hence, there exists a sequence $(y_n)_n \in \ell_p(\ell_p)$, $y_n = \sum_{k=1}^{\infty} y_k^n e_k$ such that $P_2 f(e_1)(B_{\ell_1}) \subset p\text{-co}\{y_n\}$. For each $j \in \sigma_m$, there exists $(\alpha_n^j)_n \in B_{\ell_q}$ such that $P_2 f(e_1)(e_j) = (\frac{1}{m!})^{1/p} e_j = \sum_{n=1}^{\infty} \alpha_n^j y_n$. As in Example 3.7, we conclude that $(\frac{1}{m!})^{1/p} = \sum_{n=1}^{\infty} \alpha_n^j y_n^j$, if $j \in \sigma_m$.

Hence

$$\begin{aligned} \sum_{m \geq 2} \sum_{j \in \sigma_m} \left(\left(\frac{1}{m!} \right)^{1/p} \right)^p &= \sum_{m \geq 2} \sum_{j \in \sigma_m} \left| \sum_{n=1}^{\infty} \alpha_n^j y_n^j \right|^p \\ &\leq \sum_{m \geq 2} \sum_{j \in \sigma_m} \left(\sum_{n=1}^{\infty} |\alpha_n^j|^q \right)^{p/q} \sum_{n=1}^{\infty} |y_n^j|^p \\ &\leq \sum_{m \geq 2} \sum_{j \in \sigma_m} \sum_{n=1}^{\infty} |y_n^j|^p \\ &\leq \|(y_n)_n\|_{\ell_p(\ell_p)}^p < \infty, \end{aligned}$$

which is a contradiction since $\sum_{m \geq 2} \sum_{j \in \sigma_m} ((\frac{1}{m!})^{1/p})^p$ is not convergent. Therefore, f cannot be p -compact at e_1 , and the result is proved. \square

4. The p -approximation property and p -compact mappings

The concept of p -compact sets leads naturally to that of p -approximation property. A Banach space E has the p -approximation property if the identity can be uniformly approximated by finite rank operators on p -compact sets. Since p -compact sets are compact, every space with the approximation property has the p -approximation property. Then, this property can be seen as a way to weaken the classical approximation property.

The p -approximation property has been studied in [10,12] related with p -compact linear operators and in [2] related with nonlinear mappings. The relation between the approximation property and compact holomorphic mappings was first addressed in [5]. Here, we are concerned with the study of the p -approximation property and its relation with p -compact polynomials and holomorphic functions in the spirit of [2] and [5].

We start by characterizing the notion of a homogeneous polynomial P being p -compact in terms of different conditions of continuity satisfied by P' the transpose of P . The first proposition gives an answer to [2, Problem 5.8] and should be compared with [5, Proposition 3.2].

Before going on, some words are needed on the topologies which we use. We denote by $\mathcal{P}_c^{(m)}(E)$ the space $\mathcal{P}^{(m)}(E)$ considered with the uniform convergence on compact sets of E , if $m = 1$ we simply write E'_c . When compact sets are replaced by p -compact sets we use the notation $\mathcal{P}_{cp}^{(m)}(E)$ and E'_{cp} . By the Ascoli theorem, any set $L \subset \mathcal{P}_c^{(m)}(E)$ is relatively compact if and only if $\sup_{P \in L} \|P\|$ is finite. Also, if $L \subset \mathcal{P}_{cp}^{(m)}(E)$ is relatively compact we have that L is point-wise bounded and then, by the principle of uniform boundedness, L is relatively compact in $\mathcal{P}_c^{(m)}(E)$. Now we have:

Proposition 4.1. *Let E and F be Banach spaces, $1 \leq p < \infty$, and $P \in \mathcal{P}^{(m)}(E; F)$. The following statements are equivalent.*

- (i) $P \in \mathcal{P}_{K_p}^{(m)}(E; F)$.
- (ii) $P' : F'_{cp} \rightarrow \mathcal{P}^{(m)}(E)$ is continuous.
- (iii) $P' : F'_{cp} \rightarrow \mathcal{P}_c^{(m)}(E)$ is compact.
- (iv) $P' : F'_{cp} \rightarrow \mathcal{P}_{cq}^{(m)}(E)$ is compact for any q , $1 \leq q < \infty$.
- (v) $P' : F'_{cp} \rightarrow \mathcal{P}_{cq}^{(m)}(E)$ is compact for some q , $1 \leq q < \infty$.

Proof. Suppose (i) holds, then $K = \overline{P(B_E)}$ is p -compact and its polar set K° is a neighborhood in F'_{cp} . For $y' \in K^\circ$ we have that $\|P'(y')\| = \sup_{x \in B_E} |y'(Px)| \leq 1$, and $P' : F'_{cp} \rightarrow \mathcal{P}^{(m)}(E)$ is continuous.

Now suppose (ii) holds, then there exists a p -compact set $K \subset F$ such that $P'(K^\circ)$ is equicontinuous in $\mathcal{P}^{(m)}(E)$. By the Ascoli theorem, $P'(K^\circ)$ is relatively compact in $\mathcal{P}_c^{(m)}(E)$ and $P' : F'_{cp} \rightarrow \mathcal{P}_c^{(m)}(E)$ is compact.

The continuity of the identity map $\mathcal{P}_c^{(m)}(E) \leftrightarrow \mathcal{P}_{cq}^{(m)}(E)$ gives that (iii) implies (iv), for all $1 \leq q < \infty$. Obviously, (iv) implies (v). To complete the proof, suppose (v) holds. Then, there exist an absolutely convex p -compact set $K \subset F$ and a compact set $L \subset \mathcal{P}_{cq}^{(m)}(E)$ such that $P'(K^\circ) \subset L$ and therefore, there exists $c > 0$ such that $\sup_{y' \in K^\circ} \|P'(y')\| \leq c$. Note that for any $x \in c^{-\frac{1}{m}} B_E$ and $y' \in K^\circ$ we have that $|P'(y')(x)| = |y'(Px)| \leq 1$. Then $P(x) \in K$, for all $x \in c^{-\frac{1}{m}} B_E$ and P is p -compact. \square

Now, we characterize the p -approximation property on a Banach space in terms of the p -compact homogeneous polynomials with values on it. In order to do so we appeal to the notion of the ϵ -product introduced by Schwartz [29]. Recall that for E and F two locally convex spaces, $F \in E$ is defined as the space of all linear continuous operators from E'_c to F , endowed with the topology of uniform convergence on all equicontinuous sets of E' . The space $F \in E$ is also denoted by $\mathcal{L}_\epsilon(E'_c; F)$. In [5, Proposition 3.3] is shown, for all Banach spaces E and F , that $(\mathcal{P}^{(m)}(F), \|\cdot\|) \in E = \mathcal{L}_\epsilon(E'_c; (\mathcal{P}^{(m)}(F), \|\cdot\|)) = (\mathcal{P}_K^{(m)}(F; E), \|\cdot\|)$, where the isomorphism is given by the transposition $P \leftrightarrow P'$. As a consequence, it is proved that $\mathcal{P}^{(m)}(F)$ has the approximation property if and only if $\mathcal{P}^{(m)}(F) \otimes E$ is $\|\cdot\|$ -dense in $\mathcal{P}_K^{(m)}(F; E)$ for all Banach spaces E and all $m \in \mathbb{N}$. We have the following result.

Proposition 4.2. *Let E and F be Banach spaces. Then $(\mathcal{P}_{K_p}^{(m)}(F; E), \|\cdot\|)$ is isometrically isomorphic to $\mathcal{L}_\epsilon(E'_{cp}; (\mathcal{P}^{(m)}(F), \|\cdot\|))$.*

As a consequence, E has the p -approximation property if and only if $\mathcal{P}^{(m)}(F) \otimes E$ is $\|\cdot\|$ -dense in $\mathcal{P}_{K_p}^{(m)}(F; E)$ for all Banach spaces F and all $m \in \mathbb{N}$.

Proof. Note that [(i) implies (ii)] of Proposition 4.1, says that the transposition operator maps a p -compact polynomial into a linear operator in $\mathcal{L}_\epsilon(E'_{cp}; (\mathcal{P}^{(m)}(F), \|\cdot\|))$. Now, take T in $\mathcal{L}_\epsilon(E'_{cp}; (\mathcal{P}^{(m)}(F), \|\cdot\|))$. Since the identity map $\iota : E'_c \rightarrow E'_{cp}$ is continuous, T belongs to $\mathcal{L}_\epsilon(E'_c; (\mathcal{P}^{(m)}(F), \|\cdot\|))$. By [5, Proposition 3.3], we have that $T = P'$ for some $P \in \mathcal{P}_K^{(m)}(F; E)$. In particular, $P' : E'_{cp} \rightarrow \mathcal{P}^{(m)}(F)$ is continuous and by [(ii) implies (i)] of Proposition 4.1, P is p -compact.

For the second statement, if E has the p -approximation property, $G \otimes E$ is dense in $\mathcal{L}_\epsilon(E'_{cp}; G)$, for every locally convex space G [22]. In particular we may consider $G = (\mathcal{P}^{(m)}(F), \|\cdot\|)$. Conversely, with $m = 1$ we have that $F' \otimes E$ is $\|\cdot\|$ -dense in $\mathcal{K}_p(F; E)$ for every Banach space F . Now, an application of [12, Theorem 2.1] completes the proof. \square

At the light of [5, Proposition 3.3], it would be expected to obtain a result of the type $\mathcal{P}^{(m)}E$ has the p -approximation property if and only if $\mathcal{P}^{(m)}E \otimes F$ is $\|\cdot\|$ -dense in $\mathcal{P}_{K_p}^{(m)}E; F$ for all Banach spaces F and all $m \in \mathbb{N}$. Unfortunately, our characterization is not as direct as we wanted and requires the following notion.

Definition 4.3. Let E be a Banach space, \mathcal{A} an operator ideal and α a norm on \mathcal{A} . We say that E has the (\mathcal{A}, α) -approximation property if $F' \otimes E$ is α -dense in $\mathcal{A}(F, E)$, for all Banach spaces F .

The relation between an ideal \mathcal{A} with the ideal of those operators whose transpose belongs to \mathcal{A} leads us to work with the ideal of quasi- p -nuclear operators \mathcal{QN}_p .

Proposition 4.4. Let E be a Banach space and fix $m \in \mathbb{N}$. Then,

- (a) $\mathcal{P}^{(m)}E \otimes F$ is $\|\cdot\|$ -dense in $\mathcal{P}_{K_p}^{(m)}E; F$, for all Banach spaces F if and only if $\mathcal{P}^{(m)}E$ has the $(\mathcal{QN}_p, \|\cdot\|)$ -approximation property.
- (b) $\mathcal{P}^{(m)}E$ has the p -approximation property if and only if $\mathcal{P}^{(m)}E \otimes F$ is $\|\cdot\|$ -dense in $\{P \in \mathcal{P}(E; F): L_p \in \mathcal{QN}_p(\otimes_{\pi_s}^m E; F)\}$, for all Banach spaces F .

Proof. The space $\mathcal{P}^{(m)}E$, or equivalently $(\otimes_{\pi_s}^m E)'$, has the $(\mathcal{QN}_p, \|\cdot\|)$ -approximation property if and only if $(\otimes_{\pi_s}^m E)' \otimes F$ is $\|\cdot\|$ -dense in $\mathcal{K}_p(\otimes_{\pi_s}^m E; F)$ for all Banach spaces F [22]. In virtue of Lemma 2.1, it is equivalent to have that $\mathcal{P}^{(m)}E \otimes F$ is $\|\cdot\|$ -dense in $\mathcal{P}_{K_p}^{(m)}E; F$. Then, statement (a) is proved. Note that (a) can be reformulated saying that $\mathcal{P}^{(m)}E$ has the $(\mathcal{QN}_p, \|\cdot\|)$ -approximation property if and only if $\mathcal{P}^{(m)}E \otimes F$ is $\|\cdot\|$ -dense in $\{P \in \mathcal{P}(E; F): L_p \in \mathcal{K}_p(\otimes_{\pi_s}^m E; F)\}$, for all Banach spaces F .

For the proof of (b), we use that the p -approximation property corresponds to the $(\mathcal{A}, \|\cdot\|)$ -approximation property for the ideal $\mathcal{A} = \mathcal{K}_p$, of p -compact operators. The result follows proceeding as before if the ideal \mathcal{K}_p and its dual ideal \mathcal{QN}_p are interchanged. \square

Now, we change our study to that of p -compact holomorphic mappings. Aron and Schottenloher described the space of compact holomorphic functions considered with τ_w , the Nachbin topology [26], via the ϵ -product. Namely, they show that $(\mathcal{H}_K(E; F), \tau_w) = \mathcal{L}_\epsilon(F'_{cp}; (\mathcal{H}(E), \tau_w))$, where the isomorphism is given by the transposition map $f \mapsto f'$ [5, Theorem 4.1]. The authors use this equivalence to obtain, in presence of the approximation property, results on density similar to that of Proposition 4.2. Recall that $f': F' \rightarrow \mathcal{H}(E)$ denotes the linear operator given by $f'(y') = y' \circ f$. With the next proposition we try to clarify the relationship between p -compact holomorphic mappings and the ϵ -product. The result obtained gives, somehow, a partial answer to [2, Problem 5.6].

Proposition 4.5. Let E and F be Banach spaces. Then,

- (a) $(\mathcal{H}_{K_p}(E; F), \tau_w)$ is topologically isomorphic to a subspace of $\mathcal{L}_\epsilon(F'_{cp}; (\mathcal{H}(E), \tau_w))$.
- (b) $\mathcal{L}_\epsilon(F'_{cp}; (\mathcal{H}(E), \tau_w))$ is topologically isomorphic to a subspace of $\{f \in \mathcal{H}(E; F): P_m f(x) \in \mathcal{P}_{K_p}^{(m)}E; F, \forall x \in E, \forall m \in \mathbb{N}\}$, considered with the Nachbin topology, τ_w .

Proof. To prove (a), fix f in $\mathcal{H}_{K_p}(E; F)$ and consider q any τ_w -continuous seminorm on $\mathcal{H}(E)$. By [16, Proposition 3.47], we may consider only the seminorms such that, for $g \in \mathcal{H}(E)$,

$$q(g) = \sum_{m=0}^{\infty} \|P_m g(0)\|_{K+a_m B_E},$$

with $K \subset E$ an absolutely convex compact set and $(a_m)_m$ a sequence in c_0^+ . There exists $V \subset E$, an open set such that $2K \subset V$ and $f(V) \subset F$ is p -compact. Fix $m_0 \in \mathbb{N}$ such that $2K + 2a_m B_E \subset V$, for all $m \geq m_0$. Now, choose $c > 0$ such that $c(2K + 2a_m B_E) \subset 2K + 2a_{m_0} B_E \subset V$, for all $m < m_0$. The polar set of $f(V)$, $f(V)^\circ$, is a neighborhood in F'_{cp} . By the Cauchy inequalities for entire functions, we have for all $y' \in f(V)^\circ$,

$$\begin{aligned} q(f'(y')) &= \sum_{m=0}^{\infty} \|P_m(y' \circ f)(0)\|_{K+a_m B_E} \\ &= \sum_{m=0}^{\infty} \frac{1}{2^m} \|P_m(y' \circ f)(0)\|_{2K+2a_m B_E} \\ &= \sum_{m < m_0} \frac{1}{2^m} \|P_m(y' \circ f)(0)\|_{2K+2a_m B_E} + \sum_{m \geq m_0} \frac{1}{2^m} \|P_m(y' \circ f)(0)\|_{2K+2a_m B_E} \end{aligned}$$

$$\begin{aligned} &\leq \sum_{m < m_0} \frac{1}{(2c)^m} \|P_m(y' \circ f)(0)\|_{c(2K+2a_m B_E)} + \sum_{m \geq m_0} \frac{1}{2^m} \|P_m(y' \circ f)(0)\|_{2K+2a_m B_E} \\ &\leq \sum_{m < m_0} \frac{1}{(2c)^m} \|y' \circ f\|_{c(2K+2a_m B_E)} + \sum_{m \geq m_0} \frac{1}{2^m} \|y' \circ f\|_{2K+2a_m B_E} \\ &\leq \sum_{m < m_0} \frac{1}{(2c)^m} \|y' \circ f\|_V + \sum_{m \geq m_0} \frac{1}{2^m} \|y' \circ f\|_V \\ &\leq \sum_{m < m_0} \frac{1}{(2c)^m} + \sum_{m \geq m_0} \frac{1}{2^m} < \infty. \end{aligned}$$

Then $f' \in \mathcal{L}(F'_{cp}; (\mathcal{H}(E), \tau_\omega))$. Again, we use the continuity of the identity map $\iota : F'_c \rightarrow F'_{cp}$ now, [5, Theorem 4.1] implies the result.

To prove that (b) holds, take $T \in \mathcal{L}(F'_{cp}; (\mathcal{H}(E), \tau_\omega))$ which, in particular, is an operator in $\mathcal{L}(F'_c; (\mathcal{H}(E), \tau_\omega))$. By [5, Theorem 4.1], $T = f'$ for some $f \in \mathcal{H}_K(E; F)$. By virtue of Proposition 4.1, it is enough to show that $(P_m f(x))' : F'_{cp} \rightarrow (\mathcal{P}^m E, \|\cdot\|)$ is continuous, for each $m \in \mathbb{N}$. Consider $D_x^m : (\mathcal{H}(E), \tau_\omega) \rightarrow (\mathcal{P}^m E, \|\cdot\|)$ the continuous projection given by $D_x^m(g) = P_m g(x)$, for all $g \in \mathcal{H}(E)$. Note that $(P_m f(x))'$ and $D_x^m \circ f'$ coincide as linear operators. Hence, the result follows. \square

Example 3.7 shows that there exists an entire function $f : \ell_1 \rightarrow \ell_p$, so that every homogeneous polynomial in its Taylor series expansion at y is q -compact, for any $y \in \ell_1$, for all $1 \leq q < \infty$, but f fails to be q -compact at y , for every y and every $q \leq p$. However, we have the following result.

Lemma 4.6. *Let E and F be Banach spaces. Then,*

$$\mathcal{H}_{K_p}(E; F) \text{ is } \tau_\omega\text{-dense in } \{f \in \mathcal{H}(E; F) : P_m f(x) \in \mathcal{P}_{K_p}({}^m E; F), \forall x \in E, \forall m \in \mathbb{N}\}.$$

Proof. Fix $f \in \mathcal{H}(E; F)$ so that $P_m f(x) \in \mathcal{P}_{K_p}({}^m E; F)$ for all $x \in E$ and for all m . Let $\varepsilon > 0$ and let q be any τ_ω -continuous seminorm on $\mathcal{H}(E; F)$ of the form

$$q(g) = \sum_{m=0}^{\infty} \|P_m g(0)\|_{K+a_m B_E},$$

with $K \subset E$ absolutely convex and compact and $(a_m)_m \in c_0^+$. Consider $m_0 \in \mathbb{N}$ such that $\sum_{m \geq m_0} \|P_m f(0)\|_{K+a_m B_E} < \varepsilon$. Now, let $f_0 = \sum_{m < m_0} P_m f(0)$, which is p -compact. Note that $q(f - f_0) \leq \varepsilon$ and the lemma follows. \square

Proposition 4.7. *Let E be a Banach space. Then, the following statements are equivalent.*

- (i) E has the p -approximation property.
- (ii) $\mathcal{H}(F) \otimes E$ is τ_ω -dense in $\mathcal{H}_{K_p}(F; E)$ for all Banach spaces F .

Proof. If E has the p -approximation property, $E \otimes G$ is dense in $\mathcal{L}_\epsilon(E'_c; G)$ for all locally convex space G [22], in particular if we consider $G = (\mathcal{H}(F), \tau_\omega)$. Applying Proposition 4.5(a), we have the first assertion.

For the converse, put $\mathcal{H}_0 = \{f \in \mathcal{H}(F; E) : P_m f(x) \in \mathcal{P}_{K_p}({}^m F; E), \forall x \in E, \forall m \in \mathbb{N}\}$. By Lemma 4.6, $\mathcal{H}(F) \otimes E$ is τ_ω -dense in \mathcal{H}_0 . Now, take $T \in \mathcal{K}_p(F; E)$ and $\varepsilon > 0$. Since $T \in \mathcal{H}_0$ and $q(f) = \|P_1 f(0)\|$ is a τ_ω -continuous seminorm, there exists $g \in \mathcal{H}(F) \otimes E$ such that $q(T - g) \leq \varepsilon$. But $q(T - g) = \|T - P_1 g(0)\|$ and since $P_1 g(0) \in F' \otimes E$, we have shown that $F' \otimes E$ is $\|\cdot\|$ -dense in $\mathcal{K}_p(F; E)$. By [12, Theorem 2.1], E has the p -approximation property. \square

5. Holomorphy types and topologies

In this section we show that p -compact holomorphic functions fit into the framework of holomorphy types. Our notation and terminology follow that given in [15]. Since $\mathcal{P}_{K_p}({}^m E; F)$ is a subspace of $\mathcal{P}({}^m E; F)$ and $\mathcal{P}_{K_p}({}^0 E; F) = F$, the first two conditions in the definition of a holomorphy type are fulfilled. Therefore, we only need to corroborate that the sequence $(\mathcal{P}_{K_p}({}^m E; F), \kappa_p)_m$ satisfies the third condition. Indeed, this last condition is also fulfilled if we show

$$\kappa_p(P_j(P)(a)) \leq (2e)^m \kappa_p(P) \|a\|^{m-j}, \tag{4}$$

for every $P \in \mathcal{P}_{K_p}({}^m E; F)$, for all $j = 1, \dots, m$ and for all m , where $P_j(P)(a)$ denotes the j -component in the expansion of P at a .

To give a simple proof of the fact that the sequence $(\mathcal{P}_{\mathcal{K}_p}(^m E; F), \kappa_p)_m$ satisfies the inequalities given in (4) we use the following notation. Let $P \in \mathcal{P}(^m E; F)$ and fix $a \in E$, we denote by P_{a^j} the $(m - j)$ -homogeneous polynomial defined as

$$P_{a^j}(x) := \check{P}(a^j, x^{m-j}),$$

for all $x \in E$ and $j < m$. Note that, for any $l < j < m$, we have that $P_{a^j} = (P_{a^{j-l}})_{a^l}$ and that $P_j(P)(a) = \binom{m}{m-j} P_{a^{m-j}}$. We appeal to the description of P_a given in [8, Corollary 1.8, b)]:

$$P_a(x) = \check{P}(a, x^{m-1}) = \frac{1}{m^2} \frac{1}{(m-1)^{m-1}} \sum_{j=1}^{m-1} P((m-1)r^j x + a), \tag{5}$$

where $r \in \mathbb{C}$ is such that $r^m = 1$ and $r^j \neq 1$ for $j < m$.

Theorem 5.1. *For any Banach spaces E and F , the sequence $(\mathcal{P}_{\mathcal{K}_p}(^m E; F), \kappa_p)_m$ is a holomorphy type from E to F .*

Proof. If $P \in \mathcal{P}_{\mathcal{K}_p}(^m E; F)$ by [2, Proposition 3.5] or Proposition 3.4 we have that $P_j(P)(a) \in \mathcal{P}_{\mathcal{K}_p}(^j E; F)$ for all $a \in E$, for all $j \leq m$. To prove the holomorphy type structure, we show that $\kappa_p(P_j(P)(a)) \leq 2^m e^m \|a\|^{m-j} \kappa_p(P)$, for all $j \leq m$.

Fix $a \in E$. If we show that $\kappa_p(P_a) \leq e \|a\| \kappa_p(P)$ then the proof is complete using a generalized inductive reasoning. Indeed, suppose that for any p -compact homogeneous polynomial Q , of degree less than m , the inequality $\kappa_p(Q_a) \leq e \|a\| \kappa_p(Q)$ holds. Then, since $P_{a^l} = (P_{a^{l-1}})_a$ and $P_j(P)(a) = \binom{m}{m-j} P_{a^{m-j}}$, we obtain

$$\begin{aligned} \kappa_p(P_j(P)(a)) &= \binom{m}{m-j} \kappa_p(P_{a^{m-j}}) = \binom{m}{m-j} \kappa_p((P_{a^{m-j-1}})_a) \\ &\leq \binom{m}{m-j} e \|a\| \kappa_p(P_{a^{m-j-1}}) \\ &\leq \binom{m}{m-j} e^{m-j} \|a\|^{m-j} \kappa_p(P) \\ &\leq 2^m e^m \|a\|^{m-j} \kappa_p(P). \end{aligned}$$

Now, take $P \in \mathcal{P}_{\mathcal{K}_p}(^m E; F)$. Then

$$\kappa_p(P_a) = m_p(\check{P}(a, B_E^{m-1})) = \|a\| m_p\left(\check{P}\left(\frac{a}{\|a\|}, B_E^{m-1}\right)\right). \tag{6}$$

Using (5) and Lemma 3.1 we have

$$\|a\| m_p\left(\check{P}\left(\frac{a}{\|a\|}, B_E^{m-1}\right)\right) \leq \frac{\|a\|}{m^2(m-1)^{m-1}} \sum_{j=1}^{m-1} m_p\left(P\left((m-1)r^j B_E + \frac{a}{\|a\|}\right)\right).$$

Since $\sup\{\|x\|: x \in (m-1)r^j B_E + \frac{a}{\|a\|}\} = m$,

$$\begin{aligned} \|a\| m_p\left(\check{P}\left(\frac{a}{\|a\|}, B_E^{m-1}\right)\right) &\leq \frac{\|a\|}{m^2(m-1)^{m-1}} \sum_{j=1}^{m-1} m^m m_p\left(P\left(\frac{1}{m}\left((m-1)r^j B_E + \frac{a}{\|a\|}\right)\right)\right) \\ &\leq \|a\| \left(\frac{m}{m-1}\right)^{m-1} \kappa_p(P) \leq e \|a\| \kappa_p(P). \end{aligned} \tag{7}$$

Combining (6) and (7) we get that $\kappa_p(P_a) \leq e \|a\| \kappa_p(P)$, as desired. \square

A function $f \in \mathcal{H}(E; F)$ is said to be of holomorphic type κ_p at a , if there exist $c_1, c_2 > 0$ such that each component of the Taylor series expansion of f at a is a p -compact polynomial satisfying that $\kappa_p(P_m f(a)) \leq c_1 c_2^m$. Now, we have the following result.

Corollary 5.2. *Let f be a function in $\mathcal{H}(E; F)$, then $f \in \mathcal{H}_{\mathcal{K}_p}(E; F)$ if and only if f is of κ_p -holomorphy type.*

Proof. It follows from Theorem 5.1 and [2, Proposition 3.5] or Proposition 3.4. \square

Remark 5.3. Theorem 5.1 can be improved. Indeed, the same proof of Theorem 5.1 shows that the sequence $(\mathcal{P}_{\mathcal{K}_p}(^m E; F))_m$ is a coherent sequence associated to the operator ideal $\mathcal{K}_p(E; F)$ (see [8] for definitions).

Since $\mathcal{H}_{K_p}(E, F)$ is a holomorphy type, following [27] we have a natural topology defined on $\mathcal{H}_{K_p}(E, F)$ denoted by τ_{ω, m_p} . This topology may be generated by different families of continuous seminorms. The original set of seminorms used to define τ_{ω, m_p} corresponds to the family of seminorms given below in Theorem 5.5, item (c). Our aim is to characterize the κ_p -approximation property of a Banach space E in an analogous way to [5, Theorem 4.1]. In order to do so, we give different descriptions of τ_{ω, m_p} . First, we need the following result.

Proposition 5.4. *Let E and F be Banach spaces. Then, $f \in \mathcal{H}_{K_p}(E; F)$ if and only if, for all m , $P_m f(0) \in \mathcal{P}_{K_p}({}^m E; F)$ and for any absolutely convex compact set K , there exists $\varepsilon > 0$ such that $\sum_{m=0}^{\infty} m_p(P_m f(0)(K + \varepsilon B_E)) < \infty$.*

Proof. Take $f \in \mathcal{H}_{K_p}(E; F)$ and K an absolutely convex compact set. Then, $2K$ is also absolutely convex and compact. For each $x \in 2K$, there exists $\varepsilon_x > 0$ such that $f(x + \varepsilon_x B_E)$ is p -compact. Now, we choose $x_1, \dots, x_n \in 2K$ such that $2K \subset \bigcup_{j=1}^n (x_j + \varepsilon_{x_j} B_E)$ and with $V = \bigcup_{j=1}^n (x_j + \varepsilon_{x_j} B_E)$ we have that $f(V)$ is p -compact. Let $d = \text{dist}(2K, \mathcal{C}V) > 0$, where $\mathcal{C}V$ denotes the complement of V . Let us consider $W = 2K + dB_E$, then W is an absolutely convex open set and $2K \subset W \subset V$. Then, applying Proposition 3.3 we have

$$\sum_{n=0}^{\infty} m_p \left(P_n f(0) \left(K + \frac{d}{2} B_E \right) \right) = \sum_{n=0}^{\infty} \left(\frac{1}{2} \right)^n m_p(P_n f(0)(W)) \leq 2m_p(f(W)) < \infty,$$

which proves the first claim.

Conversely, let $f \in \mathcal{H}(E; F)$ satisfy the conditions in the proposition. We have to show that f is p -compact at x for any fixed $x \in E$. Consider the absolutely convex compact set K , given by $K = \{\lambda x : |\lambda| \leq 1\}$. Then, there exists $\varepsilon_1 > 0$ such that $\sum_{n=0}^{\infty} m_p(P_n f(0)(K + \varepsilon_1 B_E)) < \infty$. Since f is entire, by [26, Proposition 1, p. 26], there exists $\varepsilon_2 > 0$ such that $f(y) = \sum_{m=1}^{\infty} P_m f(0)(y)$ uniformly for $y \in B_{\varepsilon_2}(x)$. Let $\varepsilon = \min\{\varepsilon_1, \varepsilon_2\}$, then $f(B_{\varepsilon}(x)) \subset \{\sum_{m=0}^{\infty} x_m : x_m \in P_m f(0)(B_{\varepsilon}(x))\}$.

Also

$$\sum_{m=0}^{\infty} m_p(P_m f(0)(B_{\varepsilon}(x))) \leq \sum_{m=0}^{\infty} m_p(P_m f(0)(K + \varepsilon_1 B_E)) < \infty.$$

Now, by Lemma 3.1, f is p -compact at x , and the proof is complete. \square

The next characterization of the topology τ_{ω, m_p} associated to the holomorphy type $\mathcal{H}_{K_p}(E; F)$ follows that of [15] and [26].

Theorem 5.5. *Any of the following families of seminorms generate the topology τ_{ω, m_p} on $\mathcal{H}_{K_p}(E; F)$.*

- (a) *The seminorms q satisfying that there exists a compact set K such that for every open set $V \supset K$ there exists $C_V > 0$ so that*

$$q(f) \leq C_V m_p(f(V)) \quad \forall f \in \mathcal{H}_{K_p}(E; F).$$

In this case, we say that q is m_p -ported by compact sets.

- (b) *The seminorms q satisfying that there exists an absolutely convex compact set K such that for every absolutely convex open set $V \supset K$ there exists $C_V > 0$ so that*

$$q(f) \leq C_V m_p(f(V)) \quad \forall f \in \mathcal{H}_{K_p}(E; F).$$

In this case, we say that q is AC- m_p -ported by absolutely convex compact sets.

- (c) *The seminorms q satisfying that there exists an absolutely convex compact set K such that, for all $\varepsilon > 0$ there exists $C(\varepsilon) > 0$ so that*

$$q(f) \leq C(\varepsilon) \sum_{m=0}^{\infty} \varepsilon^m \sup_{x \in K} \kappa_p(P_m f(x)) \quad \forall f \in \mathcal{H}_{K_p}(E; F).$$

- (d) *The seminorms q satisfying that there exists an absolutely convex compact set K such that, for all $\varepsilon > 0$ there exists $C(\varepsilon) > 0$ so that*

$$q(f) \leq C(\varepsilon) \sum_{m=0}^{\infty} m_p(P_m f(0)(K + \varepsilon B_E)) \quad \forall f \in \mathcal{H}_{K_p}(E; F).$$

- (e) *The seminorms of the form*

$$q(f) = \sum_{m=0}^{\infty} m_p(P_m f(0)(K + a_m B_E)),$$

where K ranges over all the absolutely convex compact sets and $(a_m)_m \in c_0^+$.

Proof. First note that if f is p -compact and K is a compact set, there exists an open set $V \supset K$ such that $f(V)$ is p -compact. Then, seminorms in (a) and (b) are well defined on $\mathcal{H}_{K_p}(E; F)$. Also, in virtue of Proposition 5.4, seminorms in (d) and (e) are well defined. Standard arguments show that seminorms in (a) and (b) define the same topology.

Now we show that seminorms in (b) and (c) coincide. Let q be a seminorm and let K be an absolutely convex compact set satisfying the conditions in (c). Let $V \supset K$ be any absolutely convex open set and take $d = \text{dist}(K, CV) > 0$. By Proposition 3.3, since $K + dB_E \subset V$, we get

$$m_p(P_m f(x)(dB_E)) \leq m_p(f(x + dB_E)) \leq m_p(f(V)),$$

for all $x \in K$ and $f \in \mathcal{H}_{K_p}(E; F)$. Thus,

$$d^m \sup_{x \in K} \kappa_p(P_m f(x)) \leq m_p(f(V)),$$

for each m . Hence

$$q(f) \leq C \left(\frac{d}{2}\right) \sum_{m=0}^{\infty} \left(\frac{d}{2}\right)^m \sup_{x \in K} \kappa_p(P_m f(x)) \leq 2C \left(\frac{d}{2}\right) m_p(f(V)),$$

which shows that q is AC- m_p -ported by K .

Conversely, let q be a seminorm, let K be an absolutely convex compact set satisfying the conditions in (b). Fix $\varepsilon > 0$ and take x_1, \dots, x_n in K such that $K \subset V$ with $V = \bigcup_{j=1}^n B_\varepsilon(x_j)$. As we did before, we may find an absolutely convex open set W so that $K \subset W \subset V$. Let $f \in \mathcal{H}_{K_p}(E; F)$, without loss of generality we may assume that $\varepsilon < r_p(f, x)$ for all $x \in K$. By Remark 3.5, we obtain

$$m_p(f(B_\varepsilon(x_j))) \leq \sum_{m=0}^{\infty} \varepsilon^m \kappa_p(P_m f(x_j)) \leq \sum_{m=0}^{\infty} \varepsilon^m \sup_{x \in K} \kappa_p(P_m f(x)).$$

As q is AC- m_p -ported by K , $q(f) \leq C_W m_p(f(W)) \leq C_W m_p(f(V))$ and therefore

$$\begin{aligned} q(f) &\leq C_W \sum_{j=1}^n m_p(f(B_\varepsilon(x_j))) \\ &\leq C_W \sum_{j=1}^n \sum_{m=0}^{\infty} \varepsilon^m \sup_{x \in K} \kappa_p(P_m f(x)) \\ &= nC_W \sum_{m=0}^{\infty} \varepsilon^m \sup_{x \in K} \kappa_p(P_m f(x)). \end{aligned}$$

Thus q belongs to the family in (c). If $\varepsilon \geq r_p(f, x)$, then $\sum_{m \geq 0} \varepsilon^m \sup_{x \in K} \kappa_p(P_m f(x)) = \infty$ and the inequality follows.

By the proof of [15, Proposition 4], we have that seminorms in (d) and (e) generate the same topology. Finally, we show that seminorms in (d) and (b) are equivalent. The proof of Proposition 5.4 shows that seminorms in (d) are AC- m_p -ported by absolutely convex compact sets.

To conclude the proof, consider a seminorm q and an absolutely convex compact set K satisfying conditions in (b). We borrow some ideas of [16, Chapter 3]. For each m , let W_m be the absolutely convex open set defined by $W_m = K + (\frac{1}{2})^m B_E$. Since q is AC- m_p -ported by K , for each $m \in \mathbb{N}$, there exists a constant $C_m = C_{W_m}$ such that $q(f) \leq C_m m_p(f(W_m))$, every p -compact function f .

For $m = 1$, there exists $n_1 \in \mathbb{N}$, such that for all $n > n_1$, $C_1^{1/n} < 2$. Take $V_1 = 2W_1$. Now, if $n > n_1$ and $Q \in \mathcal{P}_{K_p}({}^n E; F)$,

$$q(Q) \leq C_1 m_p(Q(W_1)) = m_p(Q(C_1^{1/n} W_1)) \leq m_p(Q(V_1)).$$

For $m = 2$, there exists $n_2 > n_1$ such that $C_2^{1/n} \leq 2$, for all $n > n_2$. Now, take $V_2 = 2W_2$ and, as before, we have for any $Q \in \mathcal{P}_{K_p}({}^n E; F)$, with $n > n_2$,

$$q(Q) \leq C_2 m_p(Q(W_2)) = m_p(Q(C_2^{1/n} W_2)) \leq m_p(Q(V_2)).$$

Repeating this procedure we obtain a sequence of absolutely convex open sets V_j satisfying

$$\begin{aligned} q(f) &\leq \sum_{m \geq 0} q(P_m f(0)) = \sum_{m < n_1} q(P_m f(0)) + \sum_{j \geq 1} \sum_{n_j \leq m < n_{j+1}} q(P_m f(0)) \\ &\leq C_{V_1} \sum_{m < n_1} m_p(P_m f(0)(V_1)) + \sum_{j \geq 1} \sum_{n_j \leq m < n_{j+1}} m_p(P_m f(0)(V_j)) \\ &\leq C \left(\sum_{m < n_1} m_p(P_m f(0)(V_1)) + \sum_{j \geq 1} \sum_{n_j \leq m < n_{j+1}} m_p(P_m f(0)(V_j)) \right) \end{aligned}$$

where $C = \min\{1, C_{V_j}\}$ and the result follows since $V_j = 2K + (\frac{1}{2})^{j-1}B_E$ and the seminorm q is bounded above by a seminorm of the family of the form (e). Now, the proof is complete. \square

We finish this section by inspecting the κ_p -approximation property introduced in [13]. We show that p -compact homogeneous polynomials from F to E can be κ_p -approximated by polynomials in $\mathcal{P}^{(m)}(F) \otimes E$ whenever E has the κ_p -approximation property. We then obtain a similar result for p -compact holomorphic functions. What follows keeps the spirit of [5, Theorem 4.1]. Recall that a Banach space E has the κ_p -approximation property if for every Banach space F , $F' \otimes E$ is κ_p -dense in $\mathcal{K}_p(F; E)$.

Theorem 5.6. *Let E be a Banach space. The following statements are equivalent.*

- (i) E has the κ_p -approximation property.
- (ii) For all $m \in \mathbb{N}$, $\mathcal{P}^{(m)}(F) \otimes E$ is κ_p -dense in $\mathcal{P}_{\mathcal{K}_p}^{(m)}(F; E)$, for every Banach space F .
- (iii) $\mathcal{H}(F) \otimes E$ is τ_{ω, m_p} -dense in $\mathcal{H}_{\mathcal{K}_p}(F; E)$ for all Banach spaces F .

Proof. First, suppose that E has the κ_p -approximation property and fix $m \in \mathbb{N}$. Then, $(\otimes_{\pi_s}^m F)' \otimes E$ is κ_p -dense in $\mathcal{K}_p(\otimes_{\pi_s}^m F; E)$ which coincides with $(\mathcal{P}_{\mathcal{K}_p}^{(m)}(F; E), \kappa_p)$, by Lemma 2.1. Thus, (ii) is satisfied.

Now, assume (ii) holds. Take $f \in \mathcal{H}_{\mathcal{K}_p}(F, E)$, $\varepsilon > 0$. By Theorem 5.5, we may consider a τ_{ω, m_p} -continuous seminorm of the form $q(f) = \sum_{m=0}^{\infty} m_p(P_m f(0)(K + a_m B_F))$, where $K \subset F$ is an absolutely convex compact set and $(a_m)_m \in c_0^+$. Let $m_0 \in \mathbb{N}$ be such that $\sum_{m > m_0} m_p(P_m f(0)(K + a_m B_F)) \leq \frac{\varepsilon}{2}$ and let $C > 0$ be such that $\frac{1}{C}(K + a_m B_F) \subset B_F$, for all $m \leq m_0$. Given $\delta > 0$, to be chosen later, by hypothesis, we may find $Q_m \in \mathcal{P}^{(m)}(F) \otimes E$ such that $\kappa_p(P_m f(0) - Q_m) \leq \delta$, for all $m \leq m_0$. Define $g = \sum_{m=0}^{m_0} Q_m$, which belongs to $\mathcal{H}(F) \otimes E$, then

$$\begin{aligned} q(f - g) &= \sum_{m=0}^{m_0} m_p((P_m f(0) - Q_m)(K + a_m B_F)) + \sum_{m > m_0} m_p(P_m f(0)(K + a_m B_F)) \\ &\leq \sum_{m=0}^{m_0} C^m \kappa_p((P_m f(0) - Q_m)) + \frac{\varepsilon}{2}. \end{aligned}$$

Thus, $q(f - g) < \varepsilon$ for a suitable choice of δ , which proves (iii).

Finally, suppose we have (iii). Take $T \in \mathcal{K}_p(F; E)$, $\varepsilon > 0$ and the seminorm on $\mathcal{H}_{\mathcal{K}_p}(F; E)$ defined by $q(f) = \kappa_p(P_1 f(0))$. Since q is τ_{ω, m_p} -continuous, by assumption, there exist $f_1, \dots, f_n \in \mathcal{H}(F)$ and $x_1, \dots, x_n \in E$, such that $q(T - \sum_{j=1}^n f_j \otimes x_j) < \varepsilon$. In other words, $\kappa_p(T - \sum_{j=1}^n P_1 f_j(0) \otimes x_j) < \varepsilon$ which proves that $F' \otimes E$ is κ_p -dense in $\mathcal{K}_p(F, E)$. Whence, the proof is complete. \square

Acknowledgments

We wish to express our gratitude to Richard Aron for introducing us to the problem during his visit to Buenos Aires in 2007 and for many useful conversations thereafter. Also, we are grateful to Manuel Maestre for his help and generosity. Finally, we thank Nacho Zalduendo for the suggestions he made to consider holomorphy types and to Christopher Boyd for his careful reading and comments on the manuscript.

References

- [1] R. Aron, P. Berner, A Hahn–Banach extension theorem for analytic mappings, Bull. Soc. Math. France 106 (1978) 3–24.
- [2] R. Aron, M. Maestre, P. Rueda, p -Compact holomorphic mappings, RACSAM Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. 104 (2) (2010) 353–364.
- [3] R. Aron, P. Rueda, p -Compact homogeneous polynomials from an ideal point of view, Contemp. Math., Amer. Math. Soc. 547 (2011) 61–71.
- [4] R. Aron, P. Rueda, \mathcal{I} -bounded holomorphic functions, preprint.
- [5] R. Aron, M. Schottenloher, Compact holomorphic mappings and the approximation property, J. Funct. Anal. 21 (1976) 7–30.
- [6] C. Boyd, S. Dineen, P. Rueda, Weakly uniformly continuous holomorphic functions and the approximation property, Indag. Math. (N.S.) 12 (2) (2001) 147–156.
- [7] E. Çalişkan, The bounded approximation property for spaces of holomorphic mappings on infinite dimensional spaces, Math. Nachr. 279 (7) (2006) 705–715.
- [8] D. Carando, V. Dimant, S. Muro, Coherent sequences of polynomial ideals on Banach spaces, Math. Nachr. 282 (8) (2009) 1111–1133.
- [9] P.G. Casazza, Approximation properties, in: Handbook of the Geometry of Banach Spaces, vol. I, North-Holland, Amsterdam, 2001, pp. 271–316.
- [10] Y.S. Choi, J.M. Kim, The dual space of $(\mathcal{L}(X, Y); \tau_p)$ and the p -approximation property, J. Funct. Anal. 259 (2010) 2437–2454.
- [11] A.M. Davie, T.W. Gamelin, A theorem on polynomial-star approximation, Proc. Amer. Math. Soc. 106 (2) (1989) 351–356.
- [12] J.M. Delgado, E. Oja, C. Piñeiro, E. Serrano, The p -approximation property in terms of density of finite rank operators, J. Math. Anal. Appl. 354 (2009) 159–164.
- [13] J.M. Delgado, C. Piñeiro, E. Serrano, Density of finite rank operators in the Banach space of p -compact operators, J. Math. Anal. Appl. 370 (2010) 498–505.
- [14] J.M. Delgado, C. Piñeiro, E. Serrano, Operators whose adjoints are quasi p -nuclear, Studia Math. 197 (3) (2010) 291–304.
- [15] S. Dineen, Holomorphy types on a Banach space, Studia Math. 39 (1971) 241–288.
- [16] S. Dineen, Complex Analysis of Infinite Dimensional Spaces, Springer Monogr. Math., Springer, 1999.

- [17] S. Dineen, J. Mujica, The approximation property for spaces of holomorphic functions on infinite-dimensional spaces. I, *J. Approx. Theory* 126 (2) (2004) 141–156.
- [18] S. Dineen, J. Mujica, The approximation property for spaces of holomorphic functions on infinite-dimensional spaces. II, *J. Funct. Anal.* 259 (2) (2010) 545–560.
- [19] P. Enflo, A counterexample to the approximation problem in Banach spaces, *Acta Math.* 130 (1) (1973) 309–317.
- [20] D. Galicer, S. Lassalle, P. Turco, The ideal of p -compact operators: a tensor product approach, preprint.
- [21] A. Grothendieck, Produits tensoriels topologiques et espaces nucléaires, *Mem. Amer. Math. Soc.* (16) (1955), 140 pp.
- [22] S. Lassalle, P. Turco, On p -approximation properties and operator ideals, in preparation.
- [23] J. Lindenstrauss, L. Tzafriri, *Classical Banach Spaces. I*, vol. 92, Springer-Verlag, Berlin, New York, 1977.
- [24] J. Mujica, *Complex Analysis in Banach Spaces*, *Math. Stud.*, vol. 120, North-Holland, Amsterdam, 1986.
- [25] J. Mujica, Linearization of bounded holomorphic mappings on Banach spaces, *Trans. Amer. Math. Soc.* 324 (2) (1991) 867–887.
- [26] L. Nachbin, *Topology on Spaces of Holomorphic Mappings*, *Ergeb. Math. Grenzgeb.*, vol. 47, Springer-Verlag, Berlin, 1969.
- [27] L. Nachbin, Concerning holomorphy types for Banach spaces, *Studia Math.* 38 (1970) 407–412.
- [28] A. Persson, A. Pietsch, p -Nukleare und p -integrale Abbildungen in Banachräumen, *Studia Math.* 33 (1969) 19–62 (in German).
- [29] L. Schwartz, Théorie des distributions à valeurs vectorielles. I, *Ann. Inst. Fourier (Grenoble)* 7 (1957) 1–141 (in French).
- [30] D.P. Sinha, A.K. Karn, Compact operators whose adjoints factor through subspaces of ℓ_p , *Studia Math.* 150 (2002) 17–33.
- [31] D.P. Sinha, A.K. Karn, Compact operators which factor through subspaces of ℓ_p , *Math. Nachr.* 281 (2008) 412–423.