

## Gibbs random graphs on point processes

Pablo A. Ferrari, Eugene A. Pechersky, Valentin V. Sisko, and Anatoly A. Yambartsev

Citation: *Journal of Mathematical Physics* **51**, 113303 (2010); doi: 10.1063/1.3514605

View online: <http://dx.doi.org/10.1063/1.3514605>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jmp/51/11?ver=pdfcov>

Published by the [AIP Publishing](#)

---

### Articles you may be interested in

[Robustness of random graphs based on graph spectra](#)

*Chaos* **22**, 043101 (2012); 10.1063/1.4754875

[Analysis of random Landau-Lifshitz dynamics by using stochastic processes on graphs](#)

*J. Appl. Phys.* **99**, 08F301 (2006); 10.1063/1.2165582

[Random Feynman Graphs](#)

*AIP Conf. Proc.* **776**, 118 (2005); 10.1063/1.1985383

[Kinetic Theory of Random Graphs](#)

*AIP Conf. Proc.* **776**, 3 (2005); 10.1063/1.1985373

[An inequality for the Gibbs mean number of clusters](#)

*J. Math. Phys.* **46**, 023301 (2005); 10.1063/1.1845972

---

Did your publisher get  
**18 MILLION DOWNLOADS** in 2014?  
AIP Publishing did.



THERE'S POWER IN NUMBERS. Reach the world with AIP Publishing.



## Gibbs random graphs on point processes

Pablo A. Ferrari,<sup>1,4,a)</sup> Eugene A. Pechersky,<sup>2,b)</sup> Valentin V. Sisko,<sup>3,c)</sup> and Anatoly A. Yambartsev<sup>4,d)</sup>

<sup>1</sup>*Departamento de Matemática, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Pabellón 1, Ciudad Universitaria, 1428 Buenos Aires, Argentina*

<sup>2</sup>*Dobrushin Laboratory of Institute for Information Transmission Problems of Russian Academy of Sciences, 19, Bolshoj Karetny, Moscow, Russia*

<sup>3</sup>*Federal University of Fluminense, Institute of Mathematics and Statistics, Rua Mário Santos Braga, s/n, 24020-140 Niterói, RJ, Brazil*

<sup>4</sup>*Department of Statistics, Institute of Mathematics and Statistics, University of São Paulo, Rua do Matão 1010, CEP 05508-090, São Paulo SP, Brazil*

(Received 3 March 2010; accepted 12 October 2010; published online 29 November 2010)

Consider a discrete locally finite subset  $\Gamma$  of  $\mathbb{R}^d$  and the complete graph  $(\Gamma, E)$ , with vertices  $\Gamma$  and edges  $E$ . We consider Gibbs measures on the set of sub-graphs with vertices  $\Gamma$  and edges  $E' \subset E$ . The Gibbs interaction acts between open edges having a vertex in common. We study percolation properties of the Gibbs distribution of the graph ensemble. The main results concern percolation properties of the open edges in two cases: (a) when  $\Gamma$  is sampled from a homogeneous Poisson process; and (b) for a fixed  $\Gamma$  with sufficiently sparse points. © 2010 American Institute of Physics. [doi:10.1063/1.3514605]

### I. INTRODUCTION

Let a sample  $\Gamma \subset \mathbb{R}^d$  of a *point process* be a locally finite set of  $\mathbb{R}^d$ . We consider an ensemble of graphs with *vertices*  $\Gamma$  and random *edges* belonging to the set of unordered pairs of points in  $\Gamma$ . The edges can be open or closed and we study probability distributions on the set of configurations of open edges. The classical example is the *Erdős-Rényi's random graph* where each edge is open independently of the others with some probability, see Refs. 1 and 2. We introduce interactions between edges and/or vertices and study the associated Gibbs measures.

Given a configuration of open edges, two edges *collide* if both of them are open and they have a vertex in common. *Monomers* are those vertices that belong to no open edge. Each collision and each monomer contribute a positive energy; any open edge contributes a positive energy proportional to its length. The energy function  $H$  is described explicitly in (1) later. The Gibbs measure associated with  $H$  at inverse temperature  $\beta > 0$  gives more weight to configurations with few monomers, few (or no) collisions and short edges.

In Theorem 1, we give sufficient conditions on  $\Gamma$  for the existence of an infinite volume Gibbs measure. An open edge not colliding with any other edge is called *dimer*. In Theorem 2, we show that the ground states for any locally finite configuration  $\Gamma$  are composed only of monomers and dimers. Theorem 3 gives conditions for the uniqueness of the ground state. Theorem 4 shows that if  $\Gamma$  is a sample of a homogeneous Poisson process with small density and low temperature, then there is no percolation for almost all  $\Gamma$ . Non-percolation in this context is the absence of an infinite sequence of colliding open edges a.s. with respect to the Gibbs measure. Theorem 5 proves that if  $\Gamma$  is “sparse” and the temperature is sufficiently low, then there is no percolation a.s. with respect to

a) Electronic mail: pferrari@dm.uba.ar and pablo@ime.usp.br.

b) Electronic mail: pech@iitp.ru.

c) Electronic mail: valentin@mat.uff.br.

d) Electronic mail: yambar@ime.usp.br.

the Gibbs measure. In particular for  $\Gamma$  satisfying an  $\varepsilon$ -hard-core condition (that is, the ball of radius  $\varepsilon$  around each point of  $\Gamma$  has no other point of  $\Gamma$ ) for some  $\varepsilon > 0$ , there is no percolation.

## II. DEFINITIONS

Let  $\Gamma$  be a locally finite set of  $\mathbb{R}^d$  and consider the complete graph  $(\Gamma, E)$  with vertex set  $\Gamma$  and edge set  $E := \{\gamma\gamma' : \gamma, \gamma' \in \Gamma\}$ . The length of the edge  $e = \gamma\gamma'$ , is denoted by  $L(e) := |\gamma - \gamma'|$ . For a point  $\gamma \in \Gamma$ , let  $S_\gamma := \{\gamma\gamma' \in E : \gamma' \in \Gamma\}$ , the set of edges incident on  $\gamma$ .

Let  $\overline{\Omega}$  be the set of subsets of  $E$ . A configuration  $\omega \in \overline{\Omega}$  defines the sub-graph  $(\Gamma, \omega)$ . Edges  $e \in \omega$  are called *open* with respect to  $\omega$ . Edges  $e \in E \setminus \omega$  are called *closed* with respect to  $\omega$ . When it is clear from the context we use the terms open and closed with no mention to the configuration  $\omega$ . The *degree*  $d_\gamma(\omega)$  of a vertex  $\gamma \in \Gamma$  is the number of open edges in  $\omega$  containing  $\gamma$ :

$$d_\gamma(\omega) := |S_\gamma \cap \omega|.$$

### A. Gibbs measures

The set of  $\omega \in \overline{\Omega}$  with finite-degree vertices is called

$$\Omega := \{\omega : \omega \in \overline{\Omega}, d_\gamma(\omega) < \infty \text{ for all } \gamma\}.$$

Our goal is to define a Gibbs distribution on  $\Omega$  associated to the following formal Hamiltonian

$$H(\omega) := \sum_{e \in \omega} L(e) + \sum_{\gamma \in \Gamma} \phi_\gamma(\omega), \tag{1}$$

where  $\phi_\gamma(\omega)$  is the contribution of monomers and interacting edges in  $S_\gamma$  defined by

$$\phi_\gamma(\omega) := \begin{cases} h_0, & \text{if } d_\gamma(\omega) = 0; \\ 0, & \text{if } d_\gamma(\omega) = 1; \\ h_1 \binom{d_\gamma(\omega)}{2}, & \text{if } d_\gamma(\omega) \geq 2. \end{cases} \tag{2}$$

where  $0 < h_0 < h_1$  are fixed parameters. The potential  $\phi_\gamma(\omega)$  depends only on the degree  $d_\gamma(\omega)$ . Degree zero contributes  $h_0$ , degree 1 does not contribute and each pair of open edges incident on  $\gamma$  contributes  $h_1$ .

The potential function  $\phi_\gamma$  depends on infinitely many edges because to establish if  $d_\gamma(\omega) = 0$  we have to check that  $e \in \omega$  for infinitely many edges  $e \in S_\gamma$ . However, the usual Gibbs construction (including existence theorem) for this case works without special considerations.

We define a family of finite volume Gibbs measures. For bounded  $\Lambda \subset \mathbb{R}^d$  the set  $\Gamma_\Lambda := \Gamma \cap \Lambda$  is finite and so is  $E_\Lambda := \{\gamma\gamma' : \gamma, \gamma' \in \Gamma_\Lambda\}$ . The Gibbs state  $P_\Lambda$  on  $\Omega_\Lambda := \{\omega \in \overline{\Omega} : \omega \subset E_\Lambda\}$  with the zero boundary configuration is defined by

$$P_\Lambda(\omega) := \frac{\exp\{-\beta H_\Lambda(\omega)\}}{Z_\Lambda}, \quad \omega \in \Omega_\Lambda, \tag{3}$$

where the parameter  $\beta$  is called *inverse temperature*,  $Z_\Lambda$  is the normalizing constant and

$$H_\Lambda(\omega) := \sum_{e \in \omega} L(e) + \sum_{\gamma \in \Gamma_\Lambda} \phi_\gamma(\omega), \quad \omega \in \Omega_\Lambda. \tag{4}$$

Denote  $E_\Lambda$  the expectation with respect to  $P_\Lambda$ .

We only consider Gibbs distributions  $P$  on  $\overline{\Omega}$  associated with the formal Hamiltonian  $H$  defined in (1) that can be constructed as a limit along subsequences of  $P_\Lambda$ ,  $\Lambda \uparrow \mathbb{R}^d$ , where  $P_\Lambda$  is defined in (3). Since  $\overline{\Omega}$  is compact,  $P$  exists but may have infinite-degree vertices. In Theorem 1 we show that under mild conditions on  $\Gamma$ , all vertex has finite degree with  $P$ -probability 1. This implies that  $P$  is concentrated on  $\Omega$ . Uniqueness of  $P$  is not discussed in this article.

The measure  $\mathbb{P}_\Lambda$  can be seen as a measure on  $\overline{\Omega}$  concentrating mass in the set  $\Omega_\Lambda$ .

## B. Percolation

A path of length  $n$  connecting  $\gamma_0$  with  $\gamma_n$  in the graph  $(\Gamma, E)$  is a sequence of distinct vertices  $\gamma_0, \dots, \gamma_n$  and the edges connecting successive points. A path is *open* if all its edges  $\gamma_i \gamma_{i+1}$  are open. A connected component of  $\omega$  is a set of points that can be mutually connected by open paths and the open edges incident to those points. Maximal connected components of  $\omega$  are called *open clusters*. The *open cluster at  $\gamma$* , called  $C_\gamma(\omega)$ , is the maximal connected component containing  $\gamma$ . The *vertex set* of  $C_\gamma(\omega)$  is the set of all vertices in  $\Gamma$  which are connected to  $\gamma$  by open paths and the *edge set* of  $C_\gamma(\omega)$  is the set of edges of  $\omega$  which join pairs of such vertices.

We give sufficient conditions for the absence of infinite clusters with  $\mathbb{P}$  probability one; this is called *no percolation*.

## III. MAIN RESULTS

### A. Existence

Let  $\alpha > 0$ . A point set  $\Gamma$  is  $\alpha$ -homogeneous if for any  $\gamma \in \Gamma$ ,

$$T_\gamma(\alpha) := \sum_{\gamma' \in \Gamma} e^{-\alpha L(\gamma\gamma')} < \infty. \quad (5)$$

A point set  $\Gamma$  is *uniformly  $\alpha$ -homogeneous* if

$$T(\alpha) := \sup_{\gamma \in \Gamma} T_\gamma(\alpha) < \infty. \quad (6)$$

If  $\Gamma$  consists of hard core ball centers of a fixed radius then  $\Gamma$  is uniformly  $\alpha$ -homogeneous for all  $\alpha$ ; this is because the number of points grows polynomially with the distance while the weight decreases exponentially. Almost all sample  $\Gamma$  from a Poisson process is  $\alpha$ -homogeneous for all  $\alpha$  but uniformly  $\alpha$ -homogeneous for no  $\alpha$ ; see Lemma 2.

**Theorem 1:** Take  $0 < h_0 < h_1$ ,  $\beta > 0$  and a  $\beta$ -homogeneous  $\Gamma$ . Then, any Gibbs measure  $\mathbb{P}$  at inverse temperature  $\beta$  is concentrated on  $\Omega$ .

### B. Ground states

A configuration  $\widehat{\omega}$  is a *local perturbation* of  $\omega \in \Omega$  if the symmetric difference  $\widehat{\omega} \Delta \omega$  is a finite set. A configuration  $\omega \in \Omega$  is a *ground state* if

$$H(\widehat{\omega}) - H(\omega) \geq 0$$

for any local perturbation  $\widehat{\omega}$  of  $\omega$ . The difference is well defined because all but a finite number of terms vanish.

The next result says that all ground states are composed by dimers and monomers.

**Theorem 2:** For any locally finite configuration  $\Gamma$  and any  $0 < h_0 < h_1$  there exists at least one ground state. If  $\omega$  is a ground state then

$$d_\gamma(\omega) \leq 1$$

for every  $\gamma \in \Gamma$ . The length of any open edge in  $\omega$  is less than  $2h_0$ .

Let  $\pi_\lambda$  be the distribution of a homogeneous Poisson process on  $\mathbb{R}^d$  with rate  $\lambda > 0$ .

**Theorem 3:** There exists  $\lambda_g$  such that if  $\lambda < \lambda_g$ , then for  $\pi_\lambda$ -almost all  $\Gamma$  there is only one ground state.

**C. No percolation at low rate and temperature**

We consider the random graph as a dependent percolation model.<sup>4</sup> For  $\gamma \in \Gamma$ , let  $|C_\gamma(\omega)|$  be the number of vertices in  $C_\gamma(\omega)$  and let  $\theta_\gamma(\mathbf{P})$  be the probability that the open cluster at  $\gamma$  is infinite:

$$\theta_\gamma(\mathbf{P}) := \mathbf{P}(|C_\gamma(\omega)| = \infty).$$

We say that there is no percolation for  $\mathbf{P}$  if  $\theta_\gamma(\mathbf{P}) = 0$ .

Take a Poisson process of rate  $\lambda$  and a Gibbs state at inverse temperature  $\beta$ . We establish a  $(\lambda, \beta)$ -region where there is no percolation for  $\mathbf{P}$ :

**Theorem 4:** *If*

$$\lambda \left( 2h_0 + \frac{\log 2}{\beta} \right) \leq 1 \tag{7}$$

*then there is no percolation for  $\mathbf{P}$  for  $\pi_\lambda$ -almost all  $\Gamma$ .*

The next theorem is stronger but for more restricted sets  $\Gamma$ .

**Theorem 5:** *If  $\Gamma$  is such that*

$$\varepsilon := T(\beta) \exp(-\beta(h_1 - h_0)) < 1, \tag{8}$$

*then there is no percolation for  $\mathbf{P}$ .*

*Remark 1:* Fix  $\Gamma, h_0$  and  $h_1$ . Then  $T(\beta)$  is a non-increasing function of  $\beta$ . Therefore the inequality (8) is valid for all  $\beta$  big enough.

**IV. PROOFS**

Along this section  $\Gamma$  is assumed  $\beta$ -homogeneous. Let  $\mathbf{P}^0$  be the product measure on  $\bar{\Omega}$  with marginals

$$\mathbf{P}^0(e \in \omega) = \frac{e^{-\beta L(e)}}{e^{-\beta L(e)} + e^{-2\beta h_0}}. \tag{9}$$

*Lemma 1:* Any infinite volume Gibbs measure  $\mathbf{P}$  is stochastically dominated by the product measure  $\mathbf{P}^0$ : for any finite set of edges  $\zeta \subset E$ ,

$$\mathbf{P}(\omega \supset \zeta) \leq \mathbf{P}^0(\omega \supset \zeta).$$

*Proof:* We start proving that for any bounded  $\Lambda \subset \mathbb{R}^d$ ,  $\mathbf{P}_\Lambda$  is stochastically dominated by  $\mathbf{P}_\Lambda^0$  the product measure defined on  $\Omega_\Lambda$ . Holley Theorem says that if for all  $\tilde{\omega} \subset E_\Lambda$

$$\mathbf{p} := \mathbf{P}_\Lambda(e \in \omega \mid \omega \setminus \{e\} = \tilde{\omega} \setminus \{e\}) \leq \mathbf{P}_\Lambda^0(e \in \omega). \tag{10}$$

then  $\mathbf{P}_\Lambda$  is stochastically dominated by  $\mathbf{P}_\Lambda^0$ ; see Theorem 4.8 in Ref. 3. This is enough to prove the lemma because for finite  $\zeta$ ,  $\{\omega \supset \zeta\}$  is a cylinder set and the limit

$$\begin{aligned} \mathbf{P}(\omega \supset \zeta) &= \lim_{\Lambda \uparrow \mathbb{R}^d} \mathbf{P}_\Lambda(\omega \supset \zeta) \\ &\leq \lim_{\Lambda \uparrow \mathbb{R}^d} \mathbf{P}_\Lambda^0(\omega \supset \zeta) = \mathbf{P}^0(\omega \supset \zeta) \end{aligned}$$

holds along subsequences.

Now we prove (10). Let  $e = \gamma_1 \gamma_2 \in E_\Lambda$ . The conditional probability in (10) depends only on configurations on  $(S_{\gamma_1} \cup S_{\gamma_2}) \cap E_\Lambda \setminus \{e\}$ :

$$\mathbf{p} = \mathbf{P}_\Lambda(e \in \omega \mid \omega_{\gamma_1} = \tilde{\omega}_{\gamma_1}, \omega_{\gamma_2} = \tilde{\omega}_{\gamma_2}),$$

where  $\omega_\gamma = \omega \cap S_\gamma \cap E_\Lambda \setminus \{e\}$ . Consider three cases:

1.  $\tilde{\omega}_{\gamma_1} = \tilde{\omega}_{\gamma_2} = \emptyset$ ,
3.  $\tilde{\omega}_{\gamma_1} \neq \emptyset, \tilde{\omega}_{\gamma_2} = \emptyset$  (and its symmetric version  $\tilde{\omega}_{\gamma_1} = \emptyset, \tilde{\omega}_{\gamma_2} \neq \emptyset$ ).
3.  $\tilde{\omega}_{\gamma_1} \neq \emptyset, \tilde{\omega}_{\gamma_2} \neq \emptyset$ .

Case 1: The edge  $e$  is isolated of the rest, so the probability to be in  $\omega$  is

$$p = \frac{e^{-\beta L(e)}}{e^{-\beta L(e)} + e^{-2\beta h_0}} = P^0(e \in \omega).$$

Case 2: Let  $m$  be the number of open edges in  $\tilde{\omega}_{\gamma_1}$ . If  $e$  is open then it interacts with  $m$  open edges:

$$p = \frac{e^{-\beta L(e) - \beta m h_1}}{e^{-\beta L(e) - \beta m h_1} + e^{-\beta h_0}} < \frac{e^{-\beta L(e)}}{e^{-\beta L(e)} + e^{-2\beta h_0}} = P^0(e \in \omega).$$

Case 3: Let  $m$  be the number of open edges in  $\tilde{\omega}_{\gamma_1} \cup \tilde{\omega}_{\gamma_2}$ . Then,

$$p = \frac{e^{-\beta L(e) - \beta m h_1}}{e^{-\beta L(e) - \beta m h_1} + 1} < \frac{e^{-\beta L(e)}}{e^{-\beta L(e)} + 1} < \frac{e^{-\beta L(e)}}{e^{-\beta L(e)} + e^{-2\beta h_0}} = P^0(e \in \omega). \quad \square$$

*Proof of Theorem 1:* Use Lemma 1 to dominate the degree of  $\gamma$  as follows

$$\begin{aligned} Ed_\gamma &= \sum_{\gamma' \in \Gamma} P(\gamma\gamma' \in \omega) \leq \sum_{\gamma' \in \Gamma} P^0(\gamma\gamma' \in \omega) \\ &= \sum_{\gamma' \in \Gamma} \frac{e^{-\beta L(\gamma\gamma')}}{e^{-\beta L(\gamma\gamma')} + e^{-2\beta h_0}} \leq \frac{T_\gamma(\beta)}{e^{-2\beta h_0}} < \infty, \end{aligned} \tag{11}$$

because  $\Gamma$  is  $\beta$ -homogeneous by hypothesis. □

*Lemma 2:* Almost all samples  $\Gamma$  from the Poisson process distribution  $\pi_\lambda$  are  $\alpha$ -homogeneous for all  $\alpha$ .

*Proof:* Without losing generality assume  $0 \in \Gamma$ . Consider a sequence of hypercubes  $\Lambda_n = [-n^{1/d}, n^{1/d}]^d, n \geq 0$ . Any ring  $W_n = \Lambda_{n+1} \setminus \Lambda_n, n \geq 0$ , has volume 2. If  $\gamma \in W_n$  then  $L(0\gamma) \geq n^{1/d}$  and

$$\sum_{\gamma \in \Gamma} e^{-\alpha L(0\gamma)} \leq \sum_{n=0}^{\infty} |\Gamma \cap W_n| \exp(-\alpha n^{1/d}).$$

Since  $|\Gamma \cap W_n|$  is a Poisson random variable with mean 2 for all  $n$ , the sum is finite  $\pi_\lambda$ -a.s. for any  $\alpha > 0$ . □

*Proof of Theorem 2:* Assume  $\omega$  is a ground state and proceed by contradiction: assume that there exists a vertex  $\gamma \in \Gamma$  such that  $d_\gamma(\omega) \geq 2$ . Let  $\gamma\gamma' \in \omega$ . Let  $\tilde{\omega}$  be the same as  $\omega$  but without the edge  $\gamma\gamma'$ :

$$\tilde{\omega} := \omega \setminus \{\gamma\gamma'\}.$$

Then,

$$H(\omega) - H(\tilde{\omega}) = \begin{cases} L(\gamma\gamma') + (d_\gamma(\tilde{\omega}) + d_{\gamma'}(\tilde{\omega}))h_1 & \text{if } d_{\gamma'}(\tilde{\omega}) \geq 1, \\ L(\gamma\gamma') + d_\gamma(\tilde{\omega})h_1 - h_0 & \text{if } d_{\gamma'}(\tilde{\omega}) = 0. \end{cases}$$

Since  $0 < h_0 < h_1$ , we have  $H(\omega) - H(\tilde{\omega}) > 0$ , which contradicts that  $\omega$  is a ground state.

There are no edges in a ground state with length  $L$  greater than  $2h_0$ , since the energy of two monomers is  $2h_0 < L$ .

Now we prove existence of at least one ground state. Let  $(\Lambda_n)$  be a sequence of increasing cubes covering  $\mathbb{R}^d = \bigcup_n \Lambda_n$ . Let  $\omega_n$  be a configuration in  $\Lambda_n$  having the minimal energy over all configurations in  $\Lambda_n$ . There exists a subsequence  $(\omega'_i)$  of the sequence  $(\omega_n)$  (that is  $\omega'_i = \omega_{n_i}$ ) such that there exists a limit  $\lim_{i \rightarrow \infty} \omega'_i(e)$  for every  $e \in E$  (here  $w(e) = 1$  if  $e \in \omega$  and  $w(e) = 0$  otherwise). Moreover, the sequence  $(\omega'_i)$  can be chosen in such a way that  $\omega'_j(e) \equiv \text{const}$  for all  $j \geq i$  when  $e \in E_{\Lambda_i}$ . The configuration  $\omega' = \bigcup_i \omega'_i$  is one of the ground states. To see it, let  $\widehat{\omega}$  be a local perturbation of  $\omega'$ . There exists  $k$  such that  $\widehat{\omega} \Delta \omega' \subseteq E_{\Lambda_k}$ . For  $i > k$  let  $\widehat{\omega}_i$  be the restriction of  $\widehat{\omega}$  to  $E_{\Lambda_i}$ . Since  $\widehat{\omega}_i$  is a perturbation of  $\omega'_i$ ,

$$H_{\Lambda_i}(\widehat{\omega}_i) - H_{\Lambda_i}(\omega'_i) \geq 0.$$

Since no  $\omega_i$  has edges with a length greater than  $2h_0$ , there exists  $i_1 \geq i_0$  such that

$$H(\widehat{\omega}) - H(\omega') = H_{\Lambda_{i_1}}(\widehat{\omega}_{i_1}) - H_{\Lambda_{i_1}}(\omega'_{i_1}) \geq 0.$$

This proves that  $\omega'$  is a ground state. □

*Proof of Theorem 3:* Let  $\Gamma$  be a sample of  $\pi_\lambda$ , the law of a Poisson process with intensity  $\lambda$ . Let  $U$  be the union of all circles of radius  $h_0$  centered at the points of  $\Gamma$ . There exists a critical intensity  $\lambda_c$  such that for any  $\lambda < \lambda_c$  any maximal connected component of  $U$  is bounded  $\pi_\lambda$ -a.s.; see Theorem 3.3 in Ref. 5. Thus, for  $\lambda < \lambda_c$   $\pi_\lambda$ -almost-all  $\Gamma$  is a union of finite clusters  $\Gamma = \bigcup_{i=1}^\infty \Gamma_i$ ,  $|\Gamma_i| < \infty$ , and for any  $i \neq j$  and for any  $\gamma \in \Gamma_i, \gamma' \in \Gamma_j$  the distance  $|\gamma - \gamma'| > 2h_0$ . There are open edges only inside the clusters  $\Gamma_i$ . Since  $\Gamma_i$  are finite there exists a unique configuration in  $\Gamma_i$  minimizing the energy.

*The random connection model.* Choose a point configuration  $\Gamma$  with  $\pi_\lambda$  and then the edges with  $P^\circ$ . The resulting random graph is called *random-connection model* with rate  $\lambda$  and *connection function*

$$g(x) := \frac{e^{-\beta x}}{e^{-\beta x} + e^{-2\beta h_0}}. \tag{12}$$

See Chapter 6 in Ref. 5. The connection function  $g(x)$  is the probability that two points at distance  $x$  be connected. To show Theorem 4 we will dominate our graph with this model. The next lemma gives a sufficient condition for non-percolation in the random connection model.

*Lemma 3:* In the region (7) for  $\pi_\lambda$ -almost all  $\Gamma$  there is no percolation in the random-connection model with connection function (12).

*Proof:* Theorem 6.1 of Ref. 5 establishes that the random-connection model with connection function (12) does not percolate if

$$\lambda \int_0^\infty g(x) dx < 1. \tag{13}$$

For any  $\beta > 0$  and any  $h_0 > 0$  the integral of  $g(x)$  in (13) is finite and equals

$$\int_0^{2h_0} \frac{e^{-\beta x}}{e^{-\beta x} + e^{-2\beta h_0}} dx + \int_{2h_0}^\infty \frac{e^{-\beta x}}{e^{-\beta x} + e^{-2\beta h_0}} dx := J_1(\beta) + J_2(\beta).$$

The first integral on the right side of the above equality is increasing and tends to  $2h_0$  as  $\beta \rightarrow \infty$ . The second integral tends to 0 as  $\beta \rightarrow \infty$ . Fix  $\beta > 0$  and choose  $\lambda$  such that

$$\lambda < \frac{1}{2h_0 + J_2(\beta)} \leq \frac{1}{\int g(x) dx}.$$

We obtain the lemma by computing  $J_2(\beta) = \frac{\log 2}{\beta}$ . □

*Proof of Theorem 4:* Lemma 1 says that the Gibbs measure  $P$  is dominated by the product measure  $P^\circ$  defined by (9). Lemma 3 implies that the product measure  $P^\circ$  does not percolate under the conditions of the theorem. Thus the Gibbs measure  $P$  does not percolate for  $\mu_\lambda$ -a.s.  $\Gamma$ . □

*Lemma 4:* Let  $B_\gamma$  and  $D_\gamma$  be a partition of  $S_\gamma$ . Assume  $B_\gamma$  is finite and that  $\xi$  is a nonempty set of edges contained in  $B_\gamma$ . Then, for any  $e \in D_\gamma$ ,

$$P(e \in \omega \mid \omega \cap B_\gamma = \xi) \leq \exp(-\beta L(e) - \beta(h_1 - h_0)).$$

*Proof:* Consider any bounded  $\Lambda \subset \mathbb{R}^d$  such that  $\Lambda \supset B_\gamma \cup \{e\}$ . Let

$$A := \{\omega \subset E_\Lambda : \omega \cap B_\gamma = \xi, e \in \omega\},$$

$$Z := \sum_{\omega \subset E_\Lambda : \omega \cap B_\gamma = \xi} \exp(-\beta H_\Lambda(\omega))$$

and compute

$$\begin{aligned} P_\Lambda(e \in \omega \mid \omega \cap B_\gamma = \xi) &= \frac{1}{Z} \sum_{\omega \in A} \exp(-\beta H_\Lambda(\omega)) \\ &= \frac{1}{Z} \sum_{\omega \in A} \exp(-\beta(H_\Lambda(\omega) - H_\Lambda(\omega \setminus \{e\}))) \exp(-\beta H_\Lambda(\omega \setminus \{e\})) \\ &\leq \exp(-\beta L(e) - \beta(h_1 - h_0)) \frac{1}{Z} \sum_{\omega \in A} \exp(-\beta H_\Lambda(\omega \setminus \{e\})) \\ &\leq \exp(-\beta L(e) - \beta(h_1 - h_0)) P_\Lambda(e \notin \omega \mid \omega \cap B_\gamma = \xi) \\ &\leq \exp(-\beta L(e) - \beta(h_1 - h_0)). \end{aligned} \tag{14}$$

The event  $\{e \in \omega\}$  is cylindrical and so is  $\{\omega \cap B_\gamma = \xi\}$ , as  $B_\gamma$  is finite by hypothesis. Hence we can take the limit as  $\Lambda \uparrow \mathbb{R}^d$  and conclude.  $\square$

*Proof of Theorem 5:* Without losing generality we assume that the origin is a point of  $\Gamma$ ,  $0 \in \Gamma$ . Let  $C(\omega)$  be the vertex set of the graph  $C_0(\omega)$  and  $E(\omega)$  its edge set. Let us show that under the conditions of the theorem the expected number of vertices in the open cluster of the origin  $E|C|$  is finite. This is inspired in the method of generations introduced by Menshikov<sup>6</sup> to show no percolation in the site percolation model.

For a given configuration  $\omega$  let  $E(\omega)$  be the edge set of the open cluster  $C(\omega)$ , let  $V_0(\omega) = \{0\}$  and for  $n \geq 1$  define

$$V_n(\omega) := \{\gamma' \in \Gamma \setminus (V_0(\omega) \cup \dots \cup V_{n-1}(\omega)) : \text{there is a } \gamma \in V_{n-1}(\omega) \text{ such that } \gamma\gamma' \in E(\omega)\}, \tag{15}$$

$$O_n(\omega) := \{\gamma\gamma' \in E(\omega) : \gamma \in V_{n-1}(\omega), \gamma' \in V_{n-1}(\omega) \cup V_n(\omega)\}.$$

Each vertex in  $V_n(\omega)$  is attained with at least a path of  $n$  distinct open edges starting from the origin but cannot be attained with a shorter open path. Each edge in  $O_n(\omega)$  is the  $n$ -th step of an open self avoiding path starting from the origin and not belonging to a shorter open path. Since  $(V_n, n \geq 0)$  is a partition of the set of vertices of the open cluster at 0,

$$E|C| = \sum_{n=0}^{\infty} E|V_n|. \tag{16}$$

If we show that for  $\varepsilon$  defined in (8) and  $n \geq 0$ ,

$$E|V_{n+1}| \leq \varepsilon E|V_n|, \tag{17}$$

then the sum in (16) is bounded by  $\frac{\varepsilon}{1-\varepsilon}$  and the theorem is proven.  $\square$



*Proof of (17):* Call  $\tilde{O}_n(\omega) := O_1(\omega) \cup \dots \cup O_n(\omega)$ . The edges of the  $(n + 1)$ -th generation is the union of the edges belonging to the stars centered at the points of  $V_n$  and not contained in previous generations:

$$O_{n+1}(\omega) = \bigcup_{\gamma \in V_n(\omega)} (\omega \cap S_\gamma \setminus \tilde{O}_n(\omega)). \tag{18}$$

For  $n \geq 1$  define

$$F_n(\omega) := \{\gamma\gamma' \in E \setminus E(\omega) : \gamma \in V_{n-1}(\omega), \gamma' \in \Gamma\} \quad \square$$

the set of closed edges incident to the vertices in  $V_{n-1}$ . Let  $\tilde{F}_n := F_1 \cup \dots \cup F_n$ .

Let  $\xi$  be a possible set of open edges for the first  $n$  generations:  $\xi = \tilde{O}_n(\xi)$ . For any  $\omega$ , if  $\xi = \tilde{O}_n(\xi) = \tilde{O}_n(\omega)$ , then  $\tilde{F}_n(\xi) = \tilde{F}_n(\omega)$ . Since every vertex has finite degree, the set of possible  $\xi$  is countable and

$$\sum_{\xi} \mathbf{P}(\tilde{O}_n(\omega) = \xi) = 1,$$

where the sum runs over finite  $\xi \subset E$  such that  $\xi = \tilde{O}_n(\xi)$ .

Fix  $\gamma \in V_n(\xi)$  and let  $B_\gamma(\xi) := S_\gamma \cap (\tilde{O}_n(\xi) \cup \tilde{F}_n(\xi))$ , the set of edges incident to  $\gamma$  already determined by  $\xi$  and  $D_\gamma(\xi) := S_\gamma \setminus B_\gamma(\xi)$  the edges in  $S_\gamma$  that are not fixed by  $\xi$ . Since  $\xi$  is finite, so is  $B_\gamma(\xi)$ . Take  $e \in D_\gamma(\xi)$ . Then,

$$\begin{aligned} \mathbf{P}(e \in \omega \mid \tilde{O}_n(\omega) = \xi) &= \mathbf{P}(e \in \omega \mid \omega \cap B_\gamma(\xi) = \xi \cap B_\gamma(\xi)) \\ &\leq \exp(-\beta L(e) - \beta(h_1 - h_0)), \end{aligned}$$

by Lemma 4; indeed  $\xi \cap B_\gamma(\xi)$  is not empty. The obtained bound does not depend on  $\xi$ . Since  $|V_n(\xi)| \leq |O_n(\xi)|$  for all  $\xi$  and  $n \geq 1$ ,

$$\begin{aligned} \mathbf{E}(|V_{n+1}| \mid \tilde{O}_n(\omega) = \xi) &\leq \mathbf{E}(|O_{n+1}| \mid \tilde{O}_n(\omega) = \xi) \\ &= \sum_{\gamma \in V_n(\xi)} \sum_{e \in D_\gamma(\xi)} \mathbf{P}(e \in \omega \mid \tilde{O}_n(\omega) = \xi) \\ &\leq \sum_{\gamma \in V_n(\xi)} \sum_{e \in E_\gamma} \exp(-\beta L(e) - \beta(h_1 - h_0)) \\ &\leq T(\beta) \exp(-\beta(h_1 - h_0)) |V_n(\xi)| = \varepsilon |V_n(\xi)|. \end{aligned}$$

Hence,

$$\mathbf{E}|V_{n+1}| = \sum_{\xi} \mathbf{E}(|V_{n+1}| \mid \tilde{O}_n(\omega) = \xi) \mathbf{P}(\tilde{O}_n(\omega) = \xi) \leq \varepsilon \mathbf{E}|V_n|. \quad \square$$

### V. FINAL REMARKS

Since the ground state of Gibbs Random Graph does not percolate, the theorems about non-percolation show a kind of “stability” of the ground states.

Sufficient conditions for the existence of an infinite open cluster and monotonicity of  $\theta_\gamma$  as function of  $\beta$  (or  $(\lambda, \beta)$ ) are open problems.

### ACKNOWLEDGMENTS

The work of E.A.P. was partly supported by CNPq grants 300576/92-7 and 662177/96-7, (PRONEX) and FAPESP grant 99/11962-9, RFBR grants 07-01-92216 and 08-01-00105.

The work of V.V.S. was supported by FAPERJ (grants E-26/170.008/2008 and E-26/110.982/2008) and CNPq (grants 471891/2006-1, 309397/2008-1 and 471946/2008-7).

The work of A.Ya. was partly supported by FAPERJ grant E26/170.008/2008, and CNPq 306092/2007-7.

- <sup>1</sup>B. Bollobás, *Random Graphs*, 2nd ed. (Cambridge University Press, Cambridge, 2001), vol.73.
- <sup>2</sup>R. Durrett, *Random Graph Dynamics* (Cambridge University Press, Cambridge, 2007).
- <sup>3</sup>H.-O. Georgii, O. Häggström, and C. Maes, "The random geometry of equilibrium phases." *In phase transitions and critical phenomena*, Vol. 18. *Phase Transit. Crit. Phenom.*, vol. 18. Academic Press, San Diego, CA, 1–142.
- <sup>4</sup>G. R. Grimmett, and D. J. A. Welsh, *Disorder in Physical Systems*. edited by (The Clarendon Press Oxford University Press, New York, 1990).
- <sup>5</sup>R. Meester, and R. Roy, *Continuum percolation*. Cambridge Tracts in Mathematics (Cambridge University Press, Cambridge, 1996), vol. 119.
- <sup>6</sup>M. V. Men'shikov, S. A. Molchanov, and A. F. Sidorenko, (1986). Percolation theory and some applications. In *Probability theory. Mathematical statistics. Theoretical cybernetics*, Vol. 24 (Russian). *Itogi Nauki i Tekhniki. Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform.*, Moscow, 53–110, i. Translated in *J. Soviet Math.* **42**(4), 1766 (1988).