

# Distribution of spring phytoplankton (mainly diatoms) in the upper 50 m of the Southwestern Atlantic Ocean (30–61°S)

HÉCTOR F. OLGUÍN<sup>1,2\*</sup>, DEMETRIO BOLTOVSKOY<sup>1,2,3</sup>, CARINA B. LANGE<sup>4</sup> AND FREDERICO BRANDINI<sup>5</sup>

<sup>1</sup>FACULTAD DE CIENCIAS EXACTAS Y NATURALES, UNIVERSIDAD DE BUENOS AIRES, CIUDAD UNIVERSITARIA, C1428EHA-BUENOS AIRES, ARGENTINA,

<sup>2</sup>MUSEO ARGENTINO DE CIENCIAS NATURALES, A. GALLARDO 470, C1405DJR-BUENOS AIRES, ARGENTINA, <sup>3</sup>CONSEJO NACIONAL DE INVESTIGACIONES CIENTÍFICAS Y TÉCNICAS, ARGENTINA, <sup>4</sup>DEPARTAMENTO DE OCEANOGRAFÍA & CENTRO DE INVESTIGACIÓN OCEANOGRÁFICA EN EL PACÍFICO SUR-ORIENTAL (FONDAP-COPAS), UNIVERSIDAD DE CONCEPCIÓN, CASILLA 160-C-CONCEPCIÓN, CHILE AND <sup>5</sup>CENTRO DE ESTUDOS DO MAR, UNIVERSIDADE FEDERAL DO PARANÁ, AV. BEIRA MAR S/N, PONTAL DO SUL, PR 83255-000, BRAZIL

\*CORRESPONDING AUTHOR: holguin@ege.fcen.uba.ar

Received February 22, 2006; accepted in principle July 10, 2006; accepted for publication August 31, 2006; published online September 13, 2006

Communicating editor: K.J. Flynn

*This is the first study on diatom spatial patterns in relation to major oceanographic features along a megascale transect in the Southwestern Atlantic Ocean and provides a comparison with diatom distribution in surface sediments. Absolute abundances of diatoms, silicoflagellates and dinoflagellates (>10-µm fraction) were assessed in 80 bottle samples from 5 to 50 m, retrieved in November 1993 at 20 stations (30–61°S) along 53°W. Siliceous phytoplankton were scarce in the northern half of the transect and in the south of 57°S (100–150 cells L<sup>-1</sup>), with a strong peak in the vicinity of the Polar Front (~200 000 cells L<sup>-1</sup>), whereas dinoflagellates were more abundant at the northern stations (up to 24 000 cells L<sup>-1</sup>). In the south of 50°S phytoplanktonic cell densities were loosely (but significantly,  $r = 0.54$ ,  $P < 0.01$ ) associated with chlorophyll *a*, whereas in the north of this latitude, this relationship disappeared ( $r = 0.018$ ,  $P > 0.1$ ). In total, 191 diatoms and 4 silicoflagellates were recorded. Changes in diatom assemblage compositions along the transect allowed identification of five discrete areas: Subtropical (29°S), Northern Transitional (34–41°S), Southern Transitional (43–48°S), Subantarctic (49–54°S) and Antarctic (55–59°S), each characterized by a set of typical species. Diversity changed little with latitude, but numbers of species were higher in the north of 40°S. Comparison of diatom assemblage makeup in the plankton and in the surface sediments shows very strong disagreements, whereby cold water species are very significantly over-represented in the sedimentary record, suggesting enhanced preservation and strong subsurface equatorward advection of the cold water taxa.*

## INTRODUCTION

The zooplankton of the Southwestern Atlantic Ocean has been a subject of many focused surveys and at least two major compilation efforts (Boltovskoy, 1981, 1999). Biogeographically oriented investigations have chiefly dealt with the latitudinal distribution of species and definition of areas of enhanced faunal turnover as a function of local currents, water masses and fronts. Animal groups used in these studies included several protists (Boltovskoy, 1970; Balech, 1986; Boltovskoy *et al.*, 1996;

Thompson *et al.*, 1999, 2001; Thompson, 2001), as well as various multicellular taxa (Dadon and Boltovskoy, 1982; Mazzoni, 1983; Tarling *et al.*, 1995; Esnal, 1999; Esnal and Daponte, 1999; etc.). Phytoplankton, on the contrary, have received considerably less attention. Regional studies have mainly dealt with spatial distribution patterns of nutrients and chlorophyll (Brandini *et al.*, 2000; Barlow *et al.*, 2002), whereas data on the distribution and abundance of phytoplankton species, in particular diatoms, are restricted to a few circumscribed

investigations (Lange, 1985, 1988; Lange and Mostajo, 1985; Gayoso and Podestá, 1996; Fernandes and Brandini, 1999; Gayoso, 1999).

Field measurements of primary production and chlorophyll *a* indicate that the Southwestern Atlantic host areas of enhanced phytoplankton output suggesting that diatoms may play an important role in this biomass buildup (Hasle, 1969; Esteves *et al.*, 1992; Sullivan *et al.*, 1993; Gayoso and Podestá, 1996; Fernandes and Brandini, 1999; Brandini *et al.*, 2000; Marañón *et al.*, 2000; Barlow *et al.*, 2002). However, the lack of adequate spatial coverage in synoptic or quasi-synoptic surveys, associated with the extreme temporal and spatial variability of the physical and biological properties in the dynamic Brazil–Malvinas Confluence Zone (BMCZ), makes assessment of overall phytoplankton patterns very complicated.

Investigation of distributional ranges of the diatoms present in the water column is important for the assessment of the biological attributes of this complex area. Species-specific surveys allow defining functionally meaningful biogeographic zonations (Beklemishev, 1969; Boltovskoy, 1986), as well as assess the origin, path and degree of mixture of different currents and water masses (e.g. Balech, 1986). Furthermore, thorough understanding of present distributional patterns of fossilizable microplankton organisms is a necessary condition for the use of their remains in sedimentary samples for paleoceanographic studies in general and for paleoclimatic interpretations in particular (Boltovskoy, 1994; Eynaud *et al.*, 1999). In this respect, contrary to the widely accepted notion that, in most regions of the world ocean, the sedimentary imprint is an adequate proxy of water column flora, preliminary analyses of the diatoms preserved in bottom sediments of the Southwestern Atlantic suggest that the signal is strongly biased toward the cold water components (Romero and Hensen, 2002).


The main goal of this work is to assess the species-specific distribution of diatoms and silicoflagellates recorded in the upper 50 m of the water column along a N–S transect in the Southwestern Atlantic between 30 and 61°S. Densities of thecate dinoflagellates (but not species identifications) are also included. The spatial patterns defined are interpreted in terms of the dynamics and the physical and chemical characteristics of the currents and water masses that characterize the area, allowing the identification of several sectors of enhanced species change, some of which coincide with physical fronts described for the area. Distributional ranges of selected species and species groups are compared in detail with data from the surface sediments.

## Hydrology of the surface waters

The upper level circulation in the study area (47–57°W, 30–61°S) is dominated by two ocean currents flowing in opposite directions: the northbound Malvinas (Falklands) Current, which branches off the Antarctic Circumpolar Current (ACC or West Wind Drift), transporting cool, nutrient-rich Subantarctic waters along the western margin of the Argentine basin (Peterson and Stramma, 1991; Peterson, 1992), and the southbound Brazil Current (Fig. 1), which is warm, saline and relatively oligotrophic (Deacon, 1982; Peterson and Stramma, 1991). The area where the two come in contact with each other is the BMCZ, a very dynamic area between 30 and 46°S characterized by the interleaving and mixture of Subantarctic and Subtropical waters, as well as strong mesoscale features such as thermohaline fronts, rings, filaments and eddies (Olson *et al.*, 1988; Peterson, 1992; Mata and Garcia, 1996; Willson and Rees, 2000). Here, the Brazil Current detaches from the coast and starts meandering, turning first to the southeast (SE) and then to the northeast (NE) (Goni *et al.*, 1996), whereas the Malvinas Current makes a cyclonic loop returning to the SE.

The above-described hydrographic features define zones of enhanced horizontal gradients in several physical properties. Among the most conspicuous are the Subantarctic Front and the Polar Front (Fig. 1). The Subantarctic Front bounds the Malvinas Current up to ~40°S following the 2000 m isobath (Peterson, 1992). The Polar Front or Antarctic Convergence according to Deacon (Deacon, 1982), located between 50 and 60°S depending on season and longitude, defines the area where cold (up to ~4°C) waters of the ACC sink under the less dense Subantarctic surface waters and move north as Antarctic intermediate waters, probably as far north as 20–25°S (Lenz, 1975; Tsuchiya *et al.*, 1994). The area encompassed by the Subantarctic Front and the Polar Front is also known as the Polar Front Zone (Fig. 1).

In addition to these chiefly oceanic features, the area hosts several fronts associated with coastal processes. The Shelf Break Front is a stretch located along the Argentine continental shelf-break extending between ~37 and 50°S (Guerrero *et al.*, 1999; Acha *et al.*, 2004). It is a major oceanographic feature along the southern South American Atlantic margin defined by the contact between Subantarctic shelf waters and cooler and more saline waters of the Malvinas Current. Along this feature, active vertical mixing is responsible for enhanced primary production estimated at 350–450 g C m<sup>-2</sup> year<sup>-1</sup> (Negri, 1993) and the concomitant high concentrations of zooplankton (Ciechomski and Sánchez, 1983; Dadon, 1984).



**Fig. 1.** Location of sampling stations and general surface circulation pattern, modified from Peterson and Stramma (Peterson and Stramma, 1991).

At 35–36°S the mouth of the Rio de la Plata discharges  $\sim 22\,300\text{ m}^3\text{ s}^{-1}$  of fresh water into the sea (Guerrero *et al.*, 1997). The area influenced by fresh water runoff varies seasonally and geographically; during the winter, it extends up to 1300 km to the NE of the estuary, whereas in the summer, it mainly spreads offshore and along the coast reaching  $\sim 31^\circ\text{S}$  (Piola *et al.*, 2000).

## METHOD

Within the framework of the Tango Batucada International Agreement (TABIA) project, a joint Brazilian–Argentine research program based on chemical and plankton sampling in the South Atlantic, water bottle (Niskin) samples were collected at 20 stations along a N–S transect located at 47–57°W and 30–61°S (Fig. 1,

Table I), onboard RV ‘Barão de Teffé’, in November 1993. Discrete samples were collected at 5, 10, 25 and 50 m in the water column. For each depth, a volume of 3–8 L was fractionated through a 10- $\mu\text{m}$  mesh by reverse filtration and used for the estimation of the composition and abundance of the phytoplankton. Thus, all data reported here refer to the >10- $\mu\text{m}$  fraction. Samples were preserved with 5% neutralized formaldehyde in sea water. Diatom and silicoflagellate counts and identifications were performed under an inverted microscope equipped with phase-contrast illumination in Utermöhl settling chambers (Utermöhl, 1958). The number of transects scanned in each chamber varied with algal density; at least 300 cells were counted and identified for each major group reported here (only relative abundance data were available for station 1). Diatoms were also identified in subsamples processed to clean the frustules from organic matter (Simonsen, 1974) and mounted in permanent slides with Naphrax mounting medium. Diatom identifications were chiefly based on Hustedt (Hustedt, 1930), Cupp (Cupp, 1943), Simonsen (Simonsen, 1974), Hasle and Syvertsen (Hasle and Syvertsen, 1996) and Sar *et al.* (Sar *et al.*, 2001, 2002); for silicoflagellates, the work of Takahashi and Blackwelder (Takahashi and Blackwelder, 1992) was used.

Temperature, nutrient and chlorophyll *a* (by fluorometry after extraction with 90% acetone) data used in this work have been reported by Brandini *et al.* (Brandini *et al.*, 2000), where the corresponding methods are also detailed. The positions of the main fronts traversed during the cruise were defined in a previous work chiefly on the basis of surface and vertical temperature profiles (Brandini *et al.*, 2000).

Because the aim of this survey was assessing geographic (latitudinal) shifts in diatom assemblages in association with major hydrographic features, for most biogeographic analyses, the species-specific quantitative values of each vertical profile were pooled (weighted mean). This procedure was further justified by the fact that the overall structure of the database was very strongly governed by the latitudinal gradient, whereas depth had a minor influence on the composition of the assemblages. This assumption was confirmed by the results of a cluster analysis using all the data points (i.e. non-pooled data), where sample-to-sample similarities clearly followed a latitudinal gradient rather than vertical patterns (Fig. 2).

Cluster analyses were based on Jaccard’s index and the unweighted pair group method with arithmetic mean (UPGMA) algorithm (Romesburg, 1984).

Although all our taxonomic information (and, consequently, the biogeographic zonations proposed) is based on traditional, light-microscope systematics, we are aware of the fact that, in the last years, diatom taxonomy has

*Table I: Date, location of sampling stations, number of diatom species and phytoplankton (>10 µm) densities*

| Station | Date (November 1993) | Latitude S | Longitude W | Number of diatom species | Diatoms | Silicoflagellates | Thecate dinoflagellates |
|---------|----------------------|------------|-------------|--------------------------|---------|-------------------|-------------------------|
| 1       | 6                    | 29°42'06"  | 47°16'07"   | 81                       | ND      | ND                | ND                      |
| 2       | 10                   | 33°59'03"  | 51°14'12"   | 87                       | 56 234  | 325               | 1780                    |
| 3       | 11                   | 35°00'03"  | 51°23'36"   | 69                       | 519     | 175               | 521                     |
| 5       | 12                   | 37°35'12"  | 51°38'00"   | 54                       | 69      | 4                 | 15                      |
| 6       | 12                   | 37°15'06"  | 51°23'06"   | 36                       | 934     | 1454              | 13 405                  |
| 7       | 12                   | 37°01'54"  | 51°12'00"   | 68                       | 16 451  | 371               | 949                     |
| 8       | 13                   | 39°02'00"  | 52°07'00"   | 64                       | 18 647  | 193               | 1329                    |
| 9       | 14                   | 40°59'00"  | 52°40'00"   | 37                       | 161     | 4                 | 131                     |
| 10      | 15                   | 42°59'00"  | 53°08'06"   | 34                       | 766     | 61                | 205                     |
| 11      | 15                   | 45°00'12"  | 53°31'00"   | 45                       | 2030    | 100               | 945                     |
| 12      | 16                   | 47°00'00"  | 53°50'00"   | 37                       | 647     | 48                | 396                     |
| 13      | 17                   | 49°36'00"  | 54°12'00"   | 37                       | 7645    | 199               | 702                     |
| 14      | 18                   | 53°05'00"  | 54°52'00"   | 37                       | 2221    | 461               | 2007                    |
| 15      | 19                   | 53°54'18"  | 55°01'48"   | 50                       | 20 216  | 1651              | 1181                    |
| 16      | 20                   | 55°03'00"  | 55°14'00"   | 45                       | 4252    | 999               | 379                     |
| 17      | 20                   | 56°14'24"  | 55°30'54"   | 53                       | 31 718  | 1133              | 679                     |
| 18      | 20                   | 56°20'23"  | 55°25'24"   | 56                       | 67 984  | 6453              | 1055                    |
| 19      | 20                   | 57°15'42"  | 55°43'42"   | 46                       | 35 865  | 1297              | 661                     |
| 20      | 21                   | 59°14'00"  | 56°16'00"   | 43                       | 34 035  | 974               | 1111                    |
| 21      | 21                   | 61°00'00"  | 56°46'12"   | 40                       | 2713    | 90                | 90                      |

ND, no data.

Cell numbers (cells L<sup>-1</sup>) are mean values for integrated 0–50 m water column based on depth-weighted figures for 5, 10, 25 and 50 m.

undergone changes because of new evidence from molecular studies. Thus, at least some of the taxa identified in our work are probably species groups rather than valid species (e.g., Lundholm and Moestrup, 2002; Lundholm *et al.*, 2002, 2003; Sarno *et al.*, 2005). Despite these shortcomings, we contend that the taxonomic changes involved do not affect significantly biogeographic zonations derived from numerical analyses. The outcome of multivariate techniques are very resilient, and meaningful trends emerge even when the databases used contain significant amounts of noise derived chiefly from the spurious absences of the uncommon species (Romesburg, 1984). Using the smaller ranges of the intervening taxa, rather than the larger ones of the corresponding species groups, would most probably add detail to the pattern rather than modify the overall scheme radically.

## RESULTS AND DISCUSSION


### Temperature

Sea surface temperature (SST) varied between 24°C at the northern end of the transect and 1°C at the

south of the Polar Front (Fig. 3). SSTs were >18°C at stations 1–3, located in the north of the Brazil Current Front in Subtropical waters. At ~38°S, a strong drop in the SST and steepening of the isotherms indicated the presence of a lateral intrusion of colder waters. Temperatures decreased sharply within the BMCZ from 18°C at its northern end to ~7°C at 50°S, where the southern boundary of the BMCZ or Subantarctic Front (Peterson and Stramma, 1991; Saraceno *et al.*, 2004) or Falkland Escarpment Front (cf. Ikeda *et al.*, 1989) was identified. In Subantarctic waters of the Malvinas Current temperatures were homogeneous both horizontally and vertically, ranging between 7 and 5°C. The Polar Front was detected at ~56°S with SSTs decreasing sharply from 5 to 2°C and isotherms deepening to 200 m (Fig. 3 and Brandini *et al.*, 2000).

### Nutrients, chlorophyll *a* and phytoplankton densities

Nutrient concentrations along the transect (not shown) increased gradually from north to south, with low values in the upper oligotrophic waters of the Brazil Current




**Fig. 2.** Similarity between all samples investigated as shown by a cluster analysis based on their diatom inventories [Jaccard's index and unweighted pair group method with arithmetic mean (UPGMA)]. Notice that latitude, rather than depth, clearly structures the database (cf. Fig. 5).

and throughout most of the BMCZ and relatively high ones in the south of 56°S (see Fig. 3 in Brandini *et al.*, 2000).

Chlorophyll *a* concentrations varied between 0.14 and  $>3 \mu\text{g L}^{-1}$ ; peak values usually occurred in the upper 25 m of the water column, chiefly in the area of the BMCZ (36–46°S) and in the vicinity of the Polar Front (Fig. 3); no major peaks below 50 m were observed (see Fig. 2 in Brandini *et al.*, 2000). This pattern agrees with previous surveys (Sullivan *et al.*, 1993; Podestá, 1997; Marañón *et al.*, 2000). Highest value ( $3.58 \mu\text{g L}^{-1}$ ) were recorded at station 6 (37°15'); interestingly, the water temperature at this site was noticeably lower than both north and south of it (Fig. 3), and the plankton were dominated by cold water diatoms (*Asteromphalus hyalinus*, *Asteromphalus parvulus* and *Fragilariopsis ritscheri*), dinoflagellates (*Ceratium lineatum*, E. Balech, Buenos Aires, personal communication), tintinnids (Thompson *et al.*, 1999), foraminifers (Boltovskoy *et al.*, 1996), etc. These records suggest that the area was under the influence of laterally advected waters of the colder Malvinas Current (Brandini *et al.*, 2000). High chlorophyll *a* values may also have been enhanced by nutrients flushed out by the Río de la Plata estuary whose offshore influence was indicated by the presence of neritic diatoms (*Thalassiosira decipiens*) and fresh water Desmidiaceae.

With the exception of station 2, where diatoms were very abundant in the 5–25-m layer, phytoplankton were scarce in the northern half of the transect; concentrations of diatoms were  $\sim 5000\text{--}10\,000 \text{ cells L}^{-1}$ , whereas those of the silicoflagellates varied around  $100\text{--}150 \text{ cells L}^{-1}$  (Fig. 3). At 56°S, in the vicinity of the Polar Front, there was a very strong peak in diatom ( $181\,000 \text{ cells L}^{-1}$ ) and silicoflagellate abundances ( $10\,353 \text{ cells L}^{-1}$ ), both at station 18 at 5 m (Fig. 3), dropping again at the south of 57°S. As opposed to the silica-bearing cells, dinoflagellate densities were lower at the southern end of the transect than farther north; peak values for this group ( $24\,000 \text{ cells L}^{-1}$ ) occurred at station 6 (Fig. 3). In all cases, maximum densities were recorded in the uppermost 25 m. On average, diatoms accounted for 73% of all cells recorded (range: 6–96%); with lowest values at stations 6 and 9, where dinoflagellates and unidentified algae were dominant. Silicoflagellates represented on average 5.4% (0–18%) and dinoflagellates 17% (1.4–85%) of the numeric density.

Phytoplankton concentrations assessed in this work were generally lower than those reported by Fernandes and Brandini (Fernandes and Brandini, 1999) for the same area and by Gayoso and Podestá (Gayoso and Podestá, 1996) in the BMCZ, which is not surprising given the remarkable temporal and geographic variability in ambient conditions and plankton concentrations characteristic of the Confluence area (Gayoso and Podestá, 1996; Willson and Rees, 2000; Barlow *et al.*, 2002; Saraceno *et al.*, 2005); however, in all surveys, diatoms were largely dominant.



**Fig. 3.** Vertical profiles of temperature, chlorophyll *a* concentrations and cell numbers along the transect; chlorophyll data from Brandini *et al.* (Brandini *et al.*, 2000).

Peaks in pigment values ( $3.58 \mu\text{g L}^{-1}$  at station 6; Fig. 3) were characterized by moderate cell densities dominated by dinoflagellates ( $<25\,000 \text{ cells L}^{-1}$ ). Conversely, in areas with the highest cell numbers (stations 2 and 18), chlorophyll *a* values were low ( $<0.6 \mu\text{g L}^{-1}$ ; Fig. 3). Thus, when considering the entire transect, and especially in the north of  $50^\circ\text{S}$ , phytoplankton cell densities were uncoupled with concentrations of chlorophyll *a* ( $r = -0.003$ ;  $P > 0.1$ ). This lack of correlation in the north of  $50^\circ\text{S}$  may be because of the generally low association between phytoplanktonic biomass and chlorophyll *a* content (Kruskopf and Flynn, 2006). Alternatively, it may indicate that, in addition to the contribution of dinoflagellate pigment at station 6, nanophytoplankton ( $<10 \mu\text{m}$ ) represented the bulk of primary producers. In the south of  $50^\circ\text{S}$ , on the contrary, association between chlorophyll *a* and phytoplankton cell numbers was much higher ( $r = 0.54$ ;  $P < 0.01$ ), probably because the taxonomic composition of the phytoplankton was more homogeneous here than in the northern section, and/or because diatom cells  $>10 \mu\text{m}$  hosted most of the chlorophyll in this area. This pattern agrees with previous observations that, within the BMCZ, ubiquitous nanoplanktonic diatoms and flagellates are an important contributor to total chlorophyll, whereas in the north of  $35^\circ\text{S}$  picoplankton ( $<2 \mu\text{m}$ ) dominate phytoplankton biomass (Barlow *et al.*, 2002; Gibb *et al.*, 2000; Marañón *et al.*, 2000).

### Species composition and biogeographic patterns

In total, 191 diatom species were identified (Appendix), ranging between 69 (station 2, 10 m) and 9 (station 9 at 5 and 50 m); the overall mean was 31 diatom species per sample. Most of these, however, were represented by very few individuals; on average, for the entire collection six species accounted for 50% of all diatom cells counted (36 species to reach 90% of the individuals). *Fragilariopsis kerguelensis* was by far the most abundant diatom, both in terms of absolute density (average for the entire collection:  $\sim 4000 \text{ cells L}^{-1}$ ) and in terms of relative abundance (mean: 23%). In agreement with its high proportions in sedimentary materials of this area (Crosta *et al.*, 2005), in the south of  $47^\circ\text{S}$ , *F. kerguelensis* usually accounted for  $>30\%$  of the diatoms recorded. In terms of cell numbers, it was followed by *Thalassionema nitzschioides*, *Pseudonitzschia lineola*, *Chaetoceros rostratus*, *Leptocylindrus danicus*, etc. However, because throughout the collection densities were poorly correlated with occurrences, taxa most often present in the samples were often different from these: *Azpeitia tabularis* (present in 84% of the samples), *T. nitzschioides* (79%), *Corethron criophilum* (76%),

*F. kerguelensis* (75%), *Actinocyclus curvatus* (67%), etc. Thirty-six species were recorded only once.

Four silicoflagellate species were identified in the collection, of which *D. speculum* was the most abundant reaching over  $10\,000 \text{ cells L}^{-1}$  in the vicinity of the Polar Front (Fig. 3). To decrease numerical importance, it was followed by *Dictyocha messanensis*, *Distephanus pulchra* and *Dictyocha mandrai*, none of which exceeded  $700 \text{ cells L}^{-1}$ . *Distephanus pulchra* was restricted to the northern sector of the area surveyed (north of  $40^\circ\text{S}$ ), whereas the other three had a more cosmopolitan distribution, ranging from  $30$  to  $54\text{--}55^\circ\text{S}$  (*D. messanensis*, *D. mandrai*) and to  $61^\circ\text{S}$  (*D. speculum*). These distributional patterns are in general agreement with the ecological preferences of these species reported for the South Atlantic (Eynaud *et al.*, 1999) and for other areas of the World Ocean (Poelchau, 1976; Takahashi, 1987; Takahashi and Blackwelder, 1992). To establish biogeographic diatom-based zones along the transect, we performed a cluster analysis using depth-integrated diatom data (Fig. 5). The first-order change in species composition was located between stations 12 and 13, at  $\sim 48^\circ\text{S}$ . The northern area was further divided into three subzones: Subtropical (station 1), Transition North (stations 2–5 and 7–9) and Transition South (stations 6 and 10–12). The southern area showed two groups of stations: Subantarctic (stations 13–15) and Antarctic (stations 16–21).

To define diatom assemblages characteristic of each one of these geographic areas, we selected species according to the following criteria: (i) The relative abundance of the species should exceed 0.5% of all diatoms in at least one sample; (ii) the mean relative abundance of the species in the samples included in a particular geographical group should be  $>2$  times higher than that in the entire collection; (iii) the species should be present in  $>80\%$  of the samples of the geographical group it characterizes and (iv) the species should be present in  $<25\%$  of the samples of any other group. These criteria were fulfilled by 42 diatom taxa. In addition to these, we also selected a set of nine diatoms clearly associated with the northern warm waters, stations 1–9 (mean relative abundance at stations 1–9  $>1.4$  times higher than at stations 13–21; present in  $>75\%$  of stations 1–9 and  $<15\%$  of stations 13–21) and a set of seven diatoms associated with the cold southern waters (stations 13–21, criteria opposed to above). Finally, we also identified a set of nine cosmopolitan or widely distributed diatoms composed of those taxa that were present in  $>30\%$  of the samples of every geographical group defined (with the exception of the northernmost sample group which consisted of a single sample).

This procedure allowed pinpointing a total of 67 relatively abundant diatom species with reasonably

well-defined distributional ranges (Fig. 4). These 67 diatoms accounted for 72% of all the specimens identified. For the remaining 124 taxa, latitudinal distribution and abundance values were not consistent enough to derive their biogeographic affinities.

The results of these analyses are described as follows:

#### *Station 1 (Subtropical)*

A total of 81 diatom species were identified in this area. Twenty of them were restricted to station 1, but only two species were relatively abundant (>1% of the assemblage): *Chaetoceros danicus* and *Mastogloia rostrata*. The remaining 18 diatoms recorded at station 1 scarcely represented 3.6% of the cells identified in this sample. In addition to the above two, this area was characterized by 14 species most of which peaked at station 1 but were also present farther south, in the Northern Transitional area (stations 2–9). None of them were recorded in the south of 41°S (Fig. 4). These 16 characteristic species were generally represented by low numbers of individuals (0.2–4%) of the entire assemblage at station 1.

#### *Stations 2–5, 7–9 (Northern Transitional)*

In total, 119 diatom species were recorded in these samples, of which 20 were not found elsewhere; these restricted species were invariably of low abundance (mean proportion of the assemblages in these samples: 0.1%). Thirteen diatoms were found to adequately characterize this area; among these, *Chaetoceros contortus*, *Pseudo-nitzschia multiseries* and *C. rostratus* were particularly abundant (Fig. 4). With the exception of *Pseudo-nitzschia fraudulenta*, whose range extended to 55°S, these diatoms were practically absent in the south of 41°S.

#### *Stations 6, 10–12 (Southern Transitional)*

Although the area encompassed by these samples yielded 75 species, only one (*Coscinodiscus granii*) was restricted to it, and also a single diatom was found to characterize these samples (*Detonula pumila*) (Fig. 4).

#### *Stations 13–15 (Subantarctic)*

Only two species were restricted to the Subantarctic area (*Stephanopyxis turris* and *Thalassiosira allenii*) of 61 recorded in samples 13–15. No characteristic diatoms were identified according to the criteria defined above. *Eucampia antarctica* was particularly abundant in these samples (up to 14%), but it was also present in sizeable proportions in most Antarctic samples as well.

#### *Station 16–21 (Antarctic)*

Seventy-five diatoms were recorded in the Antarctic area; 20 of which were not found in the north of 53°S. Characteristic taxa included 12 species, mostly restricted

to the Antarctic and Subantarctic, but some with sporadic occurrences as far north as 36°S (Fig. 4).

Several species were not unequivocally associated with a single group of stations defined by the clustering technique but were clearly more abundant at either the northern or the southern part of the transect. Nine diatoms were consistently present in considerable numbers in samples 1–5 and 7–9 and absent from Subantarctic and Antarctic waters (labeled ‘warm water’ in Fig. 4), and seven other species depicted the opposite trend (labeled ‘cold water’ in Fig. 4).

Finally, nine more diatoms were present throughout the transect at 80% of the stations; several of these were among the most abundant species in the collection (e.g. *T. nitzschiioides*, *Nitzschia sicula*, etc.) (labeled ‘Cosmopolitan’ in Fig. 4).

In general terms, the ecological settings of the species recorded—as derived from their occurrence in our samples—agree well with previous information (Venrick, 1971; Simonsen, 1974; Hasle, 1976; Lange, 1985; Hasle and Syvertsen, 1996). Of the 63 species ascribed to warm and temperate waters by previous authors (Appendix), in our materials, 57 were either restricted to or very clearly peaked at stations of the northern half of the transect. *Nitzschia bicapitata* group and *N. sicula* peaked in warm and temperate waters as well but ranged as far south as 54–59°S, for which reason they were tentatively included in the cosmopolitan group (Fig. 4). The remaining four species (*Chaetoceros peruvianus*, *Lauderia annulata*, *S. turris* and *T. allenii*) in our materials had high or maximum relative abundances in Subantarctic or Antarctic waters; however, all four were very scarce in our samples (none exceeded 0.4% of the overall diatom assemblage at any given station), for which reason their distribution patterns cannot be adequately interpreted.

Similarly, very few of the 54 diatoms ascribed to the cold waters by previous authors had a conspicuous presence outside of the Subantarctic and Antarctic areas in our materials. The only notable exception is that of *A. tabularis*, which showed peaks in abundance around the Subantarctic Front but was rather abundant throughout the entire transect (Fig. 4). Hasle and Syvertsen (Hasle and Syvertsen, 1996) considered *A. tabularis* as a Subantarctic species; yet, its pattern in sediments of the South Atlantic indicates that it is cold tolerant but clearly associated with Tropical–Subtropical waters (Romero *et al.*, 2005).

For the 51 taxa regarded as cosmopolitan by previous authors, agreement with our results was somewhat lower. In our materials, *Actinocyclus octonarius*, *Chaetoceros affinis*, *Chaetoceros concavicornis*, *Coscinodiscus centralis*, *Coscinodiscus marginatus*, *Coscinodiscus radiatus*, *P. fraudulenta*, *P. multiseries*, *Pseudo-nitzschia pungens* and *Skeletonema costatum s.l.* were clearly associated with the northernmost warm waters




Fig. 4. Distribution of a selected group of diatom species and of the four silicoflagellates recorded in the area investigated. Notice that station sequence is modified to illustrate diatom assemblage zones.

(Fig. 4), whereas *P. lineola* was noticeably more abundant at the southern end of the transect. *Detonula pumila*, also cosmopolitan according to previous information (Hasle, 1976; Hasle and Syvertsen, 1996), was in our materials practically restricted to the Southern Transitional area. These disagreements are most probably because literature-based information on the ecological affinities of these species was often based on general review works (Hasle, 1976; Priddle and Fryxell, 1985; Medlin and Priddle, 1990; Hasle and Syvertsen, 1996) or on surveys from other areas (Simonsen, 1974), where different ecological settings may have led to a different interpretation of the environmental signal conveyed by their presence. Disagreements with the data of Romero and Hensen (Romero and Hensen, 2002) and Romero *et al.* (Romero *et al.*, 2005), in turn, respond to the blurring and distortion of the distribution pattern in sediments (see below).


The above-described biogeographic areas based on diatom assemblages reflect latitudinal changes in temperature and nutrients, as well as the location of fronts and water masses (Fig. 1). The assemblage recorded at station 1 (~30°S) is dominated by taxa typical of the Subtropical oligotrophic waters of the Brazil Current. Stations 2–12 are also clearly dominated by warm water diatoms, but the presence of cold water and cosmopolitan forms is here more conspicuous, especially in the southern part of the area (stations 10–12, Southern Transitional). As shown in Fig. 5, these stations are located in the BMCZ—Transition Zone or Subtropical–Subantarctic Convergence Zone (cf. Boltovskoy, 1970)—a large area of mixture of warm and cold waters with isolated cells and tongues of pure Subantarctic and Subtropical input (Boltovskoy, 1970; Boltovskoy *et al.*, 1999). Our data indicate that, at the time of the cruise, the transition from Subtropical to Subantarctic conditions was relatively smooth. However, sharp discontinuities in this trend were also present, as indicated by the dominance of cold water diatoms at station 6 (Figs 4 and 5), most probably because of lateral advection of colder Malvinas waters offshore (Fig. 3 and Brandini *et al.*, 2000). A similar cold water signal was also detected at this site in other surveys of the same collection, based on planktonic foraminifers (Boltovskoy *et al.*, 1996) and on tintinnids (Thompson, 2001). The southern limit of the Transition (BMCZ) was defined by the Subantarctic Front, located between our sampling stations 12 and 13 (Fig. 5). This front was characterized by a sharp decrease in temperature, a rise in the concentration of nutrients (Brandini *et al.*, 2000) and a strong species change whereby cold water taxa became clearly dominant in the assemblage (Fig. 5, right-hand panel). Interestingly, this area did not seem to host

characteristic species; most of the diatoms present in Subantarctic waters were also abundant in farther south or dwelled throughout the transect investigated (Fig. 4). Finally, in south of the Polar Front, a distinct Antarctic assemblage occurred; most of these species clearly peaked at the southernmost end but were also present, albeit usually in small proportions, in Subantarctic and Transitional waters (Fig. 4).

Comparison of the biogeographic zonation of the area based on our diatom data with those based on planktonic foraminifers (Boltovskoy *et al.*, 1996) and tintinnids (Thompson, 2001) recovered from the same samples shows very good overall agreement. All three groups indicate a sharp break between stations 1 and 2, and the Transitional–Subantarctic and Subantarctic–Antarctic boundaries are also either coincident or very close by (Fig. 6). On the contrary, ecological affinities of the species whose distributional ranges define the boundaries in question vary strongly between groups. Differences are particularly noticeable with the foraminifers: although diatom assemblages show a strong dominance of warm water cells from 30 to ~43°S (station 10, Fig. 5), warm water foraminifers drop to near-zero values around 37°S (Boltovskoy *et al.*, 1996). This difference is partly because foraminiferal assemblages are depth stratified, with deeper strata hosting higher proportions of cold water species than the surficial ones (Boltovskoy *et al.*, 1996). Alternatively, the higher lifespans of foraminifers, around 2–4 weeks (Spindler *et al.*, 1979; Hemleben *et al.*, 1989), as compared with those of the diatoms and the tintinnids, a few hours to 1–2 days (Guillard and Kilham, 1977; Burkill, 1982; Heinbokel, 1987) could facilitate extended expatriation through longer survival. This assumption is supported by the fact that, as mentioned above, cold water foraminifers have extended northward ranges at depth; it is conceivable that these specimens are expatriates that have already reached their sterile range but are still present in the water column because of their long lifespans.

### Specific diversity

Diatom diversity (as measured by the Shannon–Wiener index) and number of species were not associated with depth (Fig. 7) but rather followed a latitudinal pattern. The latitudinal variation in specific diversity depicted three more-or-less well-defined peaks: at 30°S (station 2), at 46°S (station 11) and at 56°S (stations 17 and 18) (Fig. 7, upper panel). Comparison of this pattern with the distribution of the total number of species recorded at each station shows that although the first diversity peak is clearly associated with a higher number of taxa at the northernmost end of the transect, increases in the



**Fig. 5.** Geographic zonation of the area based on the diatom depth-integrated data for the sites surveyed [Jaccard's index and unweighted pair group method with arithmetic mean (UPGMA)]. Curves on right-hand side of the graph show proportions of warm water (Subtropical, Northern Transitional and warm water in Fig. 4) and cold water (Antarctic and cold water in Fig. 4) individuals at each site.

number of species at the other two are very minor (Fig. 7, lower panel). These slight increases in species number combined with an increase in the evenness contributed to boosting diversity index values.

The north–south trend in diatom species numbers does not agree with the widely acknowledged pattern of more or less gradually decreasing species richness figures toward higher latitudes (Ricklefs and Schluter, 1993; Briggs, 1996; Hayek and Buzas, 1997; Thompson, 2004).


Furthermore, diatom values also differ from data for the zooplankton of the same area. On the basis of compiled data for 16 zooplankton groups (1432 species in total), Boltovskoy *et al.* (Boltovskoy *et al.*, 2003) calculated the numbers of species inhabiting the different biogeographic areas of the South Atlantic. For the zooplankton, species numbers drop gradually from the Central–Subtropical area to the Antarctic, where inventories are ~40% of those in the subtropics (Fig. 8). For the diatoms, on the

|    |    |    | Diatoms | Foraminifers | Tintinnids |              |
|----|----|----|---------|--------------|------------|--------------|
| 1  | 1  | 1  |         |              |            | Subtropical  |
| 2  | 2  | 2  |         |              |            |              |
| 3  | 3  | 3  |         |              |            |              |
| 4  | 4  | 4  |         |              |            |              |
| 5  | 5  | 5  |         |              |            |              |
| 6  | 6  | 6  |         |              |            |              |
| 7  | 7  | 7  |         |              |            | Transitional |
| 8  | 8  | 8  |         |              |            |              |
| 9  | 9  | 9  |         |              |            |              |
| 10 | 10 | 10 |         |              |            |              |
| 11 | 11 | 11 |         |              |            |              |
| 12 | 12 | 12 |         |              |            |              |
| 13 | 13 | 13 |         |              |            | Subantarctic |
| 14 | 14 | 14 |         |              |            |              |
| 15 | 15 | 15 |         |              |            |              |
| 16 | 16 | 16 |         |              |            |              |
| 17 | 17 | 17 |         |              |            | Antarctic    |
| 18 | 18 | 18 |         |              |            |              |
| 19 | 19 | 19 |         |              |            |              |
| 20 | 20 | 20 |         |              |            |              |
| 21 | 21 | 21 |         |              |            |              |


**Fig. 6.** Comparison of the biogeographic zonation of the area based on diatom data (this work) with those based on planktonic foraminifers (Boltovskoy *et al.*, 1996) and tintinnids (Thompson, 2001).

contrary, inventories in the south of the subtropics remain more or less stable and are even somewhat higher in the Antarctic than in the Subantarctic (Fig. 8).

These differences have interesting implications for biogeographic and paleobiogeographic or paleoecologic purposes. Phytoplanktonic fossilizable remains have often been considered better current and water mass indicators and paleoindicators because of their inability to survive for extended periods away from the photic layer (thus minimizing transport by subsurface currents), their higher abundances and because of their shorter lifespans (thus decreasing sterile expatriation phenomena) (e.g. Boltovskoy, 1994). Whereas highly diversified and very abundant diatoms seem to be less sensitive to environmental changes in traits such as diversity and number of species,



**Fig. 7.** Diatom specific diversity, as measured by the Shannon–Wiener index, log base 2 (Shannon and Weaver, 1949) and numbers of diatom species per station (all depths pooled) along the transect surveyed.



**Fig. 8.** Numbers of zooplankton species present in the different biogeographic areas of the South Atlantic, based on data from Boltovskoy *et al.* (Boltovskoy *et al.*, 1999), compared with similar figures for the diatoms recorded in the present survey. All figures are in relative values.

thus reducing the potential usefulness of these attributes. Analysis of these diversity attributes from repeated sampling along similar transects in several years may reveal if the observed trends are persistent in that region.

### Diatom assemblages in the water column versus in the surface sediments

Romero and Hensen (Romero and Hensen, 2002) investigated the diatom assemblages retrieved in 60 surface sediment samples, 47 of which are located almost directly under our stations 1–13 (Fig. 9). This allowed a close comparison of water column versus sedimentary compositional data aimed at checking whether for the diatoms, in this area, patterns in the sediments reflect those in the overlying plankton. Romero and Hensen (Romero and Hensen, 2002) identified a total of 194 diatom species, which compares very favorably with our inventory of 191 taxa. Also, the overall numbers of diatoms characteristic of warm waters, cold waters and neritic environments were very similar in the two collections. In addition, the occurrence of freshwater diatoms in both collections was

mainly restricted to the area of influence of the Rio de la Plata outflow.


However, on a species-to-species basis, of the 209 diatoms recorded in the sediments or the water column (considering the overlapping area only, Fig. 9), only 79 (38%) were present in both collections, whereas the remaining 121 were absent from either collection (Fig. 10).

Comparison of the dominant taxa also points to very significant differences. Table II shows that 18 diatoms are necessary to account for the 10 most abundant forms in both databases. With the exception of *T. nitzschioides*, which ranked second both in the plankton and in the sediments, the remaining nine taxa varied widely in relative abundance and in rank (Table II). A similar disparity is observed when comparing the mean proportions of the taxa recorded in both collections (Fig. 11).

Although the above differences point at strong dissimilarities between the two data sets, they do not show that the temperature signal conveyed by the assemblage on the seafloor differs from the one in the water column. To assess this difference, we calculated the proportions of cells belonging to warm water and cold water species along the transect occupied. Because the locations of plankton and sediment samples are not identical, within each collection, we pooled the data in 5° latitude bins (Fig. 12). In the sediments between 36 and 41°S, the cold water signal is over twice as strong as the warm water one. In contrast, in the plankton, cold water diatoms are most abundant only in the south of 46°S (Fig. 5), whereas in the north of 41°S, the warm water signal is >6 and up to 4000 times stronger than the cold water one (Fig. 12). Although the two most abundant Antarctic diatoms in the sediments (*F. kerguelensis* and *Thalassiosira lentiginosa*) account for over half of all diatoms as far north as 35°S (Fig. 13), in the water column, they are only important in the south of 47°S.

Despite the temporal disparity of biological and sedimentary collections, our results indicate that sedimentary assemblages are a very poor proxy of diatom composition in the photic layer and that, along the entire transect, the sedimentary signal is very strongly biased toward colder waters. Romero and Hensen (Romero and Hensen, 2002) noticed this mismatch and stated that the diatom imprint in core top sediments mainly represents an Antarctic-originated signal, whereas *in situ* produced specimens supply a minor portion of the siliceous remains. This Antarctic signal can be traced as far north as 27°S, with *F. kerguelensis*, *T. lentiginosa* and the resting stage of *Thalassiosira antarctica* as indicative of the intensity of the equatorward displacement of Antarctic source water masses (Romero and Hensen, 2002).

Comparison of the planktonic and the sedimentary assemblages indicates that, as suggested by Romero and Hensen (Romero and Hensen, 2002), selective dissolution



**Fig. 9.** Location of the water column samples used in this survey and the sedimentary materials where diatoms were analyzed by Romero and Hensen (Romero and Hensen, 2002). Figure does not include sedimentary samples located too far east or north from the water column transect and therefore not used for comparative analyses.




Fig. 10. Mean proportions in the water column samples (this work) of the species absent in the sediments (Romero and Hensen, 2002) and vice versa (rs, resting spores).

Table II: Comparison of the overall percentage contributions and ranks of the 10 most abundant diatom species in the water column (this study) and the surface sediments (Romero and Hensen, 2002)


|                                      | Water column (%) | Sediment (%) | Water column (rank) | Sediment (rank) |
|--------------------------------------|------------------|--------------|---------------------|-----------------|
| <i>Fragilariopsis doliolus</i>       | 10.4             | 1.5          | 1                   | 13              |
| <i>Thalassionema nitzschioides</i>   | 9.9              | 16.7         | 2                   | 2               |
| <i>Fragilariopsis kerguelensis</i>   | 8.5              | 20.6         | 3                   | 1               |
| <i>Chaetoceros rostratus</i>         | 8.2              | NR           | 4                   | –               |
| <i>Nitzschia sicula</i>              | 5                | 0.01         | 5                   | 143             |
| <i>Thalassiosira decipiens</i>       | 4.8              | NR           | 6                   | –               |
| <i>Chaetoceros convolutus</i>        | 4.3              | NR           | 7                   | –               |
| <i>Rhizosolenia setigera</i>         | 4                | 0.1          | 8                   | 76              |
| <i>Leptocylindrus danicus</i>        | 3.9              | NR           | 9                   | –               |
| <i>Pseudo-nitzschia multiseriata</i> | 3.2              | NR           | 10                  | –               |
| <i>Azpeitia tabularis</i>            | 1.8              | 3.1          | 14                  | 6               |
| <i>Actinocyclus octonarius</i>       | 0.6              | 2.6          | 30                  | 8               |
| <i>Roperia tessellata</i>            | 0.6              | 2            | 31                  | 10              |
| <i>Cyclotella litoralis</i>          | 0.24             | 2.8          | 46                  | 7               |
| <i>Thalassiosira lentiginosa</i>     | 0.21             | 5.5          | 52                  | 3               |
| <i>Paralia sulcata</i>               | 0.01             | 3.1          | 121                 | 5               |
| <i>Thalassiosira antarctica</i>      | NR               | 5.3          | –                   | 4               |
| <i>Chaetoceros</i> sp.               | –                | 2.5          | –                   | 9               |

NR, not recorded.


plays an important role in these disagreements. Antarctic diatoms (e.g. *F. kerguelensis*, *T. lentiginosa* and several others) are known to have heavily silicified frustules and high preservation potential (Crosta *et al.*, 2005), which contributes to enhance their proportions in mixed warm water and cold water sedimentary assemblages. Seasonal variability may also play a role in these mismatching patterns insofar as much of the area transected is known to undergo important seasonal shifts, especially between 30 and 35°S (Boltovskoy, 1970). Equatorward displacement of settling frustules by northbound subsurface and deep currents has been proposed by several authors (Burckle and Stanton, 1975; Burckle and Cooke, 1983; Jones and Johnson, 1984) as an important distorting effect in the study area as well. However, in the Southwestern Atlantic southward-moving subsurface currents are probably as important as the northward ones (Reid *et al.*, 1977). Thus, the resulting displacement of a random settling particle may have an important southward component as well. We suggest that the predominantly northward displacement reflected in the plankton versus sediments mismatching pattern is largely because of a complex interaction where biological factors play an important role. Planktonic organisms carried equatorward by surface or subsurface currents can sink in the water column in search of a colder stratum where they can extend their survival. Subsequent displacements will force the organism to

continue descending (if equatorward expatriation persists), allow it to return to the surface (if it is reversed) or perish and sediment (if its tolerance threshold is exceeded). Autotrophic organisms, which depend on light for extended survival and have often a more limited capacity to maintain a given depth range, can be subject to the same phenomenon through grazing by zooplankton. As opposed to expatriation toward the equator, poleward advection of plankters should not result in a descent in the water column (as it would not contribute to extended survival), thus minimizing the occurrence of warm-water remains outside of their living ranges (Boltovskoy, 1988, 1994).


Although for most groups of pelagic organisms, distributional studies are based on samples of their living assemblages, for some provided with hard, fossilizable skeletal structures, including the diatoms, their remains in the sediments are often the main source of distributional information. As a matter of fact, for some groups, most of the information available to date, in particular on their specific distribution patterns, is based on sedimentary materials, whereas plankton-based studies are very scarce and fragmentary. This is partially because of the very significant input of geologically oriented investigations in this field but also largely because comprehensive information on the distribution of these organisms is more readily, easily and economically obtained from bottom sediments than from plankton collections. Indeed, sedimentary



**Fig. 11.** Differences in the mean relative abundance of the diatom species in the water column (this work) and in the sediments (Romero and Hensen, 2002) of the Southwestern Atlantic Ocean. Graph includes species recorded in both collections only.



**Fig. 12.** Proportions of warm water and cold water diatoms at different latitudinal offsets along the transect surveyed in the 0–50 m layer water column (our data) and in the surface sediments, data from Romero and Hensen (Romero and Hensen, 2002). Station 10 (our material) was set apart by the fact that it hosted unexpectedly large proportions of warm water diatoms, most probably because it sampled a Subtropical eddy.



**Fig. 13.** Comparison of the proportions of the two most abundant cold water diatoms in bottom sediment samples, data from Romero and Hensen (Romero and Hensen, 2002) and in the water column (this work).

materials have some advantages over water column samples, including larger sample sizes, better availability and the provision of time-integrated data (Boltovskoy, 1994), which contributed to their wide use in biogeographic surveys. Furthermore, sedimentary materials are widely used in paleobiogeographic and paleoclimatic investigations; much of our current knowledge of glacial to interglacial climatic changes in the open ocean are based on the interpretation of differences between recent versus last glacial maximum distribution patterns of coccolithophorids, diatoms, foraminifers and radiolarians (e.g. CLIMAP Project Members, 1976). Use of surface sediment materials (rather than water column data) is validated by the implicit or explicit assumption that patterns in the sediments, while not identical to those in the overlying waters, are a very good proxy of the latter. However, this assumption has seldom been explicitly addressed and demonstrated (Boltovskoy, 1994). Our results show that, in the Southwestern Atlantic Ocean, diatom distribution patterns in the sediments differ very significantly from those in the plankton, and the distortions involved are similar to those recorded for the foraminifers in the same area (Boltovskoy *et al.*, 1996), and for other planktonic organisms elsewhere (Boltovskoy, 1994).

## ACKNOWLEDGEMENTS

Thanks are due to crew members on board the Brazilian RV 'Barão de Teffé' for logistical and technical assistance. This work was supported by grants from the University of Buenos Aires, Argentina (UBA EX-040)

and from the Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brazil (CNPq PROC N°670001/93-7).

## REFERENCES

- Acha, M. E., Mianzan, H. W., Guerrero, R. A., Favero, M. and Bava, J. (2004) Marine fronts at the continental shelves of South America. Physical and ecological processes. *J. Mar. Syst.*, **44**, 83–105.
- Balech, E. (1986) De nuevo sobre la oceanografía frente a la Argentina. Publ. Serv. Hidr. Nav. Argentina, **H 645**, 1–23.
- Barlow, R. G., Aiken, J., Holligan, P. M., Cummings, D. G., Maritorena, S. and Hooker, S. (2002) Phytoplankton pigment and absorption characteristics along meridional transects in the Atlantic Ocean. *Deep-Sea Res. I*, **47**, 637–660.
- Beklemishev, C. W. (1969) *Ekologiya i biogeografiya pelagiali*. Nauka, Moskva.
- Boltovskoy, E. (1970) Masas de agua (características, distribución, movimientos) en la superficie del Atlántico sudoeste, según indicadores biológicos – Foraminíferos. *Serv. Hidrogr. Naval, Argentina*, **H 643**, 1–99.
- Boltovskoy, D. (ed.) (1981) *Atlas del Zooplankton del Atlántico Sudoccidental y Métodos de Trabajo Con el Zooplankton Marino*. Instituto Nacional de Investigación y Desarrollo Pesquero, Mar del Plata.
- Boltovskoy, D. (1986) Biogeography of the Southwestern Atlantic: current problems and prospects. In Pierrot-Bults, A. C., Van der Spoel, S., Zahuranec, B. J. and Johnson, R. K. (eds), *Pelagic Biogeography. UNESCO Technical Paper Marine Science 49*. UNESCO Press, Paris, pp. 14–24.
- Boltovskoy, D. (1988) Equatorward sedimentary shadows of near-surface oceanographic patterns. *Specul. Sci. Technol.*, **11**, 219–232.
- Boltovskoy, D. (1994) The sedimentary record of pelagic biogeography. *Prog. Oceanogr.*, **34**, 135–160.
- Boltovskoy, D. (ed.) (1999) *South Atlantic Zooplankton*. Backhuys Publishers, Leyden.
- Boltovskoy, E., Boltovskoy, D., Correa, N. and Brandini, F. (1996) Planktonic foraminifera from the southwestern Atlantic (30 to 60°S): species-specific patterns in the upper 50 m. *Mar. Micropaleontol.*, **28**, 53–72.
- Boltovskoy, D., Correa, N. and Boltovskoy, A. (2003) Marine zooplanktonic diversity: a view from the South Atlantic. *Oceanol. Acta*, **25**, 271–278.
- Boltovskoy, D., Gibbons, M. J., Hutchings, L. and Binet, D. (1999) General biological features of the South Atlantic. In Boltovskoy, D. (ed.), *South Atlantic Zooplankton*. Backhuys Publishers, Leiden, pp. 1–42.
- Brandini, F. P., Boltovskoy, D., Piola, A., Kocmur, S., Röttgers, R., Abreu, P. C. and Mendes Lopes, R. (2000) Multiannual trends in fronts and distribution of nutrients and chlorophyll in the southwestern Atlantic (30–62°S). *Deep-Sea Res. I*, **47**, 1015–1033.
- Briggs, J. C. (1996) *Global Biogeography*. Elsevier, Amsterdam.
- Burckle, L. H. and Cooke, D. W. (1983) Late Pleistocene *Eucampia antarctica* abundance stratigraphy in the Atlantic Sector of the Southern Ocean. *Micropaleontology*, **29**, 6–10.
- Burckle, L. H. and Stanton, D. (1975) Distribution of displaced diatoms in the Argentina Basin. *Nova Hedwigia Beih.*, **53**, 283–292.
- Burkill, P. (1982) Ciliates and other microplankton components of a nearshore food-web: standing stock and production processes. *Ann. Inst. Oceanogr. Paris*, **58**, 335–350.

- Ciechowski, J. D. and Sánchez, R. P. (1983) Relation between ichthyoplankton abundance and associated zooplankton biomass in shelf waters off Argentina. *Biol. Oceanogr.*, **3**, 77–101.
- CLIMAP Project Members (1976) The surface of the ice-age earth. Quantitative geologic evidence is used to reconstruct boundary conditions for the climate 18,000 years ago. *Science*, **191**, 1131–1144.
- Crosta, X., Romero, O., Armand, L. K. and Pichon, J. J. (2005) The biogeography of major diatom taxa in Southern Ocean sediments: 2. Open ocean related species. *Palaeogeogr. Palaeoclimatol.*, **223**, 66–92.
- Cupp, E. E. (1943) Marine plankton diatoms of the west coast of North America. *Bull. Scripps Inst. Oceanogr.*, **5**, 1–238.
- Dadon, J. R. (1984) Distribución y abundancia de Pteropoda Thecosomata en el Atlántico Sudoccidental. *Physis A*, **42**, 25–38.
- Dadon, J. R. and Boltovskoy, D. (1982) Zooplanktonic recurrent groups (Pteropoda, Euphausiacea, Chaetognatha) in the southwestern Atlantic Ocean. *Physis A*, **41**, 63–83.
- Deacon, G. E. R. (1982) Physical and biological zonation in the Southern ocean. *Deep-Sea Res.*, **29**, 1–15.
- Esnal, G. B. (1999) Appendicularia. In Boltovskoy D. (ed.), *South Atlantic Zooplankton*. Backhuys Publishers, Leiden, pp. 1375–1399.
- Esnal, G. B. and Daponte, M. C. (1999) Salpida. In Boltovskoy, D. (ed.), *South Atlantic Zooplankton*. Backhuys Publishers, Leiden, pp. 1423–1444.
- Esteves, J. L., Santinelli, N., Sastre, V., Díaz, R. and Rivas, O. (1992) A toxic dinoflagellate bloom and PSP production associated with upwelling in Golfo Nuevo, Patagonia, Argentina. *Hydrobiologia*, **242**, 115–122.
- Eynaud, F., Giraudeau, J., Pichon, J. J. and Pudsey, C. J. (1999) Sea-surface distribution of coccolithophores, diatoms, silicoflagellates and dinoflagellates in the South Atlantic Ocean during the late austral summer 1995. *Deep-Sea Res. I*, **46**, 451–482.
- Fernandes, L. F. and Brandini, F. P. (1999) Microplankton communities in the Southwestern Atlantic Ocean: biomass and distribution in November/1992. *Rev. Bras. Oceanogr.*, **47**, 189–205.
- Gayoso, A. M. (1999) Seasonal succession patterns of phytoplankton in the Bahía Blanca Estuary. *Bot. Mar.*, **42**, 367–375.
- Gayoso, A. M. and Podestá, G. P. (1996) Hydrography and phytoplankton of the Brazil-Malvinas Currents Confluence. *J. Plankton Res.*, **18**, 941–951.
- Gibb, S. W., Barlow, R. G., Cummings, D. G., Rees, N. W., Trees, C. C., Holligan, P. and Suggett, D. (2000) Surface phytoplankton pigment distributions in the Atlantic Ocean: an assessment of basin scale variability between 50°N and 50°S. *Prog. Oceanogr.*, **45**, 339–368.
- Goni, G., Kamholz, S., Garzoli, S. and Olson, D. (1996) Dynamics of the Brazil-Malvinas Confluence based on inverted echo sounders and altimetry. *J. Geophys. Res.*, **101**, 16273–16289.
- Guerrero, R. A., Baldoni, A. and Benavides, H. R. (1999) Oceanographic conditions at the southern end of the Argentine continental slope. *INIDEP Doc. Cient. (Mar Del Plata)*, **5**, 7–22.
- Guerrero, R. A., Lasta, C., Acha, M., Mianzan, H. and Framiñan, M. (1997) *Atlas Hidrográfico del Río de la Plata*. Comisión Administradora del Río de la Plata, Buenos Aires.
- Guillard, R. R. L. and Kilham, P. (1977) The ecology of marine planktonic diatoms. In Werner, D. (ed.), *The Biology of Diatoms. Botanical Monographs*. Vol. 13. Blackwell Science Publishers, London, 372–469.
- Hasle, G. R. (1969) An analysis of the phytoplankton of the Pacific Southern Ocean: abundance, composition and distribution during the Bratæg Expedition 1947–1948. *Hvalraed. Skr.*, **52**, 1–168.
- Hasle, G. R. (1976) The biogeography of some marine planktonic diatoms. *Deep-Sea Res.*, **23**, 319–338.
- Hasle, G. R. and Syvertsen, E. E. (1996) Marine diatoms. In Tomas, C. R. (ed.), *Identifying Marine Phytoplankton*. Academic Press, San Diego, California, pp. 5–385.
- Hayek, L. A. and Buzas, M. (1997) *Surveying Natural Populations*. Columbia University Press, New York.
- Heinbokel, J. F. (1987) Diel periodicities and rates of reproduction in natural population of tintinnines in the oligotrophic water off Hawaii, September 1982. *Mar. Microb. Food Webs*, **2**, 1–13.
- Hemleben, C., Spindler, M. and Anderson, O. R. (1989) *Modern Planktonic Foraminifera*. Springer Verlag, New York.
- Hendey, N. I. (1964) *An Introductory Account of the Smaller Algae of British Coastal Waters. Part V: Bacillariophyceae (Diatoms)*. Ministry of Agriculture, Fisheries and Food, Fishery Investigations Series IV. HMSO, London.
- Hustedt, F. (1930) Die Kieselalgen Deutschlands, Österreichs und der Schweiz. In Rabenhorst, L. (ed.), *Kryptogamen-Flora von Deutschlands, Österreichs und der Schweiz*. Akademische Verlagsgesellschaft, Leipzig, pp. 1–920.
- Hustedt, F. (1958) Diatomeen aus der Antarktis und dem Südatlantik. Deutsche Antarktische Expedition **1938/39**, **2**, 103–191.
- Ikeda, Y., Siedler, G. and Zwirz, M. (1989) On the variability of Southern Ocean Front locations between Southern Brazil and the Antarctic Peninsula. *J. Geophys. Res.*, **94**, 4757–4762.
- Jones, G. A. and Johnson, D. A. (1984) Displaced Antarctic diatoms in Vema Channel sediments: Late Pleistocene/Holocene fluctuations in AABW flow. *Mar. Geol.*, **58**, 165–186.
- Kruskopf, M. and Flynn, K. J. (2006) Chlorophyll content and fluorescence responses cannot be used to gauge reliably phytoplankton biomass, nutrient status or growth rate. *New Phytol.*, **169**, 525–536.
- Lange, K. B. (1985) Spatial and seasonal variations of diatom assemblages of the Argentinean coast (South Western Atlantic). *Oceanol. Acta*, **8**, 361–370.
- Lange, K. B. (1988) *Variaciones estacionales de las asociaciones de diatomeas planctónicas de dos áreas marinas, el Atlántico Sudoccidental y el Pacífico Nordoriental*. Thesis Doctoral. University of Buenos Aires., 142 pp.
- Lange, K. B. and Mostajo, E. L. (1985) Phytoplankton (diatoms and silicoflagellates) from the South Atlantic Ocean. *Bot. Mar.*, **28**, 469–476.
- Lenz, W. (1975) Untersuchungen zur inneren hydrographischen Struktur des sudatl. Und mittleren Atlantiks (0–2000 m Tiefe) mit zoogeographischen Anmerkungen. *Ber. Deutsch. Wiss. Kömm. Meeresforsch.*, **24**, 1–21.
- Lundholm, N., Hasle, G. R., Fryxell, G. A. and Hargraves, P. E. (2002) Morphology, phylogeny and taxonomy of species within the *Pseudo-nitzschia americana* complex (Bacillariophyceae) with descriptions of the two new species *P. brasiliana* and *P. lineae*. *Phycologia*, **41**, 480–497.
- Lundholm, N. and Moestrup, Ø. (2002) The marine diatom *Pseudo-nitzschia galaxiae* sp. nov. (Bacillariophyceae), morphology and phylogenetic relationships. *Phycologia* **41**, 594–605.
- Lundholm, N., Moestrup, Ø., Hasle, G. R. and Hoef-Emden, K. (2003) A study of the *Pseudo-nitzschia pseudodelicatissima*/cuspidata complex (Bacillariophyceae): what is *P. pseudodelicatissima*? *J. Phycol.*, **39**, 797–813.
- Marañón, E., Holligan, P. M., Varela, M., Mourinho, B. and Bale, A. J. (2000) Basin-scale variability of phytoplankton biomass, production and growth in the Atlantic Ocean. *Deep-Sea Res. I*, **47**, 825–857.
- Mata, M. M. and Garcia, C. A. E. (1996) Variabilidade da topografia oceânica superficial no Atlântico Sul Ocidental observada pela

- altimetria TOPEX/POSEIDON. *Anais do VIII Simpósio Brasileiro de Sensoriamento Remoto*, T67, Salvador, Bahia, Brasil.
- Mazzoni, H. (1983) Abundancia y distribución de Chaetognatha en el Atlántico Sudoccidental (Mar Argentino). *Physis A*, **41**, 157–171.
- Medlin, L. K. and Priddle, J. (1990) *Polar Marine Diatoms*. British Antarctic Survey, Cambridge.
- Negri, R. (1993) Fitoplancton y producción primaria en el área de la plataforma bonaerense próxima al talud continental. *Inf. Técn. INIDEP (Mar Del Plata)*, **1**, 1–7.
- Olson, D. B., Podestá, G. P., Evans, R. H. and Brown, O. B. (1988) Temporal variations in the separation of Brazil and Malvinas currents. *Deep Sea Res.*, **35**, 1971–1990.
- Peterson, R. G. (1992) The boundary currents in the western Argentine Basin. *Deep-Sea Res.*, **39**, 623–644.
- Peterson, R. G. and Stramma, L. (1991) Upper-level circulation in the South Atlantic Ocean. *Prog. Oceanogr.*, **26**, 1–73.
- Piola, A. R., Campos, E. J. D., Möller, O. O. Jr., Charo, M. and Martínez, C. (2000) The Subtropical Shelf Front off eastern South America. *J. Geophys. Res.*, **105**, 6565–6578.
- Podestá, G. P. (1997) Utilización de datos satelitarios en investigaciones oceanográficas y pesqueras en el Océano Atlántico Sudoccidental. In Boschi, E. (ed.), *El Mar Argentino y sus Recursos Pesqueros Tomo 1*. Instituto Nacional de Investigación y Desarrollo Pesquero, Mar del Plata, Argentina, pp. 195–222.
- Poelchau, H. (1976) Distribution of Holocene silicoflagellates in North Pacific sediments. *Micropaleontology*, **22**, 164–193.
- Priddle, J. and Fryxell, G. (1985) *Handbook of the Common Plankton Diatoms of the Southern Ocean*. British Antarctic Survey, Cambridge.
- Reid, J. L., Nowlin, W. D. and Patzert, W. C. (1977) On the characteristics and circulation of the Southwestern Atlantic Ocean. *J. Phys. Oceanogr.*, **7**, 62–91.
- Ricklefs, R. E. and Schluter, D. (eds) (1993) *Species Diversity in Ecological Communities*. University of Chicago Press, Chicago.
- Rines, J. E. B. (1999) Morphology and taxonomy of *Chaetoceros contortus* Schütt 1895, with preliminary observations on *Chaetoceros compressus* Lauder 1864 (Subgenus Hyalochaete, Section Compressa). *Bot. Mar.*, **42**, 539–551.
- Romero, O., Armand, L. K., Crosta, X. and Pichon, J. J. (2005) The biogeography of mayor diatom taxa in Southern Ocean surface sediments: 3. Tropical/Subtropical species. *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, **223**, 49–65.
- Romero, O. and Hensen, C. (2002) Oceanographic control of biogenic opal and diatoms in surface sediments of the southwestern Atlantic. *Mar. Geol.*, **186**, 263–280.
- Romesburg, H. C. (1984) *Cluster Analysis for Researchers*. Lifetime Learning Publications, Belmont.
- Sar, E. A., Sunesen, I. and Castañón, C. (2001) Marine diatoms from Buenos Aires coastal waters (República Argentina). I. Thalassiosiraceae. *Nova Hedvigia* **73**, 199–228.
- Sar, E. A., Sunesen, I. and Lavigne, A. S. (2002) The diatom genus *Thalassiosira*: species from the northern San Matías Gulf (Río Negro, Argentina). *Nova Hedvigia*, **74**, 373–386.
- Saraceno, M., Provost, C. and Piola, A. (2005) On the relationship between satellite retrieved surface temperature fronts and chlorophyll-a in the Western South Atlantic. *J. Geophys. Res.*, **110**, C11016, doi: 10.1029/2004JC002736.
- Saraceno, M., Provost, C., Piola, A., Bava, J. and Gagliardini, A. (2004) Brazil Malvinas Frontal System as seen from 9 years of advanced very high resolution radiometer data. *J. Geophys. Res.*, **109**, C05027, doi: 10.1029/2003JC002127.
- Sarno, S., Kooistra, W., Medlin, L., Percopo, I. and Zingone, A. (2005) Diversity in the genus *Skeletonema* (Bacillariophyceae). II. An assessment of the taxonomy of *S. costatum*-like species with the description of four new species. *J. Phycol.*, **41**, 151–176.
- Shannon, C. E. and Weaver, W. (1949) *The Mathematical Theory of Communication*. University of Illinois Press, Urbana.
- Simonsen, R. (1974) The diatom plankton of the Indian Ocean expedition of RV Meteor 1964–1965. *Meteor Forsch.-Ergeb.*, **D 19**, 1–66.
- Spindler, M., Hemleben, Ch, Bayer, U., Bé, A. W. H. and Anderson, O. R. (1979) Lunar periodicity of reproduction in the planktonic foraminifer *Hastigerina pelagica*. *Mar. Ecol. Prog. Ser.*, **1**, 61–64.
- Sullivan, C. W., Arrigo, K. R., McClain, C. R., Comiso, J. C. and Firestone, J. (1993) Distributions of phytoplankton blooms in the Southern Ocean. *Science*, **262**, 1832–1837.
- Takahashi, K. (1987) Seasonal fluxes of silicoflagellates and *Actiniscus* in the subarctic Pacific during 1982–1984. *J. Mar. Res.*, **45**, 397–425.
- Takahashi, K. and Blackwelder, P. L. (1992) The spatial distribution of silicoflagellates in the region of the Gulf Stream warm-core ring 82B: application to water mass tracer studies. *Deep-Sea Res.*, **39**, 327–346.
- Tarling, G., Ward, P., Shearer, M., Williams, J. and Symon, C. (1995) Distribution patterns of macrozooplankton assemblages in the southwest Atlantic. *Mar. Ecol. Prog. Ser.*, **120**, 29–40.
- Thompson, G. A. (2001) Distribución y abundancia del microzooplankton en el Atlántico Sudoccidental, con especial énfasis en la biogeografía de Tintinnina (Protozoa, Ciliata). Tesis de Doctorado. FCEyN, UBA.
- Thompson, G. A. (2004) Tintinnid diversity trends in the southwestern Atlantic Ocean (29–60°S). *Aquat Microb. Ecol.*, **35**, 93–103.
- Thompson, G. A., Alder, V. A. and Boltovskoy, D. (2001) Tintinnids (Ciliophora) and other microzooplankton (>30 µm) in Southwestern Atlantic shelf break waters. *Mar. Ecol.*, **22**, 343–355.
- Thompson, G. A., Alder, V. A., Boltovskoy, D. and Brandini, F. (1999) Abundance and biogeography of tintinnids (Ciliophora) and associated microzooplankton in the Southwestern Atlantic Ocean. *J. Plankton Res.*, **21**, 1265–1298.
- Tsuchiya, M., Talley, L. D. and McCartney, M. S. (1994) Water-mass distributions in the western South Atlantic. A section from South Georgia Island (54°S) northward across the equator. *J. Mar. Res.*, **52**, 55–81.
- Utermöhl, H. (1958) Zur Vervollkommnung der quantitativen Phytoplankton-Methodik. *Mitt. Int. Ver. Limnol.*, **9**, 1–38.
- Venrick, E. L. (1971) Recurrent groups of diatom species in the North Pacific. *Ecology*, **52**, 614–625.
- Willson, H. R. and Rees, N. W. (2000) Classification of mesoscale features in the Brazil-Falkland Current confluence zone. *Prog. Oceanogr.*, **45**, 415–426.

## APPENDIX 1

Species list and relative abundance data. First set of square brackets: mean relative contribution of the species to the diatom assemblage at the stations surveyed; second

set of square brackets: proportion of stations where the species was recorded. Code letters give general ecological preferences of the species as derived from the literature sources indicated. C, cold; Co, cosmopolitan; N, neritic; O, oceanic; T, temperate; W, warm; SA, subantarctic; F, freshwater. Literature sources: A, Hasle and Syvertsen (1996); B, Lange (1985); C, Simonsen (1974); D, Hustedt, (1958); E, Hendey (1964); F, Medlin and Priddle (1990); G, Priddle and Fryxell (1985); H, Rines (1999); I, Romero *et al.* (2005); J, Romero and Hensen (2002).

- Achnantes* sp. [0.011] [25]  
*Achnanthes brevipes* Agardh [0.069] [30] N Co (B)  
*Actinocyclus actinochilus* (Ehrenberg) Simonsen [0.043] [20] O C (A,B)  
*Actinocyclus cholnoky* (Grunow) Van Landingham [0.006] [5] C (6)  
*Actinocyclus curvatulus* Janisch [0.897] [95] N Co (A,B)  
*Actinocyclus exiguus* Fryxell and Semina [0.022] [10] C (A)  
*Actinocyclus octonarius* Ehrenberg [0.360] [45] N Co (A,B)  
*Actinocyclus spiritus* Watkins [0.059] [5] C (A)  
*Actinopterychus senarius* (Ehrenberg) Ehrenberg [0.007] [25] N Co (B)  
*Actinopterychus vulgaris* Schumann [0.007] [5] N T (B)  
*Alveus marinus* (Grunow) Kaczmarek and Fryxell [0.069] [15] W (A)  
*Amphora* sp. [0.015] [5] F  
*Asterolampa marylandica* Ehrenberg [0.001] [5] O W (A,C)  
*Asteromphalus arachne* Brébisson [0.005] [10] O W (C)  
*Asteromphalus flabellatus* (Brébisson) Greville [0.001] [5] W (C)  
*Asteromphalus heptactis* (Brébisson) Ralfs [0.040] [5] Co (C)  
*Asteromphalus hookeri* Ehrenberg [0.344] [45] N/O C (B,D)  
*Asteromphalus hyalinus* Karsten [0.245] [50] N/O C (B,D)  
*Asteromphalus parvulus* Karsten [0.208] [45] N/O C (B,D)  
*Asteromphalus sarcophagus* Wallich [0.035] [15] W (C)  
*Aulacoseira granulata* Thwaites [0.033] [10] F  
*Azpeitia africana* (Janisch ex Schmidt) Fryxell and Watkins [0.014] [10] W (A)  
*Azpeitia neocrenulata* (Van Landingham) Fryxell and Watkins [0.005] [5] W (A)  
*Azpeitia tabularis* (Grunow) Fryxell and Sims [1.620] [100] O C, SA (A,B)  
*Bacteriastrum comosum* Pavillard [0.009] [10] W (A)  
*Bacteriastrum delicatulum* Cleve [0.442] [30] W/T (A)  
*Bacteriastrum hyalinum* Lauder [0.466] [40] N/O W/T (A,B)  
*Banquisia belgicae* (Van Heurck) Paddock [0.079] [15] C (A)  
*Cerataulina pelagica* (Cleve) Hendey [0.074] [15] N Co (A,B)  
*Chaetoceros affinis* Lauder [0.129] [40] N Co (A,B)  
*Chaetoceros atlanticus* Cleve [0.732] [50] O Co (A,B)  
*Chaetoceros bulbosum* (Ehrenberg) Heiden [0.024] [15] C (G)  
*Chaetoceros castracanei* Karsten [0.105] [15] C (A)  
*Chaetoceros coarctatus* Lauder [0.010] [5] W (A)  
*Chaetoceros contortus* Schütt [1.680] [40] Co (H)  
*Chaetoceros concavicornis* Manguin [0.037] [45] O Co (A,B)  
*Chaetoceros constrictus* Gran [0.263] [25] N T (A,B)  
*Chaetoceros convolutus* Castracane [2.568] [15] O T (A,B)  
*Chaetoceros costatus* Pavillard [0.002] [5] W/T (A)  
*Chaetoceros criophilus* Castracane [0.463] [30] C (A)  
*Chaetoceros curvisetus* Cleve [0.179] [20] N W/ T (A,B)  
*Chaetoceros danicus* Cleve [0.201] [5] N T Co? (A,B)  
*Chaetoceros debilis* Cleve [0.204] [10] N Co (A)  
*Chaetoceros decipiens* Cleve [1.597] [60] N/O Co (A,C)  
*Chaetoceros dictyota* Ehrenberg [0.121] [25] C (A)  
*Chaetoceros didymus* Ehrenberg [0.366] [35] N W/T (A,B)  
*Chaetoceros diversus* Cleve [0.030] [20] W (A)  
*Chaetoceros flexuosus* Mangin [0.079] [30] C (A)  
*Chaetoceros lacinosus* Schütt [0.537] [30] T (A)  
*Chaetoceros lorenzianus* Grunow [0.368] [35] N/O W (A,B)  
*Chaetoceros messanensis* Castracane [0.011] [20] W (A)  
*Chaetoceros neglectus* Karsten [1.259] [35] C (A)  
*Chaetoceros pendulus* Karsten [0.014] [5]  
*Chaetoceros peruvianus* Brightwell [0.045] [30] N/O W/T (A,B)  
*Chaetoceros rostratus* Lauder [4.907] [35] W (A)  
*Chaetoceros seiracanthus* Gran [0.006] [10] W/T (A)  
*Chaetoceros similis* Cleve [0.004] [10] ? (A)  
*Climacodium frauenfeldianum* Grunow [0.020] [5] W (A)  
*Cocconeis* spp. [0.013] [10]  
*Corethron criophilum* Castracane [2.153] [85] O Co (A,B)  
*Coscinodiscus asteromphalus* Ehrenberg [0.001] [5] O Co (A,B)  
*Coscinodiscus centralis* Ehrenberg [0.058] [25] O Co (A,B)  
*Coscinodiscus granii* Gough [0.000] [5] N Co (A,B)  
*Coscinodiscus marginatus* Ehrenberg [0.065] [40] N Co (A,B)  
*Coscinodiscus perforatus* Ehrenberg [0.001] [5] N/O Co (B)  
*Coscinodiscus oculus-iridis* Ehrenberg [0.001] [5] O Co (B)  
*Coscinodiscus radiatus* Ehrenberg [0.115] [50] N/O Co (A,B)  
*Cyclotella litoralis* Lange and Syvertsen [0.145] [40] N T (A)  
*Cyclotella stelligera* Cleve and Grunow [0.050] [10]  
*Cylindrotheca closterium* (Ehrenberg) Lewin and Reinmann [0.100] [20] Co (A)  
*Cymbella* sp. [0.000] [5] (F)  
*Dactyliosolen antarcticus* Castracane [0.083] [35] O C (A,B)  
*Detonula pumila* (Castracane) Gran [0.310] [30] N Co (A,B)  
*Diploneis* sp. [0.000] [5]  
*Ditylum brightwellii* (West) Grunow [0.008] [25] N Co (A,B)  
*Endictya* sp. [0.026] [15]  
*Eucampia antarctica* (Castracane) Manguin [1.335] [40] C (A)  
*Eucampia comuta* (Cleve) Grunow [0.037] [35] O W (A,B)  
*Fragilaria striatula* Lyngbye [0.003] [5] N (E)  
*Fragilariopsis curta* (Van Heurck) Hustedt [0.243] [45] C (A)  
*Fragilariopsis doliolus* (Wallich) Medlin and Sims [6.284] [55] O W (A,C)

- Fragilariopsis kerguelensis* (O'Meara) Hustedt [20.178] [85] O C (A,B)
- Fragilariopsis rhombica* (O'Meara) Hustedt [0.700] [60] C (A)
- Fragilariopsis ritscheri* Hustedt [1.353] [50] C (A)
- Gomphonema* sp. [0.011] [10]
- Gossleriella tropica* Schütt [0.020] [5] W (A)
- Grammatophora* sp. [0.004] [5]
- Guinardia flaccida* (Castracane) Peragallo [0.001] [10] N Co (A,B)
- Hantzschia* sp. [0.030] [10] F
- Haslea trompüi* (Cleve) Simonsen [0.000] [5] C (A)
- Hemidiscus cuneiformis* Wallich [0.010] [5] W (A)
- Hemiaulus hauckii* Grunow [0.149] [25] N W/T (A,C)
- Hemiaulus membranaceus* Cleve [0.145] [15] W (A)
- Hemiaulus sinensis* Greville [0.040] [10] N W/T (A,B)
- Lauderia annulata* Cleve [0.004] [15] W/T (A)
- Leptocylindrus danicus* Cleve [2.352] [25] Co (A)
- Leptocylindrus mediterraneus* (Peragallo) Hasle [0.151] [30] Co (A)
- Licmophora* sp. [0.001] [5]
- Lioloma pacificum* (Cupp) Hasle comb. nov. [0.292] [35] W/T (A)
- Mastogloia capitata* Cleve [0.010] [10] W (C)
- Mastogloia rostrata* (Wallich) Hustedt [0.075] [5] O W (C)
- Membraneis challengeri* (Grunow) Paddock [0.099] [40] C (F)
- Membraneis imposter* Paddock [0.010] [5] C (F)
- Navicula directa* (Smih) Ralfs [0.210] [20]
- Navicula distans* (Smih) Ralfs [0.078] [15]
- Neodelphineis indica* (Taylor) Hasle [0.041] [10] W/T (A)
- Neodenticula* sp. [0.003] [10]
- Nitzschia bicapitata* Group [2.022] [85] O W/T (A,B)
- Nitzschia braarudii* Hasle [0.430] [35] W/T (A)
- Nitzschia kolaczekii* Grunow [0.100] [30] O W (C)
- Nitzschia sicula* (Castracane) Hustedt [3.066] [75] N/O W (A,B)
- Odontella mobiliensis* (Bailey) Grunow [0.000] [10] N Co (A,B)
- Paralia sulcata* (Ehrenberg) Cleve [0.009] [5] N Co (A,B)
- Pinnularia* sp. [0.008] [10] F
- Plagiotropis gaussi* (Heiden) Paddock [0.020] [10] C (F)
- Planktoniella sol* (Wallich) Schütt [0.126] [45] W (A)
- Pleurosigma directum* Grunow [0.200] [50] Co (A)
- Pleurosigma normanii* Ralfs [0.000] [5] N Co (A,B)
- Porosira pseudodenticulata* (Hustedt) Josué [0.042] [20] C (A)
- Proboscia alata* (Brightwell) Sundström [0.495] [70]
- Proboscia inermis* (Castracane) Jordan and Ligowski [0.103] [40] C (F)
- Pseudo-nitzschia delicatissima* (Cleve) Heiden [0.293] [20] Co (A)
- Pseudo-nitzschia fraudulenta* (Cleve) Hasle [1.026] [45] N/O Co (A,B)
- Pseudo-nitzschia granii* (Hasle) Hasle [0.018] [5] C/T (A)
- Pseudo-nitzschia heimii* Manguin [1.704] [40] Co (A)
- Pseudo-nitzschia inflatula* var. *capitata* Simonsen [0.012] [15]
- Pseudo-nitzschia lineola* (Cleve) Hasle [2.319] [40] Co (A)
- Pseudo-nitzschia multiseriata* (Hasle) Hasle [1.903] [35] Co (A)
- Pseudo-nitzschia pseudodelicatissima* (Hasle) Hasle [1.934] [70] Co (A)
- Pseudo-nitzschia pungens* (Grunow ex Cleve) Hasle [0.097] [10] N Co (A,B)
- Pseudo-nitzschia seriata* (Cleve) Peragallo [0.131] [15] T (A)
- Pseudo-nitzschia subfraudulenta* (Hasle) Hasle [0.558] [35] W/T (A)
- Pseudo-nitzschia turgidula* (Hustedt) Hasle [0.250] [10] Co (A)
- Pseudo-nitzschia turgiduloides* (Hasle) Hasle [0.403] [15] C (F)
- Pseudosolenia calcar-avis* (Schulze) Sundström [0.068] [35] N/O W/T (A,B)
- Rhizosolenia antennata* f. *antennata* (Ehrenberg) Brown [0.016] [10] C (A)
- Rhizosolenia antennata* f. *semispina* Sundström [0.226] [25] C (A)
- Rhizosolenia chunii* Karsten [0.281] [40] C (F)
- Rhizosolenia clevei* var. *communis* Sundström [0.107] [20] W (A)
- Rhizosolenia crassa* Schimper [0.043] [35] C (A)
- Rhizosolenia curvata* Zacharias [0.025] [20] C (A)
- Rhizosolenia antarctica* Karsten [0.423] [25] C (D)
- Rhizosolenia hebetata* f. *semispina* (Hensen) Gran ? [0.020] [10] C (F)
- Rhizosolenia imbricata* Brightwell [0.093] [40] N Co (A,B)
- Rhizosolenia polydactyla* (Castracane) f. *polydactyla* [0.116] [30] C (A)
- Rhizosolenia pungens* Cleve-Euler [0.185] [40] N (A)
- Rhizosolenia setigera* Brightwell [2.651] [65] N Co (A,B)
- Rhizosolenia simplex* Karsten [0.157] [35] C (A)
- Rhizosolenia styliiformis* Brightwell [0.925] [45] N/O Co (B)
- Roperia tessellata* (Roper) Grunow [0.338] [50] N/O W/Co ? (A,B)
- Skeletonema costatum* s.l. (Greville) Cleve [0.560] [35] N Co (A,B) Note: recent molecular genetic studies have shown that *Skeletonema costatum*-like species comprise eight distinct entities (Sarno *et al.*, 2005)
- Stellarima microtrias* (Ehrenberg) Hasle and Sims [0.009] [10] C (A)
- Stephanodiscus* sp. [0.000] [5] F
- Stephanopyxis turris* (Arnott) Ralfs [0.014] [5] N W/T (A,B)
- Thalassionema bacillare* Kolbe [0.111] [20] W (A)
- Thalassionema frauenfeldii* (Grunow) Hallegraeff [0.044] [35] W/T (A)
- Thalassionema nitzschioides* var. *parva* (Heiden) Moreno-Ruiz [0.090] [20] W (I)

- Thalassionema nitzschioides* (Grunow) Grunow ex Hustedt [8.133] [90] N Co (A,B)  
*Thalassiosira allenii* Takano [0.002] [5] W/T (A)  
*Thalassiosira angulata* (Gregory) Hasle [0.038] [10] Co (B)  
*Thalassiosira anguste-lineata* (Schmidt) Fryxell and Hasle [0.087] [60] N Co (A,B)  
*Thalassiosira antarctica* Comber (vegetative cell) [0.404] [35] N/O C (A,B)  
*Thalassiosira dichotomica* (Kozlova) Fryxell and Hasle [0.045] [10] C (F)  
*Thalassiosira decipiens* (Grunow) Jørgensen [2.884] [35] N (A)  
*Thalassiosira delicatula* Ostenfeld in Borgert [0.003] [5] N/O Co (A,B)  
*Thalassiosira eccentrica* (Ehrenberg) Cleve [0.035] [35] N Co (A,B)  
*Thalassiosira frenguelli* Kozlova [0.211] [50] C (F)  
*Thalassiosira frenguelliopsis* Fryxell and Johansen [0.047] [30] C (F)  
*Thalassiosira gracilis* (Karsten) Hustedt [1.244] [45] N/O C (A,B)  
*Thalassiosira gravida* Cleve [0.061] [25] N/O C (A,B)  
*Thalassiosira lentiginosa* (Janisch) Fryxell [1.793] [45] O C (A,B)  
*Thalassiosira leptopus* (Grunow) Hasle and Fryxell [0.040] [5] W (A)  
*Thalassiosira lineata* Jousé [0.227] [50] O W (A,B)  
*Thalassiosira oestrupii* var. *oestrupii* (Ostenfeld) Hasle [0.312] [55] O W/T (A,B)  
*Thalassiosira oestrupii* var. *venrickae* Fryxell and Hasle [0.446] [65] W/T (A)  
*Thalassiosira oliverana* (O'Meara) Mak [0.041] [25] C (G)  
*Thalassiosira pacifica* Grand and Angst [0.006] [20] Co (A)  
*Thalassiosira perpusilla* Kozlova [0.079] [25] C (A)  
*Thalassiosira poroseriata* (Ramsfjell) Hasle [0.059] [15] Co (A)  
*Thalassiosira punctifera* (Grunow) Fryxell [0.051] [15] N/O (C)  
*Thalassiosira punctigera* (Castracane) Hasle [0.094] [15] N/O W/T (A,B)  
*Thalassiosira ritscheri* (Hustedt) Hasle [0.065] [45] C (A)  
*Thalassiosira sacketti* Fryxell [0.010] [5] W (J)  
*Thalassiosira scotia* Fryxell and Hoban [0.112] [40] C (F)  
*Thalassiosira symmetrica* Fryxell and Hasle [0.010] [5] Co (C)  
*Thalassiosira simonsenii* Hasle and Fryxell [0.001] [5] N/O W/T (A,B)  
*Thalassiosira subtilis* (Ostenfeld) Gran [0.010] [5] O W/T (A,B)  
*Thalassiosira trifulta* Fryxell [0.137] [60] N/O C (A,B)  
*Thalassiosira tumida* (Janisch) Hasle [0.186] [45] N/O C (A,B)  
*Thalassiothrix antarctica* Schimper ex Karsten [0.909] [55] N/O C (A)  
*Trichotoxon reinboldii* (Van Heurck) Reid and Round [0.303] [35] C (A)