

Partial characterizations of clique-perfect graphs I: Subclasses of claw-free graphs[☆]

Flavia Bonomo^{a,1}, Maria Chudnovsky^{b,c,2}, Guillermo Durán^{d,3}

^aDepartamento de Matemática, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina

^bDepartment of IEOR, Columbia University, New York, NY, USA

^cDepartment of Mathematics, Columbia University, New York, NY, USA

^dDepartamento de Ingeniería Industrial, Facultad de Ciencias Físicas y Matemáticas, Universidad de Chile, Santiago, Chile

Received 27 June 2005; received in revised form 12 June 2006; accepted 20 May 2007

Available online 4 September 2007

Abstract

A *clique-transversal* of a graph G is a subset of vertices that meets all the cliques of G . A *clique-independent set* is a collection of pairwise vertex-disjoint cliques. The *clique-transversal number* and *clique-independence number* of G are the sizes of a minimum clique-transversal and a maximum clique-independent set of G , respectively. A graph G is *clique-perfect* if these two numbers are equal for every induced subgraph of G . The list of minimal forbidden induced subgraphs for the class of clique-perfect graphs is not known. In this paper, we present a partial result in this direction; that is, we characterize clique-perfect graphs by a restricted list of forbidden induced subgraphs when the graph belongs to two different subclasses of claw-free graphs.

© 2007 Elsevier B.V. All rights reserved.

Keywords: Claw-free graphs; Clique-perfect graphs; Hereditary clique-Helly graphs; Line graphs; Perfect graphs

1. Introduction

Let G be a graph, with vertex set $V(G)$ and edge set $E(G)$. Denote by \overline{G} , the complement of G . Given two graphs G and G' we say that G' is *smaller* than G if $|V(G')| < |V(G)|$, and that G *contains* G' if G' is isomorphic to an induced subgraph of G . When we need to refer to the non-induced subgraph containment relation, we will say so explicitly. A *claw* is the graph isomorphic to $K_{1,3}$. A graph is *claw-free* if it does not contain a claw. The *line graph* $L(G)$ of G is the intersection graph of the edges of G . A graph F is a *line graph* if there exists a graph H such that $L(H) = F$. Clearly, line graphs are a subclass of claw-free graphs.

[☆] An extended abstract of this paper was presented at GRACO 2005 (second Brazilian Symposium on Graphs, Algorithms, and Combinatorics) and appeared, under a different title, in Electronic Notes in Discrete Mathematics 19 (2005) 95–101.

E-mail addresses: fbonomo@dm.uba.ar (F. Bonomo), mchudnov@columbia.edu (M. Chudnovsky), gduran@dii.uchile.cl (G. Durán).

¹ Partially supported by UBACyT Grant X184, PICT ANPCyT Grant 11-09112, Argentina, and CNPq under PROSUL Project Proc. 490333/2004-4, Brazil.

² This research was conducted during the period the author served as a Clay Mathematics Institute Research Fellow. The research was conducted in part at Princeton University.

³ Partially supported by FONDECyT Grant 1050747, Millennium Science Nucleus “Complex Engineering Systems”, Chile, and CNPq under PROSUL Project Proc. 490333/2004-4, Brazil.

The neighborhood of a vertex v is the set $N(v)$ consisting of all the vertices which are adjacent to v . The closed neighborhood of v is $N[v] = N(v) \cup \{v\}$. A vertex v of G is *universal* if $N[v] = V(G)$. Two vertices v and w are *twins* if $N[v] = N[w]$; and u *dominates* v if $N(v) \subseteq N[u]$.

A *complete set* or just a *complete* of G is a subset of vertices pairwise adjacent. (In particular, an empty set is a complete set.) We denote by K_n the graph induced by a complete set of size n . A *clique* is a complete set not properly contained in any other. We may also use the term clique to refer to the corresponding complete subgraph. Let X and Y be two sets of vertices of G . We say that X is *complete to* Y if every vertex in X is adjacent to every vertex in Y , and that X is *anticomplete to* Y if no vertex of X is adjacent to a vertex of Y . A *stable set* in a graph G is a subset of pairwise non-adjacent vertices of G . The *stability number* $\alpha(G)$ is the cardinality of a maximum stable set of G .

A complete of three vertices is called a *triangle*, and a stable set of three vertices is called a *triad*. Let A be a set of vertices of G , and v a vertex of G not in A . Then v is *A-complete* if it is adjacent to every vertex in A , and *A-anticomplete* if it has no neighbor in A .

A vertex is called *simplicial* if its neighbors induce a complete, and *singular* if its non-neighbors induce a complete. Equivalently, a vertex is singular if it is in no stable set of size three. The *core* of G is the subgraph induced on G by the set of non-singular vertices.

Let G be a graph and X be a subset of vertices of G . Denote by $G|X$ the subgraph of G induced by X and by $G \setminus X$ the subgraph of G induced by $V(G) \setminus X$. X is *connected*, if there is no partition of X into two non-empty sets Y and Z , such that no edge has one end in Y and the other one in Z . In this case the graph $G|X$ is also connected. X is *anticonnected* if it is connected in \bar{G} . In this case the graph $G|X$ is also anticonnected.

The set X is a *cutset* if $G \setminus X$ has more connected components than G . Let G be a connected graph, X a cutset of G , and M_1, M_2 a partition of $V(G) \setminus X$ such that M_1, M_2 are non-empty and M_1 is anticomplete to M_2 in G . In this case we say that $G = M_1 + M_2 + X$, and $M_i + X$ denotes $G|(M_i \cup X)$, for $i = 1, 2$. When $X = \{v\}$, we simplify the notation to $M_1 + M_2 + v$ and $M_i + v$, respectively.

Let X be a cutset of G . If $X = \{v\}$ we say that v is a *cutpoint*. If X is complete, it is called a *clique cutset*. A clique cutset X is *internal* if $G = M_1 + M_2 + X$ and each M_i contains at least two vertices that are not twins.

Let G be a graph and H a subgraph of G (not necessarily induced). The graph H is a *clique subgraph* of G if every clique of H is a clique of G .

A *clique cover* of a graph G is a subset of cliques covering all the vertices of G . The *clique-covering number* $k(G)$ is the cardinality of a minimum clique cover of G . The *chromatic number* of a graph G is the smallest number of colors that can be assigned to the vertices of G in such a way that no two adjacent vertices receive the same color, and is denoted by $\chi(G)$. An obvious lower bound is the maximum cardinality of the cliques of G , the *clique number* of G , denoted by $\omega(G)$.

A graph G is *perfect* if $\chi(H) = \omega(H)$ for every induced subgraph H of G . Perfect graphs are interesting from the algorithmic point of view, see [16]. While determining the clique-covering number, the independence number, the chromatic number, and the clique number of a graph are NP-complete problems, they are solvable in polynomial time for perfect graphs [17].

The *clique graph* $K(G)$ of G is the intersection graph of the cliques of G . A graph G is *K-perfect* if $K(G)$ is perfect.

A graph is *bipartite* if its vertex set can be partitioned into two stable sets. A graph is *split* if its vertex set can be partitioned into a stable set and a complete. Bipartite and split graphs are perfect.

A *hole* is a chordless cycle of length at least 4. An *antihole* is the complement of a hole. A hole or antihole is said to be *odd* if it consists of an odd number of vertices. A hole of length n is denoted by C_n .

A graph is *chordal* if it does not contain a hole. Chordal graphs can be recognized in polynomial time [25].

An *r-sun*, $r \geq 3$, is a chordal graph of $2r$ vertices whose vertex set can be partitioned into two sets: $W = \{w_1, \dots, w_r\}$ and $U = \{u_1, \dots, u_r\}$, such that W is a stable set and, for each i and j , w_j is adjacent to u_i if and only if $i = j$ or $i \equiv j + 1 \pmod{r}$. Please note that, since an *r-sun* is a chordal graph, it follows that U induces a cycle with no holes. An *r-sun* is said to be *odd* if r is odd.

A graph is *balanced* if its vertex-clique incidence matrix is balanced. A 0–1 matrix is balanced if it does not contain the incidence matrix of an odd cycle as a submatrix.

A family of sets S is said to satisfy the *Helly property* if every subfamily of it, consisting of pairwise intersecting sets, has a common element. A graph is *clique-Helly (CH)* if its cliques satisfy the Helly property, and it is *hereditary clique-Helly (HCH)* if H is CH for every induced subgraph H of G .

A *clique-transversal* of a graph G is a subset of vertices that meets all the cliques of G . A *clique-independent set* is a collection of pairwise vertex-disjoint cliques. The *clique-transversal number* and *clique-independence number* of G , denoted by $\tau_C(G)$ and $\alpha_C(G)$, are the sizes of a minimum clique-transversal and a maximum clique-independent set of G , respectively. It is easy to see that $\tau_C(G) \geq \alpha_C(G)$ for any graph G . A graph G is *clique-perfect* if $\tau_C(H) = \alpha_C(H)$ for every induced subgraph H of G . Clique-perfect graphs have been implicitly studied in [1,3,6,4,7,14,18,19]. The terminology “clique-perfect” has been introduced in [18]. There are two main open problems concerning this class of graphs:

- find all minimal forbidden induced subgraphs for the class of clique-perfect graphs, and
- is there a polynomial time recognition algorithm for this class of graphs?

In this paper, we present some results related to these problems. We characterize clique-perfect graphs by forbidden subgraphs when the graph belongs to a certain class. Both classes studied are subclasses of claw-free graphs: line graphs and HCH claw-free graphs. As corollaries of these partial characterizations, we can immediately deduce polynomial time algorithms to recognize clique-perfect graphs in these classes of graphs.

2. Preliminaries

It has been proved recently that perfect graphs can be characterized by two families of minimal forbidden induced subgraphs [9] and recognized in polynomial time [8].

Theorem 1 (*Strong perfect graph theorem, Chudnovsky et al. [9]*). *Let G be a graph. Then the following are equivalent:*

- (i) *no induced subgraph of G is an odd hole or an odd antihole.*
- (ii) *G is perfect.*

On the other hand, the problem of recognition of clique-perfect chordal graphs can be reduced to the recognition of balanced graphs, which is solvable in polynomial time [5,13].

Theorem 2 (*Lehel and Tuza [19]*). *Let G be a chordal graph. Then the following are equivalent:*

- (i) *G does not contain odd suns.*
- (ii) *G is balanced.*
- (iii) *G is clique-perfect.*

Next we define the family of the so-called “generalized suns” [4]. Let G be a graph and C be a cycle of G not necessarily induced. An edge of C is *non-proper* (or *improper*) if it forms a triangle with some vertex of C . An *r-generalized sun*, $r \geq 3$, is a graph G whose vertex set can be partitioned into two sets: a cycle C of r vertices, with all its non-proper edges $\{e_j\}_{j \in J}$ (J is permitted to be an empty set) and a stable set $U = \{u_j\}_{j \in J}$, such that, for each $j \in J$, u_j is adjacent only to the endpoints of e_j . An *r-generalized sun* is said to be *odd* if r is odd. Clearly, an odd hole is an odd generalized sun, with the set of non-proper edges J being empty. Odd suns are also odd generalized suns, since every edge of the cycle in an r -sun is a non-proper edge.

Theorem 3 (*Bonomo et al. [4]*). *Odd generalized suns and antiholes of length $t = 1, 2 \pmod 3$ ($t \geq 5$) are not clique-perfect.*

Unfortunately, odd generalized suns are not necessarily minimal (with respect to taking induced subgraphs) and besides there are other minimal non-clique-perfect graphs, for example, the following family of graphs. Define the graph S_k , $k \geq 2$, as follows: $V(S_k) = \{v_1, \dots, v_{2k}, v, v', w, w'\}$ where v_1, \dots, v_{2k} induce a path; v is adjacent to v', v_1, v_2 , and v_{2k} ; v' is adjacent to v, v_1, v_{2k-1} , and v_{2k} ; w is adjacent only to v_1 and v_2 ; and w' is adjacent only to v_{2k-1} and v_{2k} (Fig. 1).




Fig. 1. The graph S_k .




Fig. 2. Forbidden induced subgraphs for hereditary clique-Helly graphs: (left to right) 3-sun (or 0-pyramid), 1-pyramid, 2-pyramid, and 3-pyramid.

Every clique of S_k contains at least two of the vertices v_1, \dots, v_{2k}, v , so $\alpha_C(S_k) \leq k$. On the other hand, consider the following family of cliques of S_k : $\{v_{2k-1}, v_{2k}, w'\}$, $\{v_{2k}, v, v'\}$, $\{v, v_1, v'\}$, $\{v_1, v_2, w\}$, and either $\{v_2, v_{2k-1}\}$, if $k = 2$, or $\{v_2, v_3, \dots, v_{2k-2}, v_{2k-1}\}$, if $k > 2$. No vertex of S_k belongs to more than two of these $2k + 1$ cliques, so $\tau_C(S_k) \geq k + 1$.

At this time we do not know whether the list of all such forbidden graphs has a nice description. However, if we restrict our attention to certain classes of graphs (that can be described by forbidding certain induced subgraphs), we can describe all the minimal forbidden induced subgraphs.

HCH graphs are of particular interest because in this case it follows from [4] that if $K(H)$ is perfect for every induced subgraph H of G , then G is clique-perfect (the converse is not necessarily true). On the other hand, the class of HCH graphs can be characterized by forbidden induced subgraphs.

Theorem 4 (Prisner [23]). *A graph G is HCH if and only if it does not contain the graphs of Fig. 2 as induced subgraphs.*

One of our main results in this paper is a characterization of clique-perfect HCH claw-free graphs by forbidden induced subgraphs. To prove this characterization we use a recent structure theorem for claw-free graphs [11]. In order to state that theorem we need to introduce some definitions.

A graph G is *prismatic* if for every triangle T of G , every vertex of G not in T has a unique neighbor in T . A graph G is *antiprismatic* if its complement graph \bar{G} is prismatic.

Construct a graph G as follows. Take a circle C , and let $V(G)$ be a finite set of points of C . Take a set of intervals from C (an *interval* means a proper subset of C homeomorphic to $[0, 1]$) such that there are not three intervals covering C ; and say that $u, v \in V(G)$ are adjacent in G if the set of points $\{u, v\}$ of C is a subset of one of the intervals. Such a graph is called *circular interval graph*. When the set of intervals does not cover C , the graph is called *linear interval graph* (Fig. 3).

Let G be a graph and A, B be disjoint subsets of $V(G)$. The pair (A, B) is called a *homogeneous pair* in G if for every vertex $v \in V(G) \setminus (A \cup B)$, v is either A -complete or A -anticomplete and either B -complete or B -anticomplete. If, in addition, B is empty, then A is called a *homogeneous set*. Let (A, B) be a homogeneous pair, such that A, B are both completes, and A is neither complete nor anticomplete to B . In these circumstances the pair (A, B) is called a *W-join*. Note that there is no requirement that $A \cup B \neq V(G)$. The pair (A, B) is *non-dominating* if some vertex of $G \setminus (A \cup B)$ has no neighbor in $A \cup B$, and it is *coherent* if the set of all $(A \cup B)$ -complete vertices in $V(G) \setminus (A \cup B)$ is a complete.

Next, suppose that V_1, V_2 is a partition of $V(G)$ such that V_1, V_2 are non-empty and there are no edges between V_1 and V_2 . The pair (V_1, V_2) is called a *0-join* in G . Thus G admits a 0-join if and only if it is not connected.




Fig. 3. Example of a circular interval graph and its circular interval representation.

Next, suppose that V_1, V_2 is a partition of $V(G)$, and for $i = 1, 2$ there is a subset $A_i \subseteq V_i$ such that:

- for $i = 1, 2$, A_i is a complete, and $A_i, V_i \setminus A_i$ are both non-empty,
- A_1 is complete to A_2 ,
- every edge between V_1 and V_2 is between A_1 and A_2 .

In these circumstances, the pair (V_1, V_2) is a 1-join.

Now, suppose that V_0, V_1, V_2 are disjoint subsets with union $V(G)$, and for $i = 1, 2$ there are subsets A_i, B_i of V_i satisfying the following:

- for $i = 1, 2$, A_i, B_i are completes, $A_i \cap B_i = \emptyset$, and A_i, B_i , and $V_i \setminus (A_i \cup B_i)$ are all non-empty,
- A_1 is complete to A_2 , and B_1 is complete to B_2 , and there are no other edges between V_1 and V_2 ,
- V_0 is a complete, and, for $i = 1, 2$, V_0 is complete to $A_i \cup B_i$ and anticomplete to $V_i \setminus (A_i \cup B_i)$.

The triple (V_0, V_1, V_2) is called a *generalized 2-join*, and, if $V_0 = \emptyset$, the pair (V_1, V_2) is called a *2-join*. This is closely related to, but not the same as, what has been called a 2-join in other papers like [8].

The last decomposition is the following. Let (V_1, V_2) be a partition of $V(G)$, such that for $i = 1, 2$ there are completes $A_i, B_i, C_i \subseteq V_i$ with the following properties:

- for $i = 1, 2$ the sets A_i, B_i, C_i are pairwise disjoint and have union V_i ,
- V_1 is complete to V_2 except that there are no edges between A_1 and A_2 , between B_1 and B_2 , and between C_1 and C_2 ,
- V_1, V_2 are both non-empty.

In these circumstances it is said that G is a *hex-join* of $G|V_1$ and $G|V_2$. Note that if G is expressible as a hex-join as above, then the sets $A_1 \cup B_2, B_1 \cup C_2$, and $C_1 \cup A_2$ are three completes with union $V(G)$, and consequently no graph G with $\alpha(G) > 3$ is expressible as a hex-join.

Now, define classes $\mathcal{S}_0, \dots, \mathcal{S}_6$ as follows:

- \mathcal{S}_0 is the class of all line graphs.
- The *icosahedron* is the unique planar graph with 12 vertices all of degree five. For $0 \leq k \leq 3$, $icosa(-k)$ denotes the graph obtained from the icosahedron by deleting k pairwise adjacent vertices. A graph $G \in \mathcal{S}_1$ if G is isomorphic to $icosa(0)$, $icosa(-1)$, or $icosa(-2)$. As it can be seen in Fig. 4, all of them contain odd holes.
- Let H_1 be the graph with vertex set $\{v_1, \dots, v_{13}\}$, with adjacency as follows: $v_1 v_2 \dots v_6 v_1$ is a hole of length 6; v_7 is adjacent to v_1, v_2 ; v_8 is adjacent to v_4, v_5 and possibly to v_7 ; v_9 is adjacent to v_6, v_1, v_2, v_3 ; v_{10} is adjacent to v_3, v_4, v_5, v_6, v_9 ; v_{11} is adjacent to $v_3, v_4, v_6, v_1, v_9, v_{10}$; v_{12} is adjacent to $v_2, v_3, v_5, v_6, v_9, v_{10}$; and v_{13} is adjacent to $v_1, v_2, v_4, v_5, v_7, v_8$. A graph $G \in \mathcal{S}_2$ if G is isomorphic to $H_1 \setminus X$, where $X \subseteq \{v_{11}, v_{12}, v_{13}\}$. Please note that vertices $v_3 v_4 v_5 v_6 v_9 v_3$ induce a hole of length 5 in G (Fig. 5).
- \mathcal{S}_3 is the class of all circular interval graphs.
- Let H_2 be the graph with seven vertices h_0, \dots, h_6 , in which h_1, \dots, h_6 are pairwise adjacent and h_0 is adjacent to h_1 . Let H_3 be the graph obtained from the line graph $L(H_2)$ of H_2 by adding one new vertex, adjacent precisely to the members of $V(L(H_2)) = E(H_2)$ that are not incident with h_1 in H_2 . Then H_3 is claw-free. Let \mathcal{S}_4 be the class




Fig. 4. Graphs $icos(0)$, $icos(-1)$, and $icos(-2)$.




Fig. 5. Graph $H_1 \setminus \{v_{11}, v_{12}, v_{13}\}$. Every graph in \mathcal{S}_2 contains it as an induced subgraph.




Fig. 6. Graph H_4 , for $n = 2$.

of all graphs isomorphic to induced subgraphs of H_3 . Note that the vertices of H_3 corresponding to the members of $E(H_2)$ that are incident with h_1 in H_2 form a complete in H_3 . So every graph in \mathcal{S}_4 is either a line graph or it has a singular vertex.

- Let $n \geq 0$. Let $A = \{a_1, \dots, a_n\}$, $B = \{b_1, \dots, b_n\}$, $C = \{c_1, \dots, c_n\}$ be three completes, pairwise disjoint. For $1 \leq i, j \leq n$, let a_i, b_j be adjacent if and only if $i = j$, and let c_i be adjacent to a_j, b_j if and only if $i \neq j$. Let d_1, d_2, d_3, d_4, d_5 be five more vertices, where d_1 is $(A \cup B \cup C)$ -complete; d_2 is complete to $A \cup B \cup \{d_1\}$; d_3 is complete to $A \cup \{d_2\}$; d_4 is complete to $B \cup \{d_2, d_3\}$; d_5 is adjacent to d_3, d_4 ; and there are no more edges. Let the graph just constructed be H_4 . A graph $G \in \mathcal{S}_5$ if (for some n) G is isomorphic to $H_4 \setminus X$ for some $X \subseteq A \cup B \cup C$. Note that vertex d_1 is adjacent to all the vertices but the triangle formed by d_3, d_4 , and d_5 , so it is a singular vertex in G (Fig. 6).
- Let $n \geq 0$. Let $A = \{a_0, \dots, a_n\}$, $B = \{b_0, \dots, b_n\}$, $C = \{c_1, \dots, c_n\}$ be three completes, pairwise disjoint. For $0 \leq i, j \leq n$, let a_i, b_j be adjacent if and only if $i = j > 0$, and for $1 \leq i \leq n$ and $0 \leq j \leq n$ let c_i be adjacent to a_j, b_j if and only if $i \neq j \neq 0$. Let the graph just constructed be H_5 . A graph $G \in \mathcal{S}_6$ if (for some n) G is isomorphic to $H_5 \setminus X$ for some $X \subseteq (A \setminus \{a_0\}) \cup (B \setminus \{b_0\}) \cup C$, and then G is said to be 2-simplicial of antihat type (Fig. 7).




Fig. 7. Graph H_5 , for $n = 2$.




Fig. 8. Some graphs mentioned in the paper.

The structure theorem in [11] is the following:

Theorem 5 (Chudnovsky and Seymour [11]). *Let G be a claw-free graph. Then either $G \in \mathcal{S}_0 \cup \dots \cup \mathcal{S}_6$, or G admits twins, or a non-dominating W -join, or a coherent W -join, or a 0-join, or a 1-join, or a generalized 2-join, or a hex-join, or G is antiprismatic.*

In the proofs in this paper we will mention some special graphs, shown in Fig. 8, and we will use the following results on perfect graphs, cutsets, and clique graphs (some of the results below are immediate, and in these cases we do not give a proof or a reference; we state these in order to make it more convenient to refer to them in the future).

Lemma 6. *Let G be a graph and v be a simplicial vertex of G . Then G is perfect if and only if $G \setminus \{v\}$ is.*

Theorem 7 (Berge [2]). *Let G be a graph and X be a clique cutset of G such that $G = M_1 + M_2 + X$. Then the graph G is perfect if and only if the graphs $M_1 + X$ and $M_2 + X$ are.*

Theorem 8 (Tucker [27]). *Let G be a perfect graph and let $e = v_1 v_2$ be an edge of G . Assume that no vertex of G is a common neighbor of v_1 and v_2 . Then $G \setminus e$ is perfect.*

Let P be an induced path of a graph G . The length of P is the number of edges in P . The parity of P is the parity of its length. We say that P is even if its length is even, and odd otherwise.

Theorem 9. *Let G be a graph, and let $u, v \in V(G)$ be non-adjacent such that $\{u, v\}$ is a cutset of G , $G = M_1 + M_2 + \{u, v\}$. For $i = 1, 2$, let G_i be a graph obtained from $M_i + \{u, v\}$ by joining u and v by an even induced path. If G_1 and G_2 are perfect, then G is perfect.*

Proof. Suppose G_1 and G_2 are perfect, and G contains an odd hole or an odd antihole; denote it by A . Since no odd antihole of length at least 7 has a one- or two-vertex cutset, if A is an odd antihole of length at least 7, then A is contained either in G_1 or in G_2 , a contradiction. So A is an odd hole, and it is not contained in $M_i + \{u, v\}$ for $i = 1, 2$, thus $\{u, v\}$ is a cutset for A . Let A_1, A_2 be the two subpaths of A joining u and v . Then both A_1, A_2 have length at least 2, and one of them, say A_1 , is odd. But then, if A_1 is contained in $M_i + \{u, v\}$, the graph G_i contains an odd hole, a contradiction. \square

Theorem 10 (Chvátal and Sbihi [12]). *Let G be a graph and let U be a homogeneous set in G . Let G' be the graph obtained from G by deleting all but one vertex of U . Then G is perfect if and only if both G' and $G \setminus U$ are.*

Theorem 11. Let G be a graph, and let $u, v \in V(G)$ such that u dominates v . Then G is perfect if and only if both $G \setminus \{u\}$ and $G \setminus \{v\}$ are.

Proof. The “only if” part is clear, so it is enough to prove that if $G \setminus \{u\}$ and $G \setminus \{v\}$ are perfect, then so is G . Since neither odd holes nor odd antiholes contain a pair of vertices such that one of them dominates the other one, the result follows from Theorem 1. \square

Theorem 12 (Chudnovsky and Seymour [10]). Let G be a claw-free graph admitting an internal clique cutset. Then G is either a linear interval graph or G admits twins, or a 0-join, or a 1-join, or a coherent W -join.

Lemma 13. Let G be a graph and H a clique subgraph of G . Then $K(H)$ is an induced subgraph of $K(G)$.

Lemma 14. If G admits twins u, v , then $K(G) = K(G \setminus \{v\})$.

Lemma 15. If G is disconnected, then so is $K(G)$, and G is K -perfect if and only if each connected component is.

Theorem 16 (Maffray and Reed [22], Protti and Szwarcfiter [24]). Let G be a claw-free graph with no induced 3-fan, 4-wheel, or odd hole. Then $K(G)$ is bipartite.

Graphs whose line graph is perfect were characterized in [26,21].

Theorem 17 (Maffray [21], Trotter [26]). Let $G = L(H)$ be the line graph of a graph H . Then the following three conditions are equivalent:

- (i) G is a perfect graph.
- (ii) No subgraph of H is an odd cycle of length at least 5.
- (iii) Any connected subgraph H' of H satisfies at least one of the following properties:
 - H' is a bipartite graph;
 - H' is a complete of size four;
 - H' consists of exactly $p + 2$ vertices x_1, \dots, x_p, a, b , such that $\{x_1, \dots, x_p\}$ is a stable set, and $\{x_j, a, b\}$ is a triangle for each $j = 1, \dots, p$;
 - H' has a cutpoint.

3. Partial characterizations

We say that a graph is *interesting* if no induced subgraph of it is an odd generalized sun or an antihole of length greater than 5 and equal to $1, 2 \pmod{3}$. Since odd generalized suns and antiholes of length greater than 5 and equal to $1, 2 \pmod{3}$ are not clique-perfect, it follows that every clique-perfect graph is interesting. We prove that for some subclasses of claw-free graphs, this necessary condition is also sufficient.

Our two main results are the following.

Theorem 18. Let G be a line graph. Then G is clique-perfect if and only if no induced subgraph of G is an odd hole or a 3-sun.

Theorem 19. Let G be an HCH claw-free graph. Then G is clique-perfect if and only if no induced subgraph of G is an odd hole or an antihole of length 7.

We observe the following:

Proposition 20. Let S be an odd generalized r -sun, and assume that S is claw-free. Then either S is an odd hole or $r = 3$.

Proof. As in the definition of a generalized sun, let C be a cycle of S , and let $U = V(S) \setminus V(C)$ be a stable set, such that every vertex of U is complete to both ends of exactly one non-proper edge of C and has no other neighbor in $V(C)$.




Fig. 9. A clique-perfect graph that is not K -perfect.

We may assume that S is not an odd hole, and so C has at least one non-proper edge. Let c_1c_2 be a non-proper edge of C , let $c_3 \in V(C) \setminus \{c_1, c_2\}$ be such that $\{c_1, c_2, c_3\}$ is a triangle, and let u be the vertex of U adjacent to c_1 and c_2 . We may assume $r > 3$, and therefore, possibly with c_1 and c_2 switched, c_1 has a neighbor c'_2 in C , different from c_2 and c_3 . Since $\{c_1, u, c_3, c'_2\}$ does not induce a claw in S , it follows that c'_2 is adjacent to c_3 , and therefore $c_1c'_2$ is another non-proper edge of S . Let u' be the vertex of U adjacent to c_1 and c'_2 . Then $\{c_1, u, u', c_3\}$ is a claw, a contradiction. \square

Let us call a class of graphs \mathcal{C} hereditary if, for every $G \in \mathcal{C}$, every induced subgraph of G also belongs to \mathcal{C} . The following is a useful fact about HCH graphs:

Proposition 21. *Let \mathcal{L} be a hereditary graph class, such that every graph in \mathcal{L} is HCH, and every interesting graph in \mathcal{L} is K -perfect. Then every interesting graph in \mathcal{L} is clique-perfect.*

Proof. Let G be an interesting graph in \mathcal{L} . Let H be an induced subgraph of G . Since \mathcal{L} is hereditary, H is an interesting graph in \mathcal{L} , so it is K -perfect. Since every graph in \mathcal{L} is HCH, it follows that H is CH, and so $\alpha_C(H) = \alpha(K(H)) = k(K(H)) = \tau_C(H)$ [4], and the result follows. \square

In general, the class of clique-perfect graphs is neither a subclass nor a superclass of the class of K -perfect graphs. It is not difficult to verify that the 3-sun or 0-pyramid (Fig. 8) is K -perfect but it is not clique-perfect and, on the other hand, the graph in Fig. 9 is clique-perfect but its clique graph contains a hole of length 5. However, we will prove that within the classes of graphs analyzed in this paper, clique-perfect graphs are also K -perfect.

3.1. Line graphs

First, we prove that interesting line graphs are K -perfect.

Proposition 22. *A line graph is interesting if and only if it has no induced subgraph isomorphic to an odd hole or a 3-sun.*

Proof. Since no line graph contains an antihole of length at least 7, and every line graph is claw-free, the result follows from Proposition 20. \square

Note that if $G = L(H)$ then G contains no odd hole if and only if H contains no odd cycle of length at least 5 as a subgraph. A trinity is the complement of the 3-sun, and its line graph is also the 3-sun. Moreover, the trinity is the only graph whose line graph is the 3-sun. Therefore, G does not contain a 3-sun if and only if H does not contain a trinity as a subgraph.

Theorem 23. *If G is a line graph and G contains no odd holes, then $K(G)$ is perfect.*

Proof. The proof is by induction on $|V(G)|$. The theorem holds for the graph with one vertex, and in each case we will reduce the K -perfection of G to the K -perfection of some proper induced subgraphs of G . Since every induced subgraph of a line graph with no odd holes is also a line graph with no odd holes, the result will then follow from the inductive hypothesis.

Let $G = L(H)$. By Lemma 15, we may assume H is connected. Since G has no odd holes, it follows that all the odd cycles of H are triangles. So by Theorem 17 either H is a bipartite graph, or H is a complete of size four, or H consists

of exactly $p + 2$ vertices x_1, \dots, x_p, a, b , such that $\{x_1, \dots, x_p\}$ is a stable set, and $\{x_j, a, b\}$ is a triangle for each $j = 1, \dots, p$, or H has a cutpoint.

If H is bipartite then $G = K(H)$ and $K(G) = K^2(H)$ is an induced subgraph of H [15], so it is bipartite and hence perfect.

If H is a complete of size four, then $K(L(H))$ is the complement of $4K_2$, and so it is perfect (it is the complement of a bipartite graph).

If H consists of exactly $p + 2$ vertices x_1, \dots, x_p, a, b , such that $\{x_1, \dots, x_p\}$ is a stable set, and $\{x_j, a, b\}$ is a triangle for each $j = 1, \dots, p$, then all the cliques of G contain the vertex corresponding to the edge ab of H , so $K(G)$ is a complete graph, and hence perfect.

Suppose H has a cutpoint x , and let M_x be the complete subgraph of G induced by the vertices corresponding to the edges of H incident with x . Since x is a cutpoint of H , M_x is a clique of G , and let v be the vertex of $K(G)$ corresponding to M_x .

If $H = H_1 + H_2 + x$ and both H_1 and H_2 have at least one edge, then v is a cutpoint of $K(G)$, and $K(G) = M_1 + M_2 + v$, where M_i is the clique graph of the line graph of the subgraph of H formed by H_i and the edges incident with x with their respective endpoints. So the property follows from Theorem 7 by the inductive hypothesis.

Otherwise, x is adjacent to at least one vertex y of degree one in H . Let M'_x be the complete subgraph of $L(H \setminus \{y\})$ induced by the vertices corresponding to the edges of $H - \{y\}$ incident with x . If M'_x is still a clique of $L(H \setminus \{y\})$, then $K(G) = K(L(H \setminus \{y\}))$, and the property holds by the inductive hypothesis.

If M'_x is not a clique in $L(H \setminus \{y\})$, then x has degree three in H , and the other two neighbors z and w of x in H are adjacent. The cliques meeting M_x in G pairwise intersect (all of them contain the vertex corresponding to the edge wz of H), so v is simplicial in $K(G)$. On the other hand, $K(L(H \setminus \{y\})) = K(G) \setminus \{v\}$, so the property follows from Lemma 6 by the inductive hypothesis. \square

Theorem 18 is an immediate corollary of the following:

Theorem 24. *Let G be a line graph. Then the following are equivalent:*

- (i) *No induced subgraph of G is an odd hole, or a 3-sun.*
- (ii) *G is clique-perfect.*
- (iii) *G is perfect and it does not contain a 3-sun.*

Proof. The equivalence between (i) and (iii) is a corollary of Theorem 17. From Theorem 3 it follows that (ii) implies (i).

It therefore suffices to prove that (i) implies (ii). This proof is again by induction on $|V(G)|$. The class of line graphs with no odd holes or induced 3-suns is hereditary, so we only have to prove that for every graph in this class τ_C equals α_C . By Theorem 23 and Proposition 22, every such graph is K -perfect. So, by Proposition 21, an interesting HCH line graph is clique-perfect. Let $G = L(H)$ and suppose that G is not HCH. Then G contains a 0-, 1-, 2-, or 3-pyramid (see Fig. 2).

A 0-pyramid is a 3-sun. A 2-pyramid is not a line graph, and therefore is not an induced subgraph of G .

Suppose that G contains a 3-pyramid. This happens if and only if H contains a complete set of size four, say K . By Lemma 15 we may assume H is connected. We analyze how vertices of $V(H) \setminus K$ attach to K . If a vertex v is adjacent to two different vertices of K , then H contains an odd cycle as a subgraph and G contains an odd hole. If two different vertices v, w are adjacent to two different vertices of K , then H contains a trinity as a subgraph and so G contains a 3-sun. These cases can be seen in Fig. 10.

So, only one of the four vertices x_1, x_2, x_3, x_4 of K may have neighbors in $H \setminus K$, say x_1 . Let v, w, z_1, z_2, z_3 , and z_4 be the vertices of G corresponding to the edges $x_1x_2, x_3x_4, x_1x_3, x_1x_4, x_2x_4$, and x_2x_3 of H , respectively. The vertex w is adjacent in G only to z_1, z_2, z_3 , and z_4 . These four vertices induce a hole of length 4 and are adjacent also to v . So $G \setminus \{w\}$ is a clique subgraph of G (every clique of $G \setminus \{w\}$ is a clique of G). On the other hand, since x_2 has no neighbors in $H \setminus K$, all the neighbors of v other than z_3 and z_4 are vertices corresponding to edges of H containing x_1 , and they are a complete in G . This situation can be seen in Fig. 11.

By the inductive hypothesis, $G \setminus \{w\}$ is clique-perfect. Let A be a maximum clique-independent set and T be a minimum clique-transversal of $G \setminus \{w\}$. By maximality and by the structure of G , A has exactly one clique containing v .




Fig. 10. How the remaining vertices of H can be attached to the K_4 .




Fig. 11. Structure of G when H has a K_4 .




Fig. 12. Subgraph of H when H contains no K_4 and G contains a 1-pyramid.

Adding w , four new cliques appear, each one disjoint from a different one of the four cliques containing v , and adding w to T we have a clique-transversal of G , so $\alpha_C(G) = \alpha_C(G \setminus \{w\}) + 1 = \tau_C(G \setminus \{w\}) + 1 = \tau_C(G)$. So we may assume that H contains no complete set of size four.

Suppose finally that G contains a 1-pyramid. Since G contains a 1-pyramid, H contains as a subgraph a graph on five vertices v_1, \dots, v_5 where v_1 is adjacent to v_2, v_3 and v_4 , v_2 is adjacent to v_3 and v_4 , and v_3 is adjacent to v_5 (Fig. 12). Moreover, v_3 and v_4 are not adjacent because H does not contain a complete set of size four; v_1 and v_2 are not adjacent to v_5 , otherwise H contains an odd cycle as a subgraph; and v_1 and v_2 do not have other neighbors, otherwise H contains a trinity as a subgraph. Then v_1 and v_2 form a cutset in H , because if there is a path $v_3 P v_4$ in $H \setminus \{v_1, v_2\}$, then either $v_3 P v_4 v_1 v_3$ or $v_3 P v_4 v_1 v_2 v_3$ is an odd cycle in H .

Let w_1, \dots, w_5 be the vertices of G corresponding to the edges $v_1 v_3, v_2 v_3, v_1 v_4, v_2 v_4$, and $v_1 v_2$ of H , respectively. Then $w_1 w_2 w_4 w_3 w_1$ is a hole of length 4 in G , w_5 is adjacent only to w_1, w_2, w_3, w_4 , and w_2, w_3, w_5 is a cutset of G . The remaining neighbors of w_1 or w_2 are adjacent to both w_1 and w_2 , and form a non-empty complete in G (they are the vertices corresponding to the edges of H containing v_3 and not v_1 or v_2 , and there exists at least one such edge, namely the edge $v_3 v_5$). Similarly, the remaining neighbors of w_3 or w_4 are adjacent to both w_3 and w_4 , and form a (possibly empty) complete in G . The structure of G in this case can be seen in Fig. 13.

We show that $\alpha_C(G) = \alpha_C(G')$ and $\tau_C(G) = \tau_C(G')$, where G' is the line graph of the graph H' obtained from H by deleting the edges $v_2 v_3$ and $v_1 v_4$. So $G' = G \setminus \{w_2, w_3\}$.




Fig. 13. Structure of G when H has no K_4 .

Since every clique-transversal of G' either contains w_5 , or contains both w_1 and w_4 , it follows that every clique-transversal of G' is a clique-transversal of G . On the other hand, starting with a clique-transversal T of G and replacing the vertices w_2 and w_3 by w_1 and w_4 , respectively, if w_2 or w_3 belong to T , produce a clique-transversal of G' . Therefore $\tau_C(G) = \tau_C(G')$.

We claim that there is a maximum clique-independent set of G not containing either of the cliques $\{w_1, w_3, w_5\}$, $\{w_2, w_4, w_5\}$. Suppose the claim is false. Let I be a clique-independent set of G , we may assume I contains the clique $\{w_1, w_3, w_5\}$. Then I does not contain any other clique containing w_1 or w_5 ; and since the only clique containing w_2 and not w_1 is $\{w_2, w_4, w_5\}$, it follows that every clique in I is disjoint from $\{w_1, w_2, w_5\}$. But now the set obtained from I by removing the clique $\{w_1, w_3, w_5\}$ and adding the clique $\{w_1, w_2, w_5\}$ has the desired property. This proves the claim.

Let I be a maximum clique-independent set of G not containing either of the cliques $\{w_1, w_3, w_5\}$, $\{w_2, w_4, w_5\}$. Let I' be a set of cliques of G' obtained from I by replacing the clique $\{w_1, w_2, w_5\}$ by $\{w_1, w_5\}$ if $\{w_1, w_2, w_5\} \in I$, and the clique $\{w_3, w_4, w_5\}$ by $\{w_4, w_5\}$ if $\{w_3, w_4, w_5\} \in I$. On the other hand, clearly every clique-independent-set of G' gives rise to a clique-independent set of G , and therefore $\alpha_C(G) = \alpha_C(G')$.

But now, since G' is a proper induced subgraph of G , it follows inductively that $\alpha_C(G') = \tau_C(G')$, and therefore $\alpha_C(G) = \tau_C(G)$. This completes the proof of Theorem 24. \square

The recognition of clique-perfect line graphs can be solved in linear time in the following way. Given a graph G , in linear time we can obtain a graph H such that $L(H) = G$, or deduce that such a graph does not exist [20]. Now, by Theorems 24 and 17, and since G contains a 3-sun if and only if H contains a trinity as a subgraph, it suffices to check if H contains an odd cycle of length at least 5 or a trinity as a subgraph. It can be done in linear time in the number of edges of H , which is the number of vertices of G , combining the ideas in the proofs of Theorems 17 and 24.

3.2. HCH claw-free graphs

Let us first describe interesting HCH claw-free graphs.

Proposition 25. *No HCH graph contains an antihole of length at least 8. An HCH claw-free graph is interesting if and only if it does not contain an odd hole or an antihole of length 7.*

Proof. Since by Theorem 4 an HCH graph contains no induced subgraph isomorphic to one of the graphs of Fig. 2, it follows that no HCH graph contains a 3-sun. Since every antihole of length at least 8 contains a 2-pyramid, it follows that no HCH graph contains an antihole of length at least 8. Finally, since by Proposition 20, every claw-free odd generalized sun is either an odd hole or a 3-sun, it follows that an HCH claw-free graph is interesting if and only if it contains no odd hole and no antihole of length 7. \square

We will use Proposition 21 to prove the characterization for HCH claw-free graphs, so first we need to prove the following.

Theorem 26. *Let G be an interesting HCH claw-free graph. Then $K(G)$ is perfect.*

In the remainder of this section we use the structure theorem for claw-free graphs (Theorem 5) to prove that every interesting HCH claw-free graph G is K -perfect. The proof is by induction on $|V(G)|$.

3.2.1. Circular interval graphs

First we prove that clique graphs of interesting HCH circular interval graphs are perfect.

Lemma 27. *Let G be a circular interval graph. Then $K(G)$ is an induced subgraph of G .*

Proof. Let G be a circular interval graph with vertices v_1, \dots, v_n in clockwise order, say. We define a homomorphism v from $V(K(G))$ to $V(G)$ (meaning that for two distinct vertices $a, b \in V(K(G))$, $v(a) \neq v(b)$; and a is adjacent to b if and only if $v(a)$ is adjacent to $v(b)$). For every clique M of G , since no three intervals in the definition of a circular interval graph cover the circle, $M = \{v_i, \dots, v_{i+t}\}$ (where the indices are taken mod n). In this case we say that v_i is the *first vertex* of M . We define $v(M) = v_i$. Since v_i is the first vertex of a unique clique, it follows that $v(M) \neq v(M')$ if M and M' are distinct cliques of G . It remains to show that $v(M)$ is adjacent to $v(M')$ if and only if $M \cap M' \neq \emptyset$. If M and M' intersect at a vertex v_k , then the clockwise order of $v(M)$, $v(M')$, and v_k is either $v(M)$, $v(M')$, v_k or $v(M')$, $v(M)$, v_k , and in both cases $v(M)$ and $v(M')$ are adjacent. On the other hand, if there are two cliques such that $v(M)$ and $v(M')$ are adjacent, we may assume $v(M)$ appears first clockwise in the circular interval which contains both $v(M)$ and $v(M')$. Then since $v(M)$ is the first vertex of the clique M , it follows that $v(M')$ belongs to M , so M and M' intersect. \square

Proposition 28. *Let G be an HCH interesting circular interval graph. Then $K(G)$ is perfect.*

Proof. By Lemma 27, $K(G)$ is an induced subgraph of G . Since G is HCH and interesting, it contains no odd hole and no antihole of length at least 7, and therefore it is perfect by Theorem 1. \square

3.2.2. Decompositions

Now we show that if an interesting HCH claw-free graph admits one of the decompositions of Theorem 5, then either it is K -perfect or we can reduce the problem to a smaller one.

Theorem 29. *Let G be an interesting HCH claw-free graph. If G admits a 1-join, then $K(G)$ has a cutpoint v , $K(G) = H_1 + H_2 + v$, and $H_i + v$ is the clique graph of a smaller interesting HCH claw-free graph.*

Proof. Since G admits a 1-join, it follows that $V(G)$ is the disjoint union of two non-empty sets V_1 and V_2 ; each V_i contains a complete M_i , such that $M_1 \cup M_2$ is a complete, and there are no other edges from V_1 to V_2 . So $M_1 \cup M_2$ is a clique in G . Let v be the vertex of $K(G)$ corresponding to $M_1 \cup M_2$. Every other clique of G is either contained in V_1 or in V_2 , and no clique of the first type intersects a clique of the second type. So v is a cutpoint of $K(G)$, and $K(G) = H_1 + H_2 + v$, where H_1 (H_2) is the subgraph of $K(G)$ induced by the vertices corresponding to cliques of G of the first (second) type. Let G_i be the graph obtained from $G|V_i$ by adding a vertex v_i complete to M_i and with no

other neighbors in G_i . Then G_i is isomorphic to an induced subgraph of G , so it is interesting, HCH, and claw-free, and, for $i = 1, 2$, $H_i + v$ is isomorphic to $K(G_i)$ (where the vertex v is mapped to the vertex of $K(G_i)$ corresponding to the clique $M_i \cup \{v_i\}$ of G_i). This proves Theorem 29. \square

Theorem 30. *Let G be an interesting HCH claw-free graph. If G admits a generalized 2-join and no twins, 0-join or 1-join, then there exist two clique graphs of smaller interesting HCH claw-free graphs, H_1 and H_2 , such that if H_1 and H_2 are perfect, then so is $K(G)$.*

Proof. Since G admits a generalized 2-join, it follows that $V(G)$ is the disjoint union of three sets V_0, V_1 , and V_2 ; for $i = 1, 2$ each V_i contains two disjoint completes A_i, B_i , such that A_i, B_i , and $V_i \setminus (A_i \cup B_i)$ are all non-empty, $A_1 \cup A_2 \cup V_0$ and $B_1 \cup B_2 \cup V_0$ are completes, and there are no other edges from V_1 to V_2 or from V_0 to $V_1 \cup V_2$. Since G admits no twins, it follows that $|V_0| \leq 1$.

So $A_1 \cup A_2 \cup V_0$ and $B_1 \cup B_2 \cup V_0$ are cliques of G , and they correspond to vertices w_1, w_2 of $K(G)$. Every other clique of G is either contained in V_1 or in V_2 , and no clique of the first type intersects a clique of the second type. So $\{w_1, w_2\}$ is a cutset in $K(G)$.

If V_0 is non-empty, then w_1 is adjacent to w_2 and $\{w_1, w_2\}$ is a clique cutset in $K(G)$. Let $V_0 = \{v_0\}$. Now $K(G) = M_1 + M_2 + \{w_1, w_2\}$, where, for $i = 1, 2$, $H_i = M_i + \{w_1, w_2\}$ is the clique graph of the subgraph of G induced by $V_i \cup \{v_0\}$. By Theorem 7, $K(G)$ is perfect if and only if H_1 and H_2 are. So we may assume that V_0 is empty, and therefore w_1 is non-adjacent to w_2 .

We start with the following easy observation:

- (*) Let S be a graph which is either a claw, or an odd hole, or $\overline{C_7}$, or a 0-, 1-, 2-, or 3-pyramid, and suppose there exists a vertex $s \in V(S)$, whose neighborhood is the union of two non-empty completes with no edges between them. Then S is an odd hole.

Since G admits no 0-join or 1-join, for $i = 1, 2$ there exist a_i in A_i and b_i in B_i joined by an induced path with interior in $V_i \setminus (A_i \cup B_i)$. (The interior of a path is the set of vertices different from the endpoints; the interior may be empty, if a_i and b_i are adjacent.)

Then, since G contains no odd hole, for every a_i in A_i and b_i in B_i , all induced paths from a_1 to b_1 with interior in $V_1 \setminus (A_1 \cup B_1)$ and all induced paths from a_2 to b_2 with interior in $V_2 \setminus (A_2 \cup B_2)$ have the same parity.

Case 1: This parity is even. Note that in this case A_i is anticomplete to B_i . Let H be the graph obtained from $K(G)$ by adding the edge $w_1 w_2$. Since A_i is anticomplete to B_i , there is no clique in G intersecting both $A_1 \cup A_2$ and $B_1 \cup B_2$. So w_1 and w_2 have no common neighbor in $K(G)$. By Theorem 8, if H is perfect then $K(G)$ is.

Construct graphs G_i with vertex set $V_i \cup \{v_i\}$, where $G_i|V_i = G|V_i$ and v_i is complete to $A_i \cup B_i$ and has no other neighbors in G_i . Now, $H = M_1 + M_2 + \{w_1, w_2\}$, with $M_i + \{w_1, w_2\} = K(G_i)$, and $\{w_1, w_2\}$ is a clique cutset in H . By Theorem 7, it follows that if $K(G_1)$ and $K(G_2)$ are perfect then H is perfect and thus $K(G)$ is perfect.

We claim that for $i = 1, 2$ the graphs G_i are claw-free, HCH, and interesting. Suppose that G_1 , say, is not. So G_1 contains an induced subgraph S isomorphic to a claw, an odd hole, $\overline{C_7}$, or a 0-, 1-, 2-, or 3-pyramid. If $V(S)$ does not contain v_1 , then S is isomorphic to an induced subgraph of G , a contradiction. If $V(S)$ contains v_1 but has empty intersection with A_1 or B_1 , say B_1 , then S is isomorphic to an induced subgraph of G , obtained by replacing v_1 by any vertex of A_2 , a contradiction. So $V(S)$ meets both A_1 and B_1 , and therefore the neighborhood of v_1 in S can be partitioned into two non-empty completes A_S, B_S , such that A_S is anticomplete to B_S . By (*), S is an odd hole. Let $a_1 \in A_1$ and $b_1 \in B_1$ be the neighbors of v_1 in S . Then $S \setminus \{v_1\}$ is an induced odd path from a_1 to b_1 with interior in $V_1 \setminus (A_1 \cup B_1)$, a contradiction.

Case 2: This parity is odd. Construct graphs G_i with vertex set $V_i + \{v_{A,i}, v_{B,i}\}$, where $G_i|V_i = G|V_i$, $v_{A,i}$ is complete to A_i , $v_{B,i}$ is complete to B_i , $v_{A,i}$ is adjacent to $v_{B,i}$, and there are no other edges in G_i . Now, $K(G) = M_1 + M_2 + \{w_1, w_2\}$, and $K(G_i)$ is obtained from $M_i + \{w_1, w_2\}$ by joining w_1 and w_2 by an induced path of length 2. By Theorem 9, if $K(G_1)$ and $K(G_2)$ are perfect, so is $K(G)$.

We claim that both G_i are claw-free, interesting, and HCH. Suppose that G_1 contains an induced subgraph S isomorphic to a claw, an odd hole, $\overline{C_7}$, or a 0-, 1-, 2-, or 3-pyramid.

If $V(S)$ does not contain $v_{A,1}$ or $v_{B,1}$, say $v_{B,1}$, then S is isomorphic to an induced subgraph of G , obtained by replacing $v_{A,1}$ by any vertex of A_2 , a contradiction. If $V(S)$ contains $v_{A,1}$ and $v_{B,1}$ but has empty intersection with A_1

or B_1 , say B_1 , then S is isomorphic to an induced subgraph of G , obtained by replacing $v_{A,1}$ and $v_{B,1}$ by two adjacent vertices a_2, c_2 of V_2 such that $a_2 \in A_2$ and $c_2 \in V_2 \setminus A_2$ (such a pair of vertices exist because there is at least one path from A_2 to B_2 in G), a contradiction. So $V(S)$ meets both A_1 and B_1 , and the neighborhood of $v_{A,1}$ in S can be partitioned into two non-empty completes with no edges between them, namely $A_S = A_1 \cap V(S)$ and $\{v_{B,1}\}$. By (*) S is an odd hole. Let $a_1 \in A_1$ and $b_1 \in B_1$ be the neighbors of $v_{A,1}$ and $v_{B,1}$ in $V(S) \cap V_1$, respectively. Then $S \setminus \{v_{A,1}, v_{B,1}\}$ is an induced even path from a_1 to b_1 with interior in $V_1 \setminus (A_1 \cup B_1)$, a contradiction. This concludes the proof of Theorem 30. \square

Lemma 31. *Let G be an HCH graph such that \overline{G} is a bipartite graph. Then $K(G)$ is perfect.*

Proof. In this proof we use the vertices of $K(G)$ and the cliques of G interchangeably. By Theorem 1, if $K(G)$ is not perfect then it contains an odd hole or an odd antihole.

Let A, B be two disjoint completes of G such that $A \cup B = V(G)$. If there exists a vertex v of G adjacent to every other vertex in G , then v belongs to every clique of G and $K(G)$ is a complete graph, and therefore perfect. So we may assume that no vertex of A is complete to B and no vertex of B is complete to A . Then A and B are cliques of G , and every other clique of G meets both A and B . The degree of A and B in $K(G)$ is $|V(K(G))| - 1$, so they cannot be part of an odd hole or an odd antihole in $K(G)$.

It is therefore enough to show that there is no odd hole or antihole in the graph obtained from $K(G)$ by deleting the vertices A and B . We prove a stronger statement, namely that there is no induced path of length 2 in this graph. Since every hole and antihole of length at least 5 contains a two-edge path, the result follows.

Suppose for a contradiction that there are three cliques X, Y , and Z in G , each meeting both A and B , and such that X is disjoint from Z , and both $X \cap Y$ and $Y \cap Z$ are non-empty. From the symmetry we may assume that $X \cap Y$ contains a vertex $a_{xy} \in A$.

Suppose first that there is a vertex $a_{yz} \in A \cap Y \cap Z$. Let b_y be a vertex in $Y \cap B$. Since no vertex of B is complete to A , there is a vertex a in A non-adjacent to b_y . Since a_{yz} does not belong to X , there is a vertex b_x in X non-adjacent to a_{yz} , and since A is a complete, b_x belongs to B . Analogously, since a_{xy} does not belong to Z , there is a vertex b_z in $B \cap Z$ non-adjacent to a_{xy} . But now $\{a_{xy}, a_{yz}, b_y, b_z, b_x, a\}$ induce a 1-, 2-, or 3-pyramid, a contradiction.

So $A \cap Y \cap Z$ is empty, and therefore $B \cap Y \cap Z$ is non-empty, and, by the argument of the previous paragraph with A and B exchanged, $B \cap X \cap Y$ is empty. Choose b_{yz} in $B \cap Y \cap Z$. Choose a_z in $Z \cap A$, then $a_z \notin X \cup Y$. Since a_z does not belong to X , there is a vertex $b_x \in X$ non-adjacent to a_z , and, since A is a complete, b_x is in B . Since b_{yz} does not belong to X and B is a complete, there is a vertex $a_x \in A \cap X$ non-adjacent to b_{yz} ; and since a_{xy} does not belong to Z and A is a complete, there is a vertex $b_z \in B \cap Z$ non-adjacent to a_{xy} . But now $\{a_z, a_{xy}, b_{yz}, a_x, b_x, b_z\}$ induces a 2- or a 3-pyramid, a contradiction. This proves Lemma 31. \square

Theorem 32. *Let G be a connected interesting HCH claw-free graph, and suppose G admits no twins. Assume that G admits a coherent or a non-dominating W -join (A, B) . Then either $K(G)$ is perfect, or there exist induced subgraphs G_1, \dots, G_k of G , each smaller than G , such that if G_i is K -perfect for every $i = 1, \dots, k$, then so is G .*

Proof. Choose a coherent or non-dominating W -join (A, B) with $A \cup B$ minimal. Let C be the vertices complete to A and anticomplete to B , D be the vertices complete to B and anticomplete to A , E be the vertices complete to $A \cup B$, and F be the vertices anticomplete to $A \cup B$. Since the W -join (A, B) is either coherent or non-dominating, it follows that either E is a complete, or F is non-empty.

32.1. $A \cup C, B \cup D$ are both completes, and E is anticomplete to F .

Suppose not. Assume first that there exist two non-adjacent vertices c_1, c_2 in C . Choose a in A and b in B such that a is adjacent to b ; now $\{a, c_1, c_2, b\}$ is a claw, a contradiction. So C is a complete, and since A is a complete, it follows that $A \cup C$ is a complete. From the symmetry it follows that $B \cup D$ is a complete.

Next assume that there are two adjacent vertices e in E and f in F . Choose a in A and b in B such that a is not adjacent to b . Then $\{e, a, b, f\}$ is a claw, a contradiction. This proves 32.1

Let E_1 be a clique of $G|E$. Let \mathcal{L} be the set of all cliques of $G|(A \cup B)$. Let

$$U = \{E_1 \cup L : L \in \mathcal{L} \text{ and } L \neq A, B\}.$$

Since E is anticomplete to F , and every member of U meets both A and B , it follows that the members of U are cliques of G .

32.2. We may assume that $|U| \geq 2$.

Suppose $|U| \leq 1$. Since in G there is at least one edge between A and B , it follows that there is a unique clique L in $G|(A \cup B)$ meeting both A and B , and $|U| = 1$. Let $A' = A \cap L$, $B' = B \cap L$. Then A' is complete to B' , $A \setminus A'$ is anticomplete to B , and $B \setminus B'$ is anticomplete to A . Since G does not admit twins, each of A' , $A \setminus A'$, B' , $B \setminus B'$ has size at most one, and by the minimality of $A \cup B$ at most one of $A \setminus A'$, $B \setminus B'$ is non-empty. By the symmetry, we may assume that $B \setminus B'$ is empty and $|A'| = |B'| = |A \setminus A'| = 1$. Let $A' = \{a_1\}$, $B' = \{b_1\}$, and $A \setminus A' = \{a_2\}$.

If $K(G \setminus \{a_2\}) = K(G)$ then the theorem holds, so we may assume not. Therefore, there exists a subset E' of E such that $M = A \cup E'$ is a clique of G . It follows, in particular, that no vertex of C is complete to E .

If $G|E$ is complete, consider the cliques $M_1 = \{a_1, b_1\} \cup E$ and $M_2 = \{a_1, a_2\} \cup E$ of G . Since every clique of G containing a_2 also contains a_1 , then every clique of G , that has a non-empty intersection with M_2 , meets M_1 . Therefore, the vertex w_1 of $K(G)$, corresponding to M_1 , dominates the vertex w_2 of $K(G)$, corresponding to M_2 . Since $K(G) \setminus \{w_1\}$ is an induced subgraph of $K(G \setminus \{a_1\})$ and $K(G) \setminus \{w_2\} = K(G \setminus \{a_2\})$, by Theorem 11, $K(G)$ is perfect if $K(G \setminus \{a_1\})$ and $K(G \setminus \{a_2\})$ are and the theorem holds. So we may assume that E is not a complete.

Next we claim that D is empty. Since E is not a complete, there are two non-adjacent vertices e_1, e_2 in E , and let d in D . If d is non-adjacent to both of e_1 and e_2 , then $\{b_1, e_1, e_2, d\}$ is a claw, a contradiction. Otherwise, $\{b_1, e_1, e_2, d, a_1, a_2\}$ induces a 1- or 2-pyramid, a contradiction. This proves that D is empty.

Since D is empty, every clique disjoint from F contains the vertex a_1 , and, since every clique containing a vertex of F is disjoint from A, B , and E , it follows that the vertices of $K(G)$ corresponding to the cliques $\{a_1, b_1\} \cup E'$, with E' a clique of $G|E$, are simplicial in $K(G)$. By Lemma 6, $K(G)$ is perfect if and only if $K(G \setminus \{b_1\})$ is. This proves 32.2.

32.3. We may assume that no vertex of B is complete to A , and no vertex of A is complete to B .

Suppose there is a vertex $b \in B$ complete to A . Since A is not complete to B , there is a vertex $b' \in B \setminus \{b\}$. By 32.2, $|A| > 1$. But now $(A, B \setminus \{b\})$ is a coherent or non-dominating W -join in G , contrary to the minimality of $A \cup B$. This proves 32.3.

In view of 32.2 and 32.3, we henceforth assume that $|U| \geq 2$, no vertex of A is complete to B , and no vertex of B is complete to A .

32.4. $G|E$ is complete.

Since no vertex of B is complete to A , and there is at least one edge between A and B , there is a vertex $a_1 \in A$ with a neighbor b_1 and a non-neighbor b_2 in B . Since b_1 is not complete to A , there is a vertex $a_2 \in A$, non-adjacent to b_1 . Since A, B are both cliques, a_1 is adjacent to a_2 and b_1 to b_2 . If there exist two non-adjacent vertices e_1 and e_2 in E , then $\{a_1, a_2, b_1, b_2, e_1, e_2\}$ induces a 2- or a 3-pyramid in G , a contradiction. This proves 32.4.

32.5. Every vertex of $K(G) \setminus U$ with a neighbor in U is complete to U .

Throughout the proof of 32.5 we use cliques of G and vertices of $K(G)$ interchangeably.

It follows from 32.4 that $E_1 = E$. Let w be a vertex of $K(G) \setminus U$ with a neighbor in U . Since w has a neighbor in U , it follows that w meets one of A, B, E . If w meets E , then w is complete to U and the result follows. If w includes one of A, B , then since every member of U meets each of A, B , we again deduce that w is complete to U and the result follows. So we may assume that w is disjoint from E , and the sets $w \cap (A \cup B)$, $A \setminus \{w\}$, and $B \setminus \{w\}$ are all non-empty.

Assume first that w meets both A and B . Since w is a clique of G , $C \cup F$ is anticomplete to B , and $D \cup F$ is anticomplete to B , it follows that $w \subseteq A \cup B \cup E$. But now, since w is a clique, it follows that w includes E and w belongs to U , a contradiction. So we may assume that w is disjoint from at least one of A and B .

By the symmetry we may assume that w is disjoint from B , and therefore w meets A . Since $F \cup D$ is anticomplete to A , it follows that w is a subset of $A \cup C \cup E$, and, since w is a clique, w includes A , a contradiction. This proves 32.5.

32.6. U is a homogeneous set in $K(G)$ and the graph $K(G)|U$ is perfect.

It follows from 32.5 that U is a homogeneous set in $K(G)$. The graph $K(G)|U$ is isomorphic to the graph obtained from $K(G|(A \cup B \cup E))$ by deleting the vertices corresponding to the cliques $A \cup E$ and $B \cup E$. Since $\overline{G|(A \cup B \cup E)}$ is bipartite, it follows from Lemma 31 that $K(G)|U$ is perfect. This proves 32.6.

Choose $u \in U$.

32.7. If there exist $a_1, a_2 \in A$ and $b_1, b_2 \in B$ such that a_1 is adjacent to b_1 and not to b_2 , and a_2 is adjacent to b_2 and not to b_1 , then either $K(G)$ is perfect, or there is an induced subgraph G' of G such that $K(G) \setminus (U \setminus \{u\}) = K(G')$.

If there exist non-adjacent $c \in C$ and $e \in E$, then $\{a_1, a_2, e, c, b_1, b_2\}$ induces a 1-pyramid, a contradiction, so C is complete to E , and similarly D is complete to E . By 32.4, $G|E$ is complete. Since G admits no twins, $|E| \leq 1$. If $C \cup D$ is empty, then, since G is connected, F is empty, and G is the complement of a bipartite graph. By Lemma 31, $K(G)$ is perfect. So we may assume that C is non-empty, and, in particular, $A \cup E$ is not a clique of G . But now $K(G) \setminus (U \setminus \{u\}) = K(G \setminus ((A \cup B) \setminus \{a_1, b_1, b_2\}))$. This proves 32.7.

To finish the proof, let $a_1 \in A$ and $b_1 \in B$ be adjacent. By 32.3, there exist a vertex $b_2 \in B$, non-adjacent to a_1 and a vertex $a_2 \in A$ non-adjacent to b_1 . If a_2 is adjacent to b_2 , then the theorem follows from 32.6, 32.7, and Theorem 10. So we may assume that a_2 is non-adjacent to b_2 . Let $G' = G \setminus ((A \cup B) \setminus \{a_1, b_1, a_2, b_2\})$. We deduce from 32.2 that G' is smaller than G . Moreover, G' is an induced subgraph of G . But $K(G) \setminus (U \setminus \{u\}) = K(G')$, and, together with 32.6 and Theorem 10, this implies that the theorem holds. This proves Theorem 32. \square

Theorem 33. Let G be an interesting HCH claw-free graph. Suppose G admits a hex-join and no twins and every vertex of G is in a triad. Then $G = C_6$.

Proof. Since G admits a hex-join, there exist six completes $A_1, B_1, C_1, A_2, B_2, C_2$ in G such that A_1 is anticomplete to A_2 and complete to B_2 and C_2 ; B_1 is anticomplete to B_2 and complete to A_2 and C_2 ; C_1 is anticomplete to C_2 and complete to A_2 and B_2 ; $A_1 \cup B_1 \cup C_1$ and $A_2 \cup B_2 \cup C_2$ are non-empty; and $V(G) = A_1 \cup B_1 \cup C_1 \cup A_2 \cup B_2 \cup C_2$. Since every vertex of G is in a triad and no stable set of size three meets both $A_1 \cup B_1 \cup C_1$ and $A_2 \cup B_2 \cup C_2$, it follows that A_i, B_i, C_i are all non-empty.

Suppose there is an edge $a_1 b'_1$ with a_1 in A_1 and b'_1 in B_1 . Since every vertex is in a triad, there exists a stable set $\{a_2, b_2, c_2\}$ with a_2 in A_2 , b_2 in B_2 , and c_2 in C_2 , and a stable set $\{a_1, b_1, c_1\}$ with a_1 in A_1 , b_1 in B_1 , and c_1 in C_1 . Since G is interesting, $a_1 b'_1 a_2 c_1 b_2 a_1$ is not a hole in G , so b'_1 is adjacent to c_1 . But now $\{b'_1, a_1, b_1, c_1\}$ is a claw in G , a contradiction. So A_1 is anticomplete to B_1, C_1 . Since the vertices of A_1 are not twins in G , it follows that $|A_1| = 1$. From the symmetry, $|A_i| = |B_i| = |C_i| = 1$ for $i = 1, 2$, and $G = C_6$. This proves Theorem 33. \square

Theorem 34. Let G be an interesting HCH graph. Assume that G admits no twins and no coherent or non-dominating W -join, and contains no stable set of size three. Then $K(G)$ is perfect.

Proof. Since G is claw-free, we may assume G contains either a 4-wheel or a 3-fan, otherwise, by Theorem 16, $K(G)$ is bipartite.

Case 1: G contains a 4-wheel. Let $a_1 a_2 a_3 a_4 a_1$ be a hole and let c be adjacent to all a_i . We claim every vertex in G is adjacent to c . Suppose v is non-adjacent to c . Then since G contains no stable set of size three, from the symmetry we may assume v is adjacent to a_1, a_2 . But now $\{a_1, a_2, a_3, a_4, c, v\}$ induces a 1-, 2-, or 3-pyramid, a contradiction. So every clique in G contains c ; then $K(G)$ is a complete graph and the result follows. This proves Case 1.

Case 2: G contains a 3-fan and no 4-wheel. Let A_1, \dots, A_k be anticonnected sets in G , pairwise complete to each other, with $k > 2$, $|A_1| > 1$, and subject to that with maximal union, say A . (Such sets exist because there is a 3-fan. Let $c_1 c_2 c_3 c_4$ be a path and let c be adjacent to all c_i . Then $A_1 = \{c_1, c_3\}$, $A_2 = \{c_2\}$, $A_3 = \{c\}$ make a family of sets with the desired properties.)

Suppose $|A_2| > 1$. Then, since A_1, A_2 are both anticonnected, each of A_1, A_2 contains a non-edge, say $a_i b_i$. Choose a_3 in A_3 . Now $\{a_1, a_2, b_1, b_2, a_3\}$ is a 4-wheel, a contradiction. So for $2 \leq i \leq k$, $|A_i| = 1$, and let $A_i = \{a_i\}$.

(*) No vertex in $V(G) \setminus A$ is complete to more than one of A_1, \dots, A_k .

Let v be a vertex in $V(G)\setminus A$ and define $I = \{i : 1 \leq i \leq k \text{ and } v \text{ is complete to } A_i\}$ and $J = \{j : 1 \leq j \leq k \text{ and } v \text{ has a non-neighbor in } A_j\}$. Suppose $|I| > 1$. Define $A'_i = A_i$ for $t \in I$ and $A'_j = \bigcup_{j \in J} A_j \cup \{v\}$. Then $\{A'_i\}_{i \in I}$, A'_j is a collection of at least three anticonnected sets, pairwise complete to each other, but their union is a proper superset of A , contrary to the maximality of A . This proves (*).

(**) There is no C_4 in A_1 .

Otherwise, G contains a 4-wheel with center a_2 , a contradiction. This proves (**).

Since $|A_1| > 1$ and A_1 is anticonnected, A_1 contains a non-edge, and so, since there is no stable set of size three in G , every vertex of $V(G)\setminus A$ has a neighbor in A_1 . Let $A' = A \setminus A_1$. If no vertex of $V(G)\setminus A$ has a neighbor in A' , then the vertices of A' are twins (they are pairwise adjacent, complete to A_1 , and anticomplete to $V(G)\setminus A$), a contradiction.

So there exists v in $V(G)\setminus A$ with a neighbor in A_1 and a neighbor a' in A' . By (*) v has a non-neighbor a'' in A' . If v has two non-adjacent neighbors in A_1 , say x, y , then $xvya''x$ is a 4-hole and a' is complete to it, so G contains a 4-wheel, a contradiction. So the neighbors of v in A_1 are a complete. Since G has no stable set of size three, the non-neighbors of v in A_1 are a complete. Thus, $G|_{A_1}$ is the union of two completes (complement bipartite), and since it is anticonnected the bipartition is unique, say X, Y , both X and Y are non-empty, and every vertex of $V(G)\setminus A$ with a neighbor in A' is either complete to X and anticomplete to Y , or complete to Y and anticomplete to X . Let X' be the vertices with a neighbor in A' and complete to X , Y' be the vertices with a neighbor in A' and complete to Y . Then, $X' \cup Y'$ is non-empty, and since there is no stable set of size three in G , X', Y' are both completes.

For $i = 2, \dots, k$ let X_i be the vertices of X' adjacent to a_i , and let Y_i be defined similarly. By (*), $X_i \cap X_j = \emptyset$ for $i \neq j$, and the same holds for Y_i, Y_j . If there is an edge from X to Y then there is no edge from X_i to Y_i , or else G contains a 4-wheel with center a_i .

34.1. $k \leq 4$ and $X' = X_i, Y' = Y_j$ for some i different from j .

Suppose both X_2, X_3 are non-empty, choose x_2 in X_2 and x_3 in X_3 . Then $a_2x_2x_3a_3a_2$ is a hole of length 4, and every x in X is complete to it, so G contains a 4-wheel, a contradiction. So we may assume that $X' = X_2$ and, similarly, $Y' = Y_j$ for some j . If Y_2 is non-empty, then since x_2, y_2, a_3 is not a stable set of size three, x_2 is adjacent to y_2 . Since A_1 is anticonnected, there exist non-adjacent vertices $x \in X$ and $y \in Y$. But now $xx_2y_2ya_3x$ is a hole of length 5, a contradiction. So Y_2 is empty and therefore i is different from j , say $j = 3$. Since a_4, a_5 are not twins, $k \leq 4$. This proves 34.1.

By 34.1 we may assume that $X' = X_2, Y' = Y_3$. Let Z be the vertices of G with no neighbor in A' . Then, since G contains no triad, Z is a complete.

34.2. Every vertex in Z is complete to $X' \cup Y'$ and to one of X, Y .

If some vertex z in Z has a non-neighbor x_2 in X_2 , then z, x_2, a_3 is a stable set of size three, a contradiction, so Z is complete to X' , and similarly to Y' . Next suppose some vertex z in Z has a non-neighbor x in X and a non-neighbor y in Y . Then x is adjacent to y , and there is an odd antipath Q from x to y in $X \cup Y$. By (**) $X \cup Y$ contains no C_4 , so Q has length 3, say $Q = xy'x'y$. Since there is no stable set of size three, z is adjacent to y' and x' . But then $zx'xyy'z$ is a hole of length 5, a contradiction. This proves 34.2.

Let Z_x be the vertices of Z complete to X , and let $Z_y = Z \setminus Z_x$.

34.3. If Z, X', Y' are all non-empty then the theorem holds.

We may assume Z_x is non-empty. Since $a_2x_2zy_3a_3a_2$ (where $z \in Z, x_2 \in X_2$, and $y_3 \in Y_3$) is not a hole of length 5, X_2 is complete to Y_3 . Suppose z in Z_x has a neighbor y in Y . Since A_1 is anticonnected, y has a non-neighbor x in X . But now $a_3za_2y_3xyx_2a_3$ (with x_2 in X_2 and y_3 in Y_3) is an antihole of length 7, a contradiction. So Z_x is anticomplete to Y . Choose z in Z_x and non-adjacent x in X and y in Y . Then zxa_2yy_3z is a hole of length 5, a contradiction. This proves 34.3.

34.4. If Z is empty then the theorem holds.

The pairs (X, Y) and (X_2, Y_3) are coherent homogeneous pairs, and, since G does not admit twins or a coherent W -join, all four of these sets have size ≤ 1 . Every vertex of G is adjacent to a_3 , except the vertex x_2 of X_2 , if X' is non-empty. So every clique of G contains either a_3 or x_2 , and therefore $K(G)$ is perfect (it is either a complete graph, or the complement of a bipartite graph). This proves 34.4.

In view of 34.4, we henceforth assume that $Z \neq \emptyset$. By 34.3 we may assume X' is empty, and so Y' is non-empty. By 34.1 we may assume $Y' = Y_3$. Since the vertices of Y_3 are not twins, $Y_3 = \{y_3\}$.

34.5. Z is complete to Y .

Suppose not. Choose z in Z , with a non-neighbor y in Y . Then z in Z_x . Since A_1 is anticonnected, y has a non-neighbor x in X . But now zxa_2yy_3z is a hole of length 5, a contradiction. This proves 34.5.

Let M be the set of vertices in X with a neighbor in Z . Suppose some z in Z has adjacent neighbors x in X and y in Y . Then zxa_3y_3z is a hole of length 4, and y is complete to it, so G contains a 4-wheel, a contradiction. This proves that M is anticomplete to Y . Now (Z, M) is a coherent homogeneous pair, and the same for $(X \setminus M, Y)$. Since G admits no twins and no coherent W -join, all four of these sets have size ≤ 1 . Also, since a_2 and a_4 are not twins, $k = 3$. Let $Z = \{z\}$. Every vertex of G different from z is adjacent to a_3 . So every clique of G contains either a_3 or z , and then $K(G)$ is perfect (it is the complement of a bipartite graph). This completes the proof of Theorem 34. \square

Theorem 35. *Let G be an interesting HCH claw-free graph, and suppose that G is connected, does not admit a coherent or non-dominating W -join, or a 1-join, or twins. If G contains a stable set of size three and a singular vertex, then $K(G)$ is perfect.*

Proof. The proof is by induction on $|V(G)|$. Assume that, for every smaller graph G' satisfying the hypotheses of the theorem, $K(G')$ is perfect. Let v be a singular vertex in G with maximum number of neighbors (there exists at least one singular vertex in G , by hypothesis). Let A be the set of neighbors of v and B be the set of its non-neighbors. Since v is singular, B is a complete.

Since G contains a stable set of size three, and every such set meets both A and B (because B is a complete, and G is claw-free), there exist vertices in B that are non-singular. Let U be the set of all such vertices.

35.1. If U is anticomplete to A then $K(G)$ is perfect.

Let $B_2 = B \setminus U$, so every vertex of B_2 is singular, and, since G is connected, B_2 is non-empty. Let a_1, a_2 be two non-adjacent vertices in A . If $b \in B_2$ is non-adjacent to both a_1, a_2 , then $\{b, a_1, a_2\}$ is a stable set of size three, and if b is adjacent to both a_1, a_2 then $\{b, a_1, a_2, u\}$ is a claw for every $u \in U$; in both cases we get a contradiction. So every vertex in B_2 is adjacent to exactly one of a_1, a_2 . Suppose there exist v_1, v_2 in B_2 with v_i adjacent to a_i . Then $v_1v_2a_2va_1v_1$ is a hole of length 5, a contradiction. So one of a_1, a_2 is anticomplete to B_2 , and therefore the other one is complete to B_2 . Let A_1 be the vertices in A complete to B_2 , A_2 be the vertices in A anticomplete to B_2 , and $A_3 = A \setminus (A_1 \cup A_2)$. It follows from the previous argument that $A_1 \cup A_3$ and $A_2 \cup A_3$ are both completes. If A_3 is non-empty, then $|B_2| > 1$ and (A_3, B_2) is a coherent W -join, a contradiction. So we may assume A_3 is empty. Now (A_1, A_2) is a coherent homogeneous pair, and all the vertices of each of U, B_2 are twins. So all these sets have size at most one and $K(G)$ is the clique graph of an induced subgraph of a four-edge path, and hence perfect. This proves 35.1.

So we may assume that there exists a non-singular vertex u in B with a neighbor in A . Let M be the set of neighbors of u in A , N the set of non-neighbors. Since u is non-singular, N contains two non-adjacent vertices x, y . Choose m in M . If m is adjacent to both x, y then $\{m, x, y, u\}$ is a claw. If m is non-adjacent to both x, y then $\{v, x, y, m\}$ is a claw. So every vertex in M is adjacent to exactly one of x, y . So there is no complement of an odd cycle in $G|N$, and therefore the complement of $G|N$ is bipartite and N is the union of two completes.

Let M_1 be the vertices in M adjacent to x , M_2 those adjacent to y , then $M_1 \cup M_2 = M$ and $M_1 \cap M_2 = \emptyset$.

If there exist m_1 in M_1 and m_2 in M_2 such that m_1 is adjacent to m_2 , then the graph induced by $\{m_1, m_2, v, x, y, u\}$ is 3-sun, a contradiction. So there are no edges between M_1 and M_2 , M_1 is anticomplete to y , and M_2 is anticomplete to x . Since $\{v, m, m', y\}$ is not a claw for m, m' in M_1 , it follows that M_1 is a complete, and the same holds for M_2 .

Case 1: M_1 and M_2 are both non-empty. Since A contains no stable set of size three (for otherwise there would be a claw in G), every vertex in N is complete to one of M_1, M_2 . Let N_3 be the vertices complete to $M_1 \cup M_2, N_1$

the vertices of $N \setminus N_3$ complete to M_1 , and N_2 the vertices of $N \setminus N_3$ complete to M_2 . So $x \in N_1$ and $y \in N_2$. Since $\{m, n, n', u\}$ is not a claw for m in M_1 and n, n' in $N_1 \cup N_3$, it follows that $N_1 \cup N_3$ is a complete. Similarly, $N_2 \cup N_3$ is a complete. Suppose N_3 is non-empty, and choose $n \in N_3$. Then n is complete to $(A \cup \{v\}) \setminus \{n\}$, and therefore is singular (for its non-neighbors are a subset of B); and by the choice of v, n and v are twins. Since G admits no twins, it follows that N_3 is empty. Suppose some n_1 in N_1 is adjacent to n_2 in N_2 . Choose m'_1 in M_1 non-adjacent to n_2 and m'_2 in M_2 non-adjacent to n_1 . Then $m'_1 n_1 n_2 m'_2 u m'_1$ is a hole of length 5, a contradiction. So N_1 is anticomplete to N_2 . Suppose n_1 in N_1 has a neighbor m'_2 in M_2 . Then $\{m'_2, n_1, y, u\}$ is a claw, a contradiction. So N_1 is anticomplete to M_2 , and, similarly, N_2 is anticomplete to M_1 .

For $i = 1, 2$ choose m'_i in M_i , and assume that m'_i has a non-neighbor b_i in B . If m'_1 and m'_2 have a common non-neighbor $b \in B$, then $\{u, m'_1, m'_2, b\}$ is a claw, a contradiction. So there are two vertices b_1 and b_2 in B such that b_1 is non-adjacent to m'_1 and adjacent to m'_2 , and b_2 is non-adjacent to m'_2 and adjacent to m'_1 . But then $m'_1 b_2 b_1 m'_2 v m'_1$ is a hole of length 5, again a contradiction. So, exchanging M_1 and M_2 if necessary, we may assume that M_1 is complete to B , and, since G admits no twins, $|M_1| = 1$, say $M_1 = \{m_1\}$.

Let b be a vertex of B with a neighbor n_1 in N_1 . We claim that b is complete to M_2 and anticomplete to N_2 . If b has a non-neighbor m_2 in M_2 , then $n_1 b u m_2 v n_1$ is a hole of length 5; and, if b has a neighbor n_2 in N_2 , then $\{b, n_1, n_2, u\}$ is a claw; in both cases there is a contradiction. This proves the claim.

So every vertex of B is either anticomplete to N_1 , or complete to M_2 and anticomplete to N_2 . Let B_1 be the set of vertices of B with a neighbor in N_1 . Then (B_1, N_1) is a non-dominating homogeneous pair, and, since G does not admit a non-dominating W -join or twins, it follows that $|B_1| \leq 1$ and $|N_1| = 1$, say $N_1 = \{x\}$.

Assume that B_1 is non-empty, let $B_1 = \{b_1\}$. Let $B_2 = B \setminus B_1$. We claim that in this case B_2 is complete to M_2 . If b_2 in B_2 has a non-neighbor m_2 in M_2 , then $b_2 \neq b_1$ and $\{b_1, x, m_2, b_2\}$ is a claw, a contradiction. This proves the claim. But now the vertices of M_2 are all twins, and, since G does not admit twins, $|M_2| = 1$. Moreover, (B_2, N_2) is a non-dominating homogeneous pair, and, since G does not admit a non-dominating W -join or twins, it follows that $|B_2| = |N_2| = 1$, so $B_2 = \{u\}$ and $N_2 = \{y\}$. But now every clique of G contains either v or b_1 , and hence $K(G)$ is the complement of a bipartite graph, and therefore perfect. This finishes the case when B_1 is non-empty.

If B_1 is empty, $(B, M_2 \cup N_2)$ is a non-dominating homogeneous pair, and, since G does not admit a non-dominating W -join or twins, it follows that $|B| = |M_2 \cup N_2| = 1$, a contradiction because both M_2 and N_2 are non-empty. This finishes the case when both M_1 and M_2 are non-empty.

Case 2: One of M_1, M_2 is empty. We may assume that M_2 is empty, and so M is complete to x and anticomplete to y . Let N_1 be the set of vertices in N complete to M , N_2 the set of vertices in N that are anticomplete to M , and let $N_3 = N \setminus (N_1 \cup N_2)$.

We claim that $N_1 \cup N_3$ and $N_2 \cup N_3$ are both completes. Choose two different vertices n_3 in $N_3 \cup N_1$ and n_1 in N_1 , and let m be a neighbor of n_3 in M . Since $\{m, u, n_1, n_3\}$ is not a claw, n_1 is adjacent to n_3 , and therefore N_1 is a complete and N_1 is complete to N_3 . Next, choose two different vertices n_3 in $N_3 \cup N_2$ and n_2 in N_2 , and let m be a non-neighbor of n_3 in M . Since $\{v, m, n_2, n_3\}$ is not a claw, n_2 is adjacent to n_3 , and therefore N_2 is a complete and N_2 is complete to N_3 . Finally, suppose there exist two non-adjacent vertices n_3 and n'_3 in N_3 . Since $\{m, u, n_3, n'_3\}$ is not a claw for any $m \in M$, it follows that no vertex of M is adjacent to both n_3 and n'_3 . Let m be a neighbor of n_3 in M and m' be a neighbor of n'_3 in M . Then m is non-adjacent to n'_3 and m' is non-adjacent to n_3 , and the graph induced by $\{v, m, m', u, n_3, n'_3\}$ is a 3-sun, a contradiction. So N_3 is a complete. This proves the claim. Since there exist two non-adjacent vertices in N , both N_1 and N_2 are non-empty.

35.2. Let b in B be adjacent to n_3 in N_3 and to m in M . Then n_3 is non-adjacent to m .

Suppose they are adjacent. Let m' be a non-neighbor of n_3 in M , and let n_2 be in N_2 . Then $n_3 m v$ is a triangle, b is adjacent to n_3, m, n_2 is adjacent to v and n_3, m' is adjacent to v and m , and this is a 0-, 1-, or 2-pyramid, a contradiction. This proves 35.2.

35.3. Every vertex in N_1 has a non-neighbor in N_2 .

Suppose some vertex n_1 of N_1 is complete to N_2 . Then the set of non-neighbors of n_1 is included in B , and therefore n_1 is singular, and it is complete to $A \setminus \{n_1\}$. From the choice of v, n_1 has no neighbor in B , but now n_1 and v are twins, a contradiction. This proves 35.3.

35.4. M is complete to B .

Let B_1 be the set of vertices in B that are complete to M . Suppose there exists b_2 in $B \setminus B_1$, and let m be a non-neighbor of b_2 in M .

35.4.1. $|N_2| = 1$ and N_2 is anticomplete to B .

Let n be in N_2 . Since nb_2umvn is not a hole of length 5, it follows that n is non-adjacent to b_2 , and the same holds for every vertex of $B \setminus B_1$. So n is anticomplete to $B \setminus B_1$. Since $\{b_1, b_2, m, n\}$ is not a claw for $b_1 \in B_1$, it follows that n is anticomplete to B_1 , and the same holds for every vertex of N_2 . Therefore N_2 is anticomplete to B . But now $\{v\} \cup N_1 \cup N_3$ is a clique cutset separating N_2 from $M \cup B$. By Theorem 12, G is either a linear interval graph or G admits twins, or a 0-join, or a 1-join, or a coherent W -join, or it is not an internal clique cutset; and it follows from the hypotheses of the theorem and from Proposition 28 that we may assume that the last alternative holds, and $|N_2| = 1$, say $N_2 = \{n_2\}$. This proves 35.4.1.

35.4.2. B is anticomplete to N_3 .

Suppose a vertex $b \in B$ has a neighbor $n \in N_3$. By the definition of N_3 , n has a neighbor m' in M . By 35.2, m' is non-adjacent to b . But now $\{n, n_2, b, m'\}$ is a claw, a contradiction. This proves 35.4.2.

Now $M \cup N_1$ is a clique cutset separating $\{v\} \cup N_2 \cup N_3$ from B . Since $|B| > 1$ and $|\{v\} \cup N_2 \cup N_3| > 1$, it follows from Theorem 12 that G is a linear interval graph, and therefore $K(G)$ is perfect by Proposition 28. This completes the proof of 35.4.

By 35.4, for every non-singular vertex in B , the set of its neighbors in A is complete to B .

35.5. B is anticomplete to N_3 .

Suppose some vertex b in B has a neighbor n_3 in N_3 . By the definition of N_3 , n_3 has a neighbor in M , and this contradicts 35.2. This proves 35.5.

35.6. N_3 is empty and $|M| = 1$.

If N_3 is non-empty then $|M| > 1$ and (N_3, M) is a coherent homogeneous pair. So N_3 is empty, but now the vertices of M are twins, so $|M| = 1$. This proves 35.6.

It follows from 35.6 that every non-singular vertex in B has at most one neighbor in A , and, since M is complete to B and has size one, every non-singular vertex in B is complete to M and anticomplete to $A \setminus M$. Therefore, the vertices of U are all twins, and, since G admits no twins, $U = \{u\}$. Let $B_2 = B \setminus U$.

35.7. B_2 is non-empty.

Otherwise (N_1, N_2) is a coherent homogeneous pair, so each of them has size one and $K(G)$ is a three-edge path. This proves 35.7.

35.8. If n_1 in N_1 is non-adjacent to n_2 in N_2 , then every b in B_2 is adjacent to exactly one of n_1, n_2 .

Let b_2 in B_2 . Since b_2 in B_2 is singular, b_2 is adjacent to at least one of n_1, n_2 . Since $\{b_2, n_1, n_2, u\}$ is not a claw, b_2 is non-adjacent to at least one of n_1, n_2 . This proves 35.8.

35.9. No vertex of N_1 has a neighbor and a non-neighbor in B_2 .

Suppose n_1 in N_1 has a neighbor b_1 in B_2 and a non-neighbor b_2 in B_2 . By 35.3 n_1 has a non-neighbor n_2 in N_2 . By 35.8 n_2 is adjacent to b_2 and not to b_1 . But now $b_1n_1vn_2b_2b_1$ is a hole of length 5, a contradiction. This proves 35.9.

Let N_{11} be the vertices of N_1 complete to B_2 , $N_{12} = N_1 \setminus N_{11}$. So N_{12} is anticomplete to B . It follows from 35.8 that every vertex of N_2 is either complete to N_{11} or to N_{12} . Let N_{22} be the set of vertices in N_2 with a non-neighbor in N_{11} .

Then N_{22} is complete to N_{12} . Let N_{21} be the vertices in N_2 with a non-neighbor in N_{12} . Then N_{21} is complete to N_{11} . Let $N_{23} = N_2 \setminus (N_{21} \cup N_{22})$. So N_{23} is complete to N_1 . By 35.8 B_2 is anticomplete to N_{22} and complete to N_{21} . Now (B_2, N_{23}) is a coherent homogeneous pair, and all the vertices of $N_{11}, N_{12}, N_{22}, N_{21}$ are twins, so all these sets have size at most one.

Now, every clique of G contains either v or b_2 , so $K(G)$ is the complement of a bipartite graph, and hence it is perfect. This completes the proof of Theorem 35. \square

3.2.3. Basic classes

Finally, we show that, if an interesting HCH claw-free graph belongs to one of the basic classes of Theorem 5, then its clique graph is perfect.

Theorem 36. *If G is interesting HCH, antiprismatic, and every vertex of G is in a triad, then $K(G)$ is perfect.*

Proof. We prove that G contains no 4-wheel or 3-fan, and, then, by Theorem 16, $K(G)$ is bipartite.

Suppose G contains a 4-wheel. Let $a_1a_2a_3a_4a_1$ be a hole and let c be adjacent to all a_i . Since every vertex is in a triad, there are two vertices c_1, c_2 different from a_1, a_2, a_3, a_4 such that $\{c, c_1, c_2\}$ is a stable set. Since G is antiprismatic, every other vertex in G is adjacent exactly to two of $\{c, c_1, c_2\}$. In particular, each a_i is adjacent either to c_1 or to c_2 . If two consecutive vertices of the hole, for instance, a_1, a_2 , are adjacent to the same c_j , then $\{a_1, a_3, a_2, a_4, c, c_j\}$ induces a 1-, 2-, or 3-pyramid, a contradiction because G is HCH. So, without loss of generality, we may assume that a_1 and a_3 are adjacent to c_1 and not to c_2 , while a_2 and a_4 are adjacent to c_2 , and not to c_1 . But then $\{a_1, a_2, a_3, c_2\}$ induces a claw, a contradiction. This proves that G does not contain a 4-wheel.

Suppose now that G contains a 3-fan. Let $a_1a_2a_3a_4$ be an induced path and let c be adjacent to all a_i . Since every vertex is in a triad, there are two vertices c_1, c_2 different from a_1, a_2, a_3, a_4 such that $\{c, c_1, c_2\}$ is a stable set. Since G is antiprismatic, each a_i is adjacent either to c_1 or to c_2 . If a_2 and a_3 are adjacent to the same c_j , then $\{a_1, a_3, a_2, a_4, c, c_j\}$ induces a 0-, 1-, or 2-pyramid, a contradiction because G is HCH. So, without loss of generality, we may assume that a_2 is adjacent to c_1 and not c_2 , while a_3 is adjacent to c_2 and not c_1 . Since $\{a_3, a_2, c_2, a_4\}$ is not a claw, a_4 is adjacent to c_2 , and, analogously, a_1 is adjacent to c_1 . By the same argument applied to the 3-fan induced by the path $a_2ca_4c_2$ and the vertex a_3 , there is a vertex d adjacent to a_4 and c_2 but not adjacent to a_2, c , or a_3 , and so $d \notin \{a_1, a_2, a_3, a_4, c, c_1, c_2\}$ (see Fig. 14).

Since $c_1a_2a_4dc_1$ is not a hole of length 5, d is non-adjacent to c_1 . Thus, c_1, c , and d form a triad, but the vertex c_2 is adjacent only to one of them, a contradiction because G is antiprismatic. This concludes the proof of Theorem 36. \square

Theorem 37. *Let $G \in \mathcal{S}_6$ be a connected interesting HCH graph such that every vertex of G is in a triad. Then $K(G)$ is perfect.*

Proof. Let A, B , and C be the sets of vertices of the graph H_5 in the definition of the class \mathcal{S}_6 , and let A_G, B_G , and C_G be those sets intersected with $V(G)$. We remind the reader that $a_0 \in A_G$ and $b_0 \in B_G$ by the definition of \mathcal{S}_6 . Every triad in G is of the form $\{a_i, b_j, c_k\}$, since A_G, B_G and C_G are complete sets. Moreover, either $i = j = 0$ or $k = i$ and $j = 0$ or $k = j$ and $i = 0$. Since every vertex of G is in a triad, it follows that A_G, B_G , and C_G are non-empty and if $i \neq 0$ and $a_i \in A_G$, then $c_i \in C_G$. Analogously, if $i \neq 0$ and $b_i \in A_G$, then $c_i \in C_G$. Let $I_A = \{i > 0 : a_i \in A_G\}$, $I_B = \{i > 0 : b_i \in B_G\}$, and $I_C = \{i > 0 : c_i \in C_G\}$. Then $I_A \cup I_B \subseteq I_C$.




Fig. 14. Situation for the second part of the proof of Theorem 36.




Fig. 15. Last three cases for the proof of Theorem 37.

Assume first that $I_C \setminus (I_A \cup I_B)$ is non-empty. Since the set $C' = \{c_i : i \in C \setminus (I_A \cup I_B)\}$ is complete to $V(G) \setminus (C' \cup \{a_0, b_0\})$, and the only cliques containing a_0 or b_0 are A_G and B_G , respectively, it follows that every pair of cliques of G , except for the pair A_G, B_G , has non-empty intersection. Thus $K(G)$ is a split graph, hence perfect.

So we may assume that $I_A \cup I_B = I_C$. If $|I_A \cup I_B| \geq 3$, we may assume by switching A and B if necessary that $1, 2 \in I_A$ and $3 \in I_C$, and then the graph induced by $\{a_1, a_2, c_1, c_2, c_3, a_0\}$ is a 1-pyramid, a contradiction because G is HCH. On the other hand, since G is connected, both I_A and I_B are non-empty and $|I_A \cup I_B| \geq 2$. So, without loss of generality, we consider three cases: $I_A = I_B = \{1, 2\}$, $I_A = \{1, 2\}$ and $I_B = \{2\}$, $I_A = \{1\}$ and $I_B = \{2\}$. Graphs obtained in each case are depicted in Fig. 15, with their corresponding clique graphs, which are all perfect. That concludes this proof. \square

3.2.4. Proof of Theorem 19

First we prove that the clique graph of an interesting HCH claw-free graph is perfect.

Proof of Theorem 26. Let G be an interesting HCH claw-free graph. The proof is by induction on $|V(G)|$, using the decomposition of Theorem 5. Assume that for every smaller interesting HCH claw-free G' , $K(G')$ is perfect. We show that $K(G)$ is perfect.

If G admits twins, then $K(G)$ is perfect by Lemma 14, and, if G is not connected, then $K(G)$ is perfect by Lemma 15. If G is connected, admits a 1-join and no twins, then $K(G)$ is perfect by Theorems 29 and 7. If G admits no twins, 0-, or 1-joins, but admits a 2-join, then $K(G)$ is perfect by Theorem 30. If G admits a coherent or non-dominating W -join and no twins, then $K(G)$ is perfect by Theorem 32. If G contains a singular vertex, then $K(G)$ is perfect by Theorems 34 and 35. So we may assume not. If G admits a hex-join and no twins, then by Theorem 33 $G = K(G) = C_6$, and therefore $K(G)$ is perfect.

So we may assume that G admits none of the decompositions of the previous paragraph, and, by Theorem 5, G is antiprismatic, or belongs to $\mathcal{S}_0 \cup \dots \cup \mathcal{S}_6$.

If $G \in \mathcal{S}_0$, then $K(G)$ is perfect by Theorem 23. The graphs $icosa(-2)$, $icosa(-1)$, and $icosa(0)$ contain holes of length 5, and therefore are not interesting, so $G \notin \mathcal{S}_1$. $G \notin \mathcal{S}_2$, because vertices v_3, v_4, v_5, v_6, v_9 induce a hole of length 5 in H_1 (Fig. 5). If $G \in \mathcal{S}_3$, then, by Proposition 28, $K(G)$ is perfect. If $G \in \mathcal{S}_4$ then, since G does not contain a singular vertex, G is a line graph and $K(G)$ is perfect by Theorem 23. $G \notin \mathcal{S}_5$, because the vertex d_1 in the definition of the class \mathcal{S}_5 is singular. If $G \in \mathcal{S}_6$, then $K(G)$ is perfect by Theorem 37, and, finally, if G is antiprismatic, then $K(G)$ is perfect by Theorem 36. This completes the proof of Theorem 26. \square

Now, Theorem 19 is an immediate corollary of the following:

Theorem 38. *Let G be claw-free and assume that G is HCH. Then the following are equivalent:*

- (i) *No induced subgraph of G is an odd hole, or $\overline{C_7}$.*
- (ii) *G is clique-perfect.*
- (iii) *G is perfect.*

Table 1
Forbidden induced subgraphs for clique-perfect graphs in each studied class

Graph classes	Forbidden subgraphs	Reference
HCH claw-free graphs	Odd holes $\overline{C_7}$	Theorem 19
Line graphs	Odd holes 3-sun	Theorem 18

Proof. The equivalence between (i) and (iii) is a corollary of Theorem 1, because by Proposition 25 HCH graphs contain no antiholes of length at least 8. From Theorem 3 it follows that (ii) implies (i). Finally, by Theorem 26 and Propositions 21 and 25, we deduce that (i) implies (ii), and this completes the proof. \square

The recognition of clique-perfect HCH claw-free graphs can be reduced to the recognition of perfect graphs, which is solvable in polynomial time [8].

4. Summary

These results allow us to formulate partial characterizations of clique-perfect graphs by forbidden subgraphs, as is shown in Table 1.

Note that in both cases all the forbidden induced subgraphs are minimal.

Acknowledgment

We are grateful to both anonymous referees for their careful reading of the paper and for many helpful remarks. In particular, one of them pointed out to us the linear time recognition algorithm for clique-perfect line graphs, and suggested ideas that allowed us to simplify some of our original proofs.

References

- [1] V. Balachandhran, P. Nagavamsi, C. Pandu Rangan, Clique-transversal and clique-independence on comparability graphs, *Inform. Process. Lett.* 58 (1996) 181–184.
- [2] C. Berge, *Graphs and Hypergraphs*, North-Holland, Amsterdam, 1985.
- [3] C. Berge, M. Las Vergnas, Sur un théorème du type König pour hypergraphes, *Ann. New York Acad. Sci.* 175 (1970) 32–40.
- [4] F. Bonomo, G. Durán, M. Groshaus, J. Szwarcfiter, On clique-perfect and K -perfect graphs, *Ars Combin.* 80 (2006) 97–112.
- [5] F. Bonomo, G. Durán, M. Lin, J. Szwarcfiter, On balanced graphs, *Math. Programming Ser. B* 105 (2006) 233–250.
- [6] A. Brandstädt, V. Chepoi, F. Dragan, Clique r -domination and clique r -packing problems on dually chordal graphs, *SIAM J. Discrete Math.* 10 (1997) 109–127.
- [7] M. Chang, M. Farber, Zs. Tuza, Algorithmic aspects of neighbourhood numbers, *SIAM J. Discrete Math.* 6 (1993) 24–29.
- [8] M. Chudnovsky, G. Cornuéjols, X. Liu, P. Seymour, K. Vušković, Recognizing Berge graphs, *Combinatorica* 25 (2005) 143–187.
- [9] M. Chudnovsky, N. Robertson, P. Seymour, R. Thomas, The strong perfect graph theorem, *Ann. of Math.* 164 (2006) 51–229.
- [10] M. Chudnovsky, P. Seymour, Claw-free graphs. III. Circular interval graphs, submitted for publication.
- [11] M. Chudnovsky, P. Seymour, Claw-free graphs. IV. Decomposition theorem, *J. Combin. Theory Ser. B*, to appear.
- [12] V. Chvátal, N. Sbihi, Bull-free berge graphs are perfect, *Graphs Combin.* 3 (1987) 127–139.
- [13] M. Conforti, G. Cornuéjols, R. Rao, Decomposition of balanced matrices, *J. Combin. Theory Ser. B* 77 (1999) 292–406.
- [14] G. Durán, M. Lin, J. Szwarcfiter, On clique-transversal and clique-independent sets, *Ann. Oper. Res.* 116 (2002) 71–77.
- [15] F. Escalante, Über iterierte clique-graphen, *Abh. Math. Sem. Univ. Hamburg* 39 (1973) 59–68.
- [16] M. Golumbic, *Algorithmic Graph Theory and Perfect Graphs*, Academic Press, New York, 1980.
- [17] M. Grötschel, L. Lovász, A. Schrijver, The ellipsoid method and its consequences in combinatorial optimization, *Combinatorica* 1 (1981) 169–197.
- [18] V. Guruswami, C. Pandu Rangan, Algorithmic aspects of clique-transversal and clique-independent sets, *Discrete Appl. Math.* 100 (2000) 183–202.
- [19] J. Lehel, Zs. Tuza, Neighborhood perfect graphs, *Discrete Math.* 61 (1986) 93–101.
- [20] P. Lehot, An optimal algorithm to detect a line graph and output its root graph, *J. ACM* 21 (4) (1974) 569–575.
- [21] F. Maffray, Kernels in perfect line-graphs, *J. Combin. Theory Ser. B* 55 (1992) 1–8.
- [22] F. Maffray, B. Reed, A description of claw-free perfect graphs, *J. Combin. Theory Ser. B* 75 (1) (1999) 134–156.

- [23] E. Prisner, Hereditary clique-Helly graphs, *J. Combin. Math. Combin. Comput.* 14 (1993) 216–220.
- [24] F. Protti, J. Szwarcfiter, Clique-inverse graphs of bipartite graphs, *J. Combin. Math. Combin. Comput.* 40 (2002) 193–203.
- [25] D. Rose, R. Tarjan, G. Lueker, Algorithmic aspects of vertex elimination on graphs, *SIAM J. Comput.* 5 (1976) 266–283.
- [26] L. Trotter, Line perfect graphs, *Math. Programming* 12 (1977) 255–259.
- [27] A. Tucker, A reduction procedure for coloring perfect K_4 -free graphs, *J. Combin. Theory Ser. B* 43 (1987) 151–172.