

QUASILINEAR ELLIPTIC SYSTEMS OF RESONANT TYPE AND NONLINEAR EIGENVALUE PROBLEMS

PABLO L. DE NÁPOLI AND M. CRISTINA MARIANI

Received 20 February 2001

This work is devoted to the study of a quasilinear elliptic system of resonant type. We prove the existence of infinitely many solutions of a related nonlinear eigenvalue problem. Applying an abstract minimax theorem, we obtain a solution of the quasilinear system $-\Delta_p u = F_u(x, u, v)$, $-\Delta_q v = F_v(x, u, v)$, under conditions involving the first and the second eigenvalues.

1. Introduction

1.1. The problem and some previous results. We consider a gradient elliptic system

$$-\Delta_p u = F_u(x, u, v), \quad -\Delta_q v = F_v(x, u, v). \quad (1.1)$$

Elliptic problems involving the p -Laplacian have been studied by several authors (cf. [3, 7, 8, 10, 11]). We recall some results from the work of Boccardo and de Figueiredo [4].

It is well known that the solutions of (1.1) in $W = W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega)$ are the critical points of the functional

$$\Phi(u, v) = \frac{1}{p} \int_{\Omega} |\nabla u|^p + \frac{1}{q} \int_{\Omega} |\nabla v|^q - \int_{\Omega} F(x, u, v) \quad (1.2)$$

under the following three assumptions:

- (1) $\Omega \subset \mathbb{R}^N$ is a bounded domain, $1 < p, q < N$, so that the following continuous embeddings hold:

$$W_0^{1,p}(\Omega) \subset L^{p^*}(\Omega), \quad W_0^{1,q}(\Omega) \subset L^{q^*}(\Omega); \quad (1.3)$$

(2) $F : \bar{\Omega} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is C^1 and verifies the following growth condition:

$$|F(x, s, t)| \leq c(1 + |s|^{p^*} + |t|^{q^*}) \quad \forall x \in \bar{\Omega}; s, t \in \mathbb{R}; \tag{1.4}$$

(3) in order to have $\Phi \in C^1(W, \mathbb{R})$, we assume

$$\begin{aligned} |F_s(x, s, t)| &\leq C(1 + |s|^{p^*-1} + |t|^{q^*(p^*-1)/p^*}) \quad \forall x \in \bar{\Omega}; s, t \in \mathbb{R}, \\ |F_t(x, s, t)| &\leq C(1 + |t|^{q^*-1} + |s|^{p^*(q^*-1)/q^*}) \quad \forall x \in \bar{\Omega}; s, t \in \mathbb{R}. \end{aligned} \tag{1.5}$$

The geometry of Φ depends strongly on the values of α and β in the estimate

$$|F(x, s, t)| \leq c(1 + |s|^\alpha + |t|^\beta) \quad \forall x \in \bar{\Omega}; s, t \in \mathbb{R}, \tag{1.6}$$

where $\alpha \leq p^*, \beta \leq q^*$. In this work we are interested in the case $\alpha = p, \beta = q$ (systems of resonant type).

In our case, it is quite adequate to assume the following condition on F : consider the function

$$L(x, s, t) = \frac{1}{p}F_s(x, s, t)s + \frac{1}{q}F_t(x, s, t)t - F(x, s, t). \tag{1.7}$$

Assume that

$$\lim_{\|(s,t)\| \rightarrow \infty} L(x, s, t) = \pm\infty \quad \text{uniformly for } x \in \Omega. \tag{1.8}$$

This assumption implies that Φ satisfies the following compactness Cerami condition.

Definition 1.1. Let X be a Banach space and $\Phi \in C^1(X, \mathbb{R})$. Given $c \in \mathbb{R}$, we say that Φ satisfies condition (C_c) , if

- (1) any bounded sequence $(u_n) \subset X$ such that $\Phi(u_n) \rightarrow c$ and $\Phi'(u_n) \rightarrow 0$ has a convergent subsequence;
- (2) there exist constants $\delta, R, \alpha > 0$ such that

$$\|\Phi'(u)\| \|u\| \geq \alpha \quad \forall u \in \Phi^{-1}([c - \delta, c + \delta]) \text{ with } \|u\| \geq R. \tag{1.9}$$

If $\Phi \in C^1(X, \mathbb{R})$ satisfies condition (C_c) for every $c \in \mathbb{R}$, we say that Φ satisfies condition (C) .

Condition (C) was introduced by Cerami [5]. It was shown in [2] that from condition (C) it is possible to obtain a deformation lemma, that is fundamental in order to get minimax theorems.

In order to avoid resonance, Boccardo and de Figueiredo [4] introduced an assumption on F involving an eigenvalue problem

$$\begin{aligned} -\Delta_p u - aG_u(u, v) &= \lambda|u|^{p-2}u, \\ -\Delta_q v - aG_v(u, v) &= \lambda|v|^{q-2}v, \end{aligned} \tag{1.10}$$

where $a = a(x) \in L^\infty(\Omega)$ and G is a C^1 even function $G : \mathbb{R} \rightarrow [0, \infty)$ such that

$$G(c^{1/p}s, c^{1/q}t) = cG(s, t) \quad \forall c > 0, \tag{1.11}$$

$$G(s, t) \leq K \left(\frac{1}{p}|s|^p + \frac{1}{q}|t|^q \right). \tag{1.12}$$

We call such a G a (p, q) homogeneous function.

It is easy to see that (1.11) implies (1.12). A (p, q) -homogeneous function satisfies

$$\frac{1}{p}G_s(s, t)s + \frac{1}{q}G_t(s, t)t = G(s, t). \tag{1.13}$$

Examples of (p, q) homogeneous functions are

- (1) $G(s, t) = c_1|s|^p + c_2|t|^q$,
- (2) $G(s, t) = c|s|^\alpha|t|^\beta$ with $\alpha/p + \beta/q = 1$ where c, c_1, c_2 are constants.

The following results are proved in [4].

THEOREM 1.2. *Problem (1.10), with G as above, has a first eigenvalue $\lambda_1(a)$, characterized variationally by*

$$\lambda_1(a) = \inf_{(u,v) \neq (0,0)} \frac{(1/p) \int_\Omega |\nabla u|^p + (1/q) \int_\Omega |\nabla v|^q - \int_\Omega aG(u, v)}{(1/p) \int_\Omega |u|^p + (1/q) \int_\Omega |v|^q} \tag{1.14}$$

which depends continuously on a in the L^∞ -norm.

THEOREM 1.3. *Assume (1.5), (1.6) with $\alpha = p, \beta = q$, and that the following conditions hold:*

- (1) *there exist positive numbers c, R, μ , and ν such that*

$$\frac{1}{p}sF_s(x, s, t) + \frac{1}{q}tF_t(x, s, t) - F(x, s, t) \geq c(|s|^\mu + |t|^\nu) \quad \text{for } |s|, |t| > R, \tag{1.15}$$

- (2) *there exists G as above, such that*

$$\limsup_{|s|, |t| \rightarrow \infty} \frac{F(x, s, t)}{G(s, t)} \leq a(x) \in L^\infty(\Omega), \tag{1.16}$$

where $\lambda_1(a) > 0$.

Then the functional Φ is bounded from below and the infimum is achieved.

1.2. The existence of infinitely many eigenfunctions. Let \mathcal{C} be the class of compact symmetric ($C = -C$) subsets of the space W . We recall that for $C \in \mathcal{C}$ the Krasnoselskii genus $\text{gen}(C)$ is defined as the minimum integer n such that there exists an odd continuous mapping $\varphi : C \rightarrow (\mathbb{R}^n - \{0\})$ (cf. [1]). We note

$$\mathcal{C}_k = \{C \in \mathcal{C} : \text{gen}(C) \geq k\}. \tag{1.17}$$

For an arbitrary symmetric subset S of $W - \{0\}$ the genus over compact sets $\gamma(S)$ is defined by

$$\gamma(S) = \sup \{ \text{gen}(C) : C \subset S, C \in \mathcal{C}, C \text{ compact} \}. \quad (1.18)$$

Now we may state our main result on the eigenvalue problem.

THEOREM 1.4. *The eigenvalue problem (1.10), with G as above, has infinitely many eigenfunctions given by*

$$\lambda_k(a, G) = \inf_{C \in \mathcal{C}_k} \sup_{(u,v) \in C} \frac{(1/p) \int_{\Omega} |\nabla u|^p + (1/q) \int_{\Omega} |\nabla v|^q - \int_{\Omega} aG(u, v)}{(1/p) \int_{\Omega} |u|^p + (1/q) \int_{\Omega} |v|^q} \quad (1.19)$$

and $\lambda_k(a, G) \rightarrow \infty$ as $k \rightarrow \infty$. Moreover, λ_k depends continuously on a in the L^∞ -norm.

Remark 1.5. Equivalently if we define

$$S = \left\{ (u, v) \in W : \frac{1}{p} \int_{\Omega} |u|^p + \frac{1}{q} \int_{\Omega} |v|^q = 1 \right\}, \quad (1.20)$$

we have

$$\lambda_k(a, G) = \inf_{C \in \mathcal{C}_k, C \subset S} \sup_{(u,v) \in C} \frac{1}{p} \int_{\Omega} |\nabla u|^p + \frac{1}{q} \int_{\Omega} |\nabla v|^q - \int_{\Omega} aG(u, v). \quad (1.21)$$

We will write $\lambda_k(a)$ instead of $\lambda_k(a, G)$, when the dependence on the (p, q) -homogeneous function G is clear from the context.

1.3. The existence result for resonant systems. Applying [Theorem 1.4](#) and an abstract minimax principle from [\[9\]](#), we prove the following theorem.

THEOREM 1.6. *Assume that $F : \Omega \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ verifies (1.5), (1.6) with $\alpha = p$, $\beta = q$, (1.8), and that $a_1, a_2 \in L^\infty(\Omega)$ satisfy*

$$a_1(x) \leq \liminf_{|s|, |t| \rightarrow \infty} \frac{F(x, s, t)}{G_1(s, t)} \leq \limsup_{|s|, |t| \rightarrow \infty} \frac{F(x, s, t)}{G_2(s, t)} \leq a_2(x) \quad (1.22)$$

with G_1 and G_2 two (p, q) -homogeneous functions and $\lambda_1(a_1, G_1) < 0 < \lambda_2(a_2, G_2)$, where $\lambda_1(a_1, G_1), \lambda_2(a_2, G_2)$ are given by (1.19). Then problem (1.1) has at least one solution.

Remark 1.7. The conditions above could be reformulated in terms of a different eigenvalue problem, for $a \in L^\infty(\Omega), a(x) > 0$

$$-\Delta_p u = \mu a G_u(u, v), \quad -\Delta_q v = \mu a G_v(u, v). \quad (1.23)$$

This problem also has infinitely many eigenvalues given by

$$\mu_k(a) = \inf_{C \in \mathcal{C}_k} \sup_{(u,v) \in C} \frac{(1/p) \int_{\Omega} |\nabla u|^p + (1/q) \int_{\Omega} |\nabla v|^p}{\int_{\Omega} aG(u, v)}. \quad (1.24)$$

The condition $\lambda_1(a) < 0$ is equivalent to $\mu_1(a) < 1$, and the condition $\lambda_2(a) > 0$ is equivalent to $\mu_2(a) > 1$.

Remark 1.8. As an example for [Theorem 1.6](#), we may take

$$G_1(s, t) = G_2(s, t) = |s|^\alpha |t|^\beta \tag{1.25}$$

with $\alpha/p + \beta/q = 1$;

$$F(x, s, t) = \lambda |s|^\alpha |t|^\beta + c |s|^\mu |t|^\delta, \tag{1.26}$$

where $c \neq 0$ is a constant, and we assume that

$$\mu_1(1) < \lambda < \mu_2(1) \tag{1.27}$$

and $\mu < \alpha, \delta < \beta$, where $\mu_1(1), \mu_2(1)$ are defined as above (with $a \equiv 1$).

2. The eigenvalue problem

2.1. The functional framework. We apply the following abstract theorem due to Amann [1].

THEOREM 2.1. *Suppose that the following hypotheses are satisfied:*

- X is a real Banach space of infinite dimension, that is uniformly convex;
- $A : X \rightarrow X^*$ is an odd potential operator (i.e., A is the Gateaux derivative of $\mathcal{A} : X \rightarrow \mathbb{R}$) which is uniformly continuous on bounded sets, and satisfies condition (S)₁: if $u_j \rightharpoonup u$ (weakly in X) and $A(u_j) \rightarrow v$, then $u_j \rightarrow u$ (strongly in X).
- For a given constant $\alpha > 0$, the level set

$$M_\alpha = \{u \in X : \mathcal{A}(u) = \alpha\} \tag{2.1}$$

is bounded and each ray through the origin intersects M_α . Moreover, for every $u \neq 0$, $\langle A(u), u \rangle > 0$ and there exists a constant $\rho_\alpha > 0$ such that $\langle A(u), u \rangle \geq \rho_\alpha$ on M_α .

- The mapping $B : X \rightarrow X^*$ is a strongly sequentially continuous odd potential operator (with potential \mathcal{B}), such that $\mathcal{B}(u) \neq 0$ implies that $B(u) \neq 0$.

Let

$$\beta_k = \sup_{C \in \mathcal{C}, C \subset M_\alpha} \inf_{u \in C} \mathcal{B}(u). \tag{2.2}$$

Then if $\beta_k > 0$, there exists an eigenfunction $u_k \in M_\alpha$ with $\mathcal{B}(u) = \beta_k$. If

$$\gamma(\{u \in M_\alpha : \mathcal{B}(u) \neq 0\}) = \infty, \tag{2.3}$$

then there exist infinitely many eigenfunctions.

We will work in the Banach space

$$W = W_0^{1,p}(\Omega) \times W_0^{1,q}(\Omega) \tag{2.4}$$

equipped with the norm

$$\|(u, v)\|_W = \sqrt{\|u\|_p^2 + \|v\|_q^2}. \tag{2.5}$$

As each factor is uniformly convex, we can conclude that W is uniformly convex (see [6]). Given $(u^*, v^*) \in W^{-1,p'}(\Omega) \oplus W^{-1,q'}(\Omega)$ we may think of it as an element of W^* :

$$\langle (u^*, v^*), (u, v) \rangle = \langle u^*, u \rangle + \langle v^*, v \rangle. \tag{2.6}$$

Then we have $W^* \cong W^{-1,p'}(\Omega) \oplus W^{-1,q'}(\Omega)$ (isometric isomorphism), where the norm in W^* is given by

$$\|(u^*, v^*)\|_{W^*} = \sqrt{\|u^*\|^2 + \|v^*\|^2}. \tag{2.7}$$

With the notations of [Theorem 2.1](#), we define

$$\mathcal{A}_0(u, v) = \frac{1}{p} \int_{\Omega} |\nabla u|^p + \frac{1}{q} \int_{\Omega} |\nabla v|^q, \tag{2.8}$$

$$\mathcal{A}(u, v) = \mathcal{A}_0(u, v) - \int_{\Omega} aG(u, v) + M \left(\frac{1}{p} \int_{\Omega} |u|^p + \frac{1}{q} \int_{\Omega} |v|^q \right), \tag{2.9}$$

with a and G as in the statement of [Theorem 1.4](#), and M a fixed constant such that $M > K \|a\|_{L^\infty}$, where K is the constant in (1.12).

We write \mathcal{A}_a instead of \mathcal{A} when we want to remark the dependence on the weight a

$$\begin{aligned} A(u, v) &= (-\Delta_p u - aG_u(u, v) + M|u|^{p-2}u, -\Delta_q v - aG_v(u, v) + M|v|^{q-2}v), \\ \mathcal{B}(u, v) &= \frac{1}{p} \int_{\Omega} |u|^p + \frac{1}{q} \int_{\Omega} |v|^q, \\ B(u, v) &= (|u|^{p-2}u, |v|^{q-2}v). \end{aligned} \tag{2.10}$$

In order to apply [Theorem 2.1](#), we prove the following two lemmas.

LEMMA 2.2. (1) A is uniformly continuous on bounded sets.

(2) A verifies the $(S)_1$ condition.

Proof. We write $A = A_1 - A_2$, where

$$\begin{aligned} A_1(u, v) &= (-\Delta_p u, -\Delta_q v), \\ A_2(u, v) &= (aG_u(u, v) - M|u|^{p-2}u, aG_v(u, v) - M|v|^{q-2}v). \end{aligned} \tag{2.11}$$

We claim that $A_2 : W \rightarrow W^*$ verifies that: if $(u_j, v_j) \rightharpoonup (u, v)$ in W , then $A_2(u_j, v_j) \rightarrow A_2(u, v)$ in W^* .

Indeed, if $(u_j, v_j) \rightharpoonup (u, v)$, then

$$(u_j, v_j) \longrightarrow (u, v) \quad \text{in } L^p(\Omega) \times L^q(\Omega) \tag{2.12}$$

and we obtain that

$$\begin{aligned} G_u(u_j, v_j) &\longrightarrow G_u(u, v) \quad \text{in } L^{p'}(\Omega), \\ G_v(u_j, v_j) &\longrightarrow G_v(u, v) \quad \text{in } L^q(\Omega). \end{aligned} \tag{2.13}$$

Hence, $A_2(u_j, v_j) \rightarrow A_2(u, v)$ in W^* .

Let $(u_j, v_j) \rightharpoonup (u, v)$ in W such that

$$A(u_j, v_j) \longrightarrow (z, w). \tag{2.14}$$

Therefore $A_2(u_j, v_j) \rightarrow A_2(u, v)$ and then $A_1(u_j, v_j) \rightarrow (z, w) + A_2(u, v)$. Since A_1 verifies condition $(S)_1$, it follows that $(u_j, v_j) \rightarrow (u, v)$. \square

LEMMA 2.3. (1) *The set $M_\alpha = \{(u, v) \in W : \mathcal{A} = \alpha\}$ is bounded.*

(2) *Every ray $t \cdot (u, v)$ with $(u, v) \neq 0$ intersects M_α .*

(3) *There exists a constant $\rho_\alpha > 0$ such that*

$$\langle A(u, v), (u, v) \rangle \leq \rho_\alpha. \tag{2.15}$$

(4) *Condition (2.3) is satisfied.*

Proof. (1) As we have fixed $M > K\|a\|_{L^\infty}$ on M_α , then

$$\alpha = \mathcal{A}(u, v) \geq \frac{1}{p}|\nabla u|^p + \frac{1}{q}|\nabla v|^q \tag{2.16}$$

and the proof is complete.

(2) Let $f(c) = \mathcal{A}(c(u, v))$, $f(0) = 0$,

$$\begin{aligned} f(c) &= \frac{c^p}{p} \int_\Omega |\nabla u|^p + \frac{c^q}{q} \int_\Omega |\nabla v|^q \\ &\quad - \int_\Omega aG(cu, cv) + M \left(\frac{c^p}{p} \int_\Omega |u|^p + \frac{c^q}{q} \int_\Omega |v|^q \right). \end{aligned} \tag{2.17}$$

From (1.12) and the choice of M , we have

$$f(c) \geq \frac{c^p}{p} \int_\Omega |\nabla u|^p + \frac{c^q}{q} \int_\Omega |\nabla v|^q \longrightarrow +\infty \tag{2.18}$$

as $c \rightarrow \infty$. Since f is continuous, there exists $c \in \mathbb{R}$ such that $f(c) = \alpha$.

(3) We have

$$\begin{aligned} \langle A(u, v), (u, v) \rangle &= \int_{\Omega} |\nabla u|^p + \int_{\Omega} |\nabla v|^q \\ &\quad - \int_{\Omega} a[G_u(u, v)u + G_v(u, v)v] + M \left(\int_{\Omega} |u|^p + \int_{\Omega} |v|^q \right). \end{aligned} \quad (2.19)$$

Then, using (1.13)

$$\langle A(u, v), (u, v) \rangle \geq \min\{p, q\} \mathcal{A}(u, v) = \min\{p, q\} \alpha. \quad (2.20)$$

(4) In order to see that $\gamma(M_\alpha) \geq k$, it is enough to show that M_α contains subsets homeomorphic to the unit sphere in \mathbb{R}^k by an odd homeomorphism. Hence, the proof is completed. \square

2.2. The continuous dependence of $\lambda_k(a)$ on a . In this section we prove that the eigenvalue $\lambda_k(a)$ depends continuously on the weight a in the L^∞ -norm. This result will be used for proving Lemma 3.3.

PROPOSITION 2.4. *The eigenvalue $\lambda_k(a)$ depends continuously on a in the L^∞ -norm.*

Proof. We have

$$|\mathcal{A}_a(u, v) - \mathcal{A}_b(u, v)| \leq K \|a - b\|_{L^\infty} \left(\frac{1}{p} \int_{\Omega} |u|^p + \frac{1}{q} \int_{\Omega} |v|^q \right), \quad (2.21)$$

where K is given by condition (1.12), with $\mathcal{A}_a, \mathcal{A}_b$ as above. Let $\varepsilon > 0$. Then there exists $C \in \mathcal{C}_k$, $C \subset S$ such that

$$\sup_{(u, v) \in C} \mathcal{A}_a(u, v) \leq \lambda_k(a) + \frac{\varepsilon}{2}. \quad (2.22)$$

Then for any $(u, v) \in C$, if $\|a - b\|_{L^\infty} \leq \delta = \varepsilon/2K$ we get

$$\mathcal{A}_b(u, v) \leq \mathcal{A}_a(u, v) + \frac{\varepsilon}{2} \leq \lambda_k(a) + \varepsilon. \quad (2.23)$$

It follows that

$$\sup_{(u, v) \in C} \mathcal{A}_b(u, v) \leq \lambda_2(a) + \varepsilon \quad (2.24)$$

and we obtain

$$\lambda_k(b) \leq \lambda_k(a) + \varepsilon. \quad (2.25)$$

By reversing the roles of a and b , we get $|\lambda_k(a) - \lambda_k(b)| \leq \varepsilon$. \square

3. Proof of the existence theorem

3.1. A minimax principle. Our main tool for proving [Theorem 1.6](#) will be an abstract minimax principle due to El Amrouss and Moussaoui [\[9\]](#).

THEOREM 3.1. *Let Φ be a C^1 functional on X satisfying condition (C), let Q be a closed connected subset of X such that $\partial Q \cap \partial(-Q) \neq \emptyset$, and let $\beta \in \mathbb{R}$. Assume that*

- (1) *for every $K \in \mathcal{C}_2$ there exists $v_K \in K$ such that $\Phi(v_K) \geq \beta$ and $\Phi(-v_K) \geq \beta$,*
- (2) *$a = \sup_{\partial Q} \Phi < \beta$,*
- (3) *$\sup_Q \Phi < \infty$.*

Then Φ has a critical value $c \geq \beta$ given by

$$c = \inf_{h \in \Gamma} \sup_{x \in Q} \Phi(h(x)), \tag{3.1}$$

where $\Gamma = \{h \in C(X, X) : h(x) = x \text{ for every } x \in \partial Q\}$.

3.2. Compactness conditions

LEMMA 3.2. *Suppose that F satisfies [\(1.6\)](#), [\(1.8\)](#), and [\(1.22\)](#). Then the functional Φ , given by [\(1.2\)](#), satisfies the Cerami condition.*

Proof. In a similar way to [\[9, Lemma 3.1\]](#), we see that the first condition in [Definition 1.1](#) holds.

We will prove that the second condition in [Definition 1.1](#) holds, in the case $L(x, s, t) \rightarrow -\infty$ as $\|(s, t)\| \rightarrow \infty$ (the case $L(x, s, t) \rightarrow +\infty$ is similar). To do that, assume by contradiction that there exists a sequence $(u_n, v_n)_{n \in \mathbb{N}} \subset W$ such that

$$\Phi(u_n, v_n) \rightarrow c, \quad \varepsilon_n = \|\Phi'(u_n, v_n)\| \|(u_n, v_n)\| \rightarrow 0, \quad \|(u_n, v_n)\| \rightarrow \infty. \tag{3.2}$$

Therefore,

$$\left| \frac{1}{p} \langle \Phi_u(u_n, v_n), u_n \rangle + \frac{1}{q} \langle \Phi_v(u_n, v_n), v_n \rangle - \Phi(u_n, v_n) \right| \rightarrow c \tag{3.3}$$

or equivalently

$$\lim_{n \rightarrow \infty} \left| \int_{\Omega} \frac{1}{p} F_u(x, u_n, v_n) u_n + \frac{1}{q} F_v(x, u_n, v_n) v_n - F(x, u_n, v_n) \right| = c. \tag{3.4}$$

We define

$$z_n = \alpha_n^{1/p} u_n, \quad w_n = \alpha_n^{1/q} v_n, \tag{3.5}$$

where

$$\alpha_n = \frac{1}{\mathcal{A}_0(u_n, v_n)} \rightarrow 0 \tag{3.6}$$

with \mathcal{A}_0 given by definition (2.8). We have that $\mathcal{A}_0(z_n, w_n) = 1$ so (z_n, w_n) is bounded in W . After passing to a subsequence, we may assume that

$$\begin{aligned} z_n &\rightharpoonup z && \text{in } W^{1,p}(\Omega), \\ w_n &\rightharpoonup w && \text{in } W^{1,q}(\Omega), \\ z_n &\rightarrow z && \text{in } L^p(\Omega), \text{ a.e. in } \Omega, \\ w_n &\rightarrow w && \text{in } L^q(\Omega), \text{ a.e. in } \Omega. \end{aligned} \quad (3.7)$$

Now we show that $(z, w) \neq (0, 0)$

$$\frac{\Phi(u_n, v_n)}{\mathcal{A}_0(u_n, v_n)} = 1 - \frac{\int_{\Omega} F(x, u_n, v_n)}{\mathcal{A}_0(u_n, v_n)}. \quad (3.8)$$

From (1.22), we get that for any $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that

$$F(x, s, t) \leq (a_2(x) + \varepsilon)G_2(s, t) + C_\varepsilon. \quad (3.9)$$

As a consequence

$$\int_{\Omega} F(x, u_n, v_n) \leq \int_{\Omega} (a_2(x) + \varepsilon)G_2(u_n, v_n) + C_\varepsilon|\Omega|, \quad (3.10)$$

then

$$\frac{\int_{\Omega} F(x, u_n, v_n)}{\mathcal{A}_0(u_n, v_n)} \leq \alpha_n \int_{\Omega} (a_2(x) + \varepsilon)G_2(u_n, v_n) + C_\varepsilon|\Omega|\alpha_n. \quad (3.11)$$

Since

$$\alpha_n \int_{\Omega} (a_2(x) + \varepsilon)G_2(u_n, v_n) = \int_{\Omega} (a_2(x) + \varepsilon)G_2(z_n, w_n) \quad (3.12)$$

in the limit we get

$$0 \geq 1 - \int_{\Omega} (a_2(x) + \varepsilon)G_2(z, w) \quad (3.13)$$

and we conclude that $G_2(z, w) \neq 0$.

Let

$$L(x, s, t) = \frac{1}{p}F_s(x, s, t)s + \frac{1}{q}F_t(x, s, t)t - F(x, s, t). \quad (3.14)$$

By (1.8) (and since L is continuous), $L(x, s, t) \leq -M$. It follows that

$$\int_{\Omega} L(x, u_n, v_n) \leq \int_{\{G_2(z, w) \neq 0\}} L(x, u_n, v_n) + M|\{x : G_2(z(x), w(x)) = 0\}|. \quad (3.15)$$

Note that

$$\alpha_n G_2(u_n, v_n) \rightarrow G_2(z, w). \quad (3.16)$$

So in the set $\{x : G_2(z(x), w(x)) \neq 0\}$, $G_2(u_n, v_n) \rightarrow +\infty$, and then by (1.11), we have that $u_n(x), v_n(x) \rightarrow \infty$. It follows that $L(u_n, v_n) \rightarrow -\infty$ by condition (1.8). Hence the first integral tends to $-\infty$ by Fatou lemma, and we get

$$\lim_{n \rightarrow \infty} \int_{\Omega} L(x, u_n, v_n) = -\infty. \tag{3.17}$$

This contradicts (3.4), and the proof is completed. □

3.3. Geometric conditions. In this section we show that the functional Φ satisfies the geometric conditions of Theorem 3.1.

LEMMA 3.3. *Let F satisfy the assumptions of Theorem 1.6. Then the functional Φ , given by (1.2), satisfies*

- (1) *there exists $(\varphi, \psi) \in W$ such that $\Phi(c^{1/p}\varphi, c^{1/q}\psi) \rightarrow -\infty$ as $c \rightarrow +\infty$;*
- (2) *for every $K \in \mathcal{C}_2$ there exists $(u_K, v_K) \in K$ and $\beta \in \mathbb{R}$ such that $\Phi(u_K, v_K) \geq \beta$ and $\Phi(-u_K, -v_K) \geq \beta$.*

Proof. (1) As $\lambda_1(a, G_1) < 0$, we may choose $\varepsilon > 0$ such that $\lambda_1(a_1 - \varepsilon, G_1) < 0$. Let (φ, ψ) be the first eigenfunction for the problem

$$\begin{aligned} -\Delta_p u - (a_1(x) - \varepsilon)G_{1u}(u, v) &= \lambda|u|^{p-2}u && \text{in } \Omega, \\ -\Delta_q v - (a_1(x) - \varepsilon)G_{1v}(u, v) &= \lambda|v|^{q-2}v && \text{in } \Omega, \\ u = v &= 0 && \text{in } \partial\Omega, \end{aligned} \tag{3.18}$$

normalized by

$$\frac{1}{p} \int_{\Omega} |\varphi|^p + \frac{1}{q} \int_{\Omega} |\psi|^q = 1. \tag{3.19}$$

Then, using (1.13), we get

$$\frac{1}{p} \int_{\Omega} |\nabla \varphi|^p + \frac{1}{q} \int_{\Omega} |\nabla \psi|^q - \int_{\Omega} (a_1(x) - \varepsilon)G_1(u, v) = \lambda_1(a_1 - \varepsilon, G_1). \tag{3.20}$$

By (1.22), we have

$$F(x, s, t) \geq (a_1(x) - \varepsilon)G_1(s, t) - C_{\varepsilon}. \tag{3.21}$$

It follows that

$$\begin{aligned} \Phi(c^{1/p}\varphi, c^{1/q}\psi) &\leq c \left(\frac{1}{p} \int_{\Omega} |\nabla \varphi|^p + \frac{1}{q} \int_{\Omega} |\nabla \psi|^q \right. \\ &\quad \left. - \int_{\Omega} (a_1(x) - \varepsilon)G_1(\varphi, \psi) \right) + C_{\varepsilon}|\Omega| \\ &\leq c\lambda_1(a_1 - \varepsilon, G_1) + C_{\varepsilon}|\Omega|, \end{aligned} \tag{3.22}$$

and so $\Phi(c^{1/p}\varphi, c^{1/q}\psi) \rightarrow -\infty$ as $c \rightarrow +\infty$.

(2) Since $\lambda_2(a_2, G_2) > 0$, we may choose $\varepsilon > 0$ such that $\lambda_2(a_2 + \varepsilon, G_2) > 0$. Given $K \in \mathcal{C}_2$ and this $\varepsilon > 0$, we claim that there exists $(u_K, v_K) \in K$ verifying

$$\begin{aligned} \lambda_2(a_2 + \varepsilon, G_2) & \left(\frac{1}{p} \int_{\Omega} |u_K|^p + \frac{1}{q} \int_{\Omega} |v_K|^q \right) \\ & \leq \frac{1}{p} \int_{\Omega} |\nabla u_K|^p + \frac{1}{q} \int_{\Omega} |\nabla v_K|^q - \int_{\Omega} (a_2(x) + \varepsilon) G_2(u_K, v_K). \end{aligned} \tag{3.23}$$

By (1.22), we have

$$F(x, s, t) \leq (a_2(x) + \varepsilon) G_2(s, t) + C_\varepsilon. \tag{3.24}$$

It follows that

$$\begin{aligned} \Phi(u_K, v_K) & \geq \frac{1}{p} \int_{\Omega} |\nabla u_K|^p + \frac{1}{q} \int_{\Omega} |\nabla v_K|^q \\ & \quad - \int_{\Omega} (a_2(x) + \varepsilon) G_2(u_K, v_K) - C_\varepsilon |\Omega| \\ & \geq \lambda_2(a_2 + \varepsilon, G_2) \left(\frac{1}{p} \int_{\Omega} |u_K|^p + \frac{1}{q} \int_{\Omega} |v_K|^q \right) - C_\varepsilon |\Omega| \\ & \geq -C_\varepsilon |\Omega| = \beta. \end{aligned} \tag{3.25}$$

Similarly,

$$\Phi(-u_K, -v_K) \geq -C_\varepsilon |\Omega| = \beta. \tag{3.26}$$

□

3.4. Proof of Theorem 1.6. We apply Theorem 3.1. We take

$$Q = \{ (|c|^{1/p-1} c\varphi, |c|^{1/q-1} c\psi), -R \leq c \leq R \}, \tag{3.27}$$

where (φ, ψ) is given by Lemma 3.3. Q is closed and compact (it is the image of $[-R, R]$ under a continuous mapping). Also $\partial Q = \partial(-Q) = \{ (\pm R^{1/p} \varphi, \pm R^{1/q} \psi) \} \neq \emptyset$. By Lemma 3.3 if we choose R big enough, we have

$$\sup_{\partial Q} \Phi < \beta. \tag{3.28}$$

Also $\sup_Q \Phi < +\infty$ since Q is compact and Φ is continuous. The functional Φ verifies condition (C) by Lemma 3.2. Then all the conditions of Theorem 3.1 are fulfilled and the proof is completed. □

Acknowledgment

The authors thank specially Prof. J. P. Gossez and the referee for their careful reading of the manuscript and their fruitful suggestions and remarks.

References





- [1] H. Amann, *Lusternik-Schnirelman theory and non-linear eigenvalue problems*, Math. Ann. **199** (1972), 55–72.
- [2] P. Bartolo, V. Benci, and D. Fortunato, *Abstract critical point theorems and applications to some nonlinear problems with “strong” resonance at infinity*, Nonlinear Anal. **7** (1983), no. 9, 981–1012.
- [3] J. M. Bezerra do Ó, *Solutions to perturbed eigenvalue problems of the p -Laplacian in \mathbb{R}^n* , Electron. J. Differential Equations **1997** (1997), 1–15.
- [4] L. Boccardo and D. G. de Figueiredo, *Some remarks on a system of quasilinear elliptic equations*, preprint, 1997, http://www.ime.unicamp.br/rel_pesq/1997/rp51-97.html.
- [5] G. Cerami, *Un criterio di esistenza per i punti critici su varietà illimitate* [An existence criterion for the critical points on unbounded manifolds], Istit. Lombardo Accad. Sci. Lett. Rend. A **112** (1978), no. 2, 332–336 (Italian).
- [6] M. M. Day, *Some more uniformly convex spaces*, Bull. Amer. Math. Soc. **47** (1941), 504–507.
- [7] P. de Nápoli and M. C. Mariani, *Three solutions for quasilinear equations in \mathbb{R}^n near resonance*, Proceedings of the USA-Chile Workshop on Nonlinear Analysis (Viña del Mar-Valparaíso, 2000), Southwest Texas State Univ., Texas, 2001, pp. 131–140.
- [8] G. Dinca, P. Jebelean, and J. Mawhin, *Variational and Topological Methods for Dirichlet Problems with p -Laplacian*, Recherches de mathématique, Inst. de Math. Pure et Appliquée, Univ. Cath. de Louvain, 1998.
- [9] A. R. El Amrouss and M. Moussaoui, *Minimax principles for critical-point theory in applications to quasilinear boundary-value problems*, Electron. J. Differential Equations **2000** (2000), no. 18, 1–9.
- [10] J. Fleckinger, R. F. Manásevich, N. M. Stavrakakis, and F. de Thélin, *Principal eigenvalues for some quasilinear elliptic equations on \mathbf{R}^N* , Adv. Differential Equations **2** (1997), no. 6, 981–1003.
- [11] J.-P. Gossez, *Some remarks on the antimaximum principle*, Rev. Un. Mat. Argentina **41** (1998), no. 1, 79–83.

PABLO L. DE NÁPOLI: DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS Y NATURALES, UNIVERSIDAD DE BUENOS AIRES, CIUDAD UNIVERSITARIA, PABELLÓN I, (1428) BUENOS AIRES, ARGENTINA

E-mail address: pdenapo@dm.uba.ar

M. CRISTINA MARIANI: DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS Y NATURALES, UNIVERSIDAD DE BUENOS AIRES, CIUDAD UNIVERSITARIA, PABELLÓN I, (1428) BUENOS AIRES, ARGENTINA

E-mail address: mcmarian@dm.uba.ar



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

